

Fe_{0.2}Al_{1.8}Zn₁O₄ Composite: An Efficient Catalyst for the Synthesis of 1, 4-Dihydropyridine Derivatives

Vijay Kumar M. Joshi¹, Sunil U. Tekale², Sushama S. Kauthale², Sanjay K. Vyawahare³, Ashok M. Zine⁴, Sunita B. Shinde², K. L. Ameta⁵, Rajendra P. Pawar^{2*}

¹Shri Jagdishprasad Jhabarmal Timrewala University, Rajasthan, India

²Department of Chemistry, Deogiri College, Station Road, Aurangabad (MS) India

³Department of Physics, Deogiri College, Station Road, Aurangabad (MS) India

⁴Vinayakrao Patil Mahavidyalaya, Vijapur (MS) India

⁵Department of Chemistry Mody Institute of Technology and Science Lakshmangarh, Sikar, Rajasthan, India

ABSTRACT:

A simple and rapid protocol has been developed for the synthesis of 1, 4-dihydropyridine derivatives in short reaction time to afford the products in excellent yield. Operational simplicity, clean reaction, high yield, simple work up are the significant advantages of the present protocol.

Keywords: Fe_{0.2}Al_{1.8}Zn₁O₄, Aldehyde, 1, 4-Dihydropyridines, EAA.

1. INTRODUCTION (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

1. INTRODUCTION


1, 4-Dihydropyridine derivatives are the biologically active potent heterocyclic compounds possessing pharmacological and medicinal properties. Dihydropyridine (DHP) skeleton is an integral part of many commercial drugs used in the treatment of hypertension such as mlodipine, nicardipine, nifedipine etc. [1]. 1, 4-Dihydropyridines derivatives are documented in literature for anti-inflammatory [2], antiulcer [3], anticonvulsant [4], antitubercular activity [5], antimicrobial etc. activities [6]. These also act as potential antagonists [7] as well as antidyslipidemic and antioxidant agents [8]. 1, 4-Dihydropyridine and their derivatives are therapeutically significant compounds due to their ability to bind to calcium channels and thus decreasing the flow of the transmembrane calcium current [9]. Thus vast biological activities these compounds have attracted the attention of synthetic organic chemists to synthesize such molecules.

The first classical method for the synthesis of 1,4-DHPs carried out by Hantzsch and Liebig involves the one pot condensation of aldehydes with 1,3-dicarbonyl compounds and ammonia either in acetic acid or in a refluxing alcohol for a prolonged time. Due to certain disadvantages associated with this method; different protocols were developed in due course by the scientific community from academia as well as industry for the synthesis of

* Tel.: +91-(0240) 2334577; fax: +91 (0240) 2334430.

E-mail address: rppawar@yahoo.com.









these heterocyclic compounds which comprises the use of heterogeneous catalysts such as silica sulfonic acid [10], alumina [11], alumina sulfuric acid [12], HY-Zeolite [13] etc. Other Lewis acid catalysts including Mg (ClO₄)₂ [14], phenylboronic acid [15], molecular iodine [16] are also reported for their catalytic efficiencies in the synthesis of these heterocyclic compounds. Furthermore the use of ionic liquid [17], ultrasound [18] or microwave [19] irradiation is also reported. However, research is mainly focused on the modification by optimization of the Hantzsch reaction to minimize reaction time, maximize reaction conversion and offer high purity 1, 4-DHPs. Heterogeneous catalysts are always superior to **homogeneous** ones in many aspects. Currently the study of multicomponent reactions by heterogeneous catalysts has become an active part of ongoing research due to several advantages associated with the heterogeneous catalysts over homogeneous counterparts such as easy separation, recycling ability, minimization of metal traces in products after isolation etc. As a part of our research in the development of new methods for the synthesis of heterocyclic compounds [20], in present work we wish to report the synthesis of 1, 4-dihydropyridine derivatives by the condensation reaction between aldehyde, ethylacetoacetate and ammonium acetate using the composite Fe_{0.2}Al_{1.8}Zn₁O₄ as an efficient catalyst in acetonitrile under reflux condition (**Scheme 1**).



Scheme 1 Synthesis of 1,4-dihydropyridines catalyzed by Fe_{0.2}Al_{1.8}Zn₁O₄ nanocomposite catalyst

Table 1: Synthesis of 1, 4-dihydropyridines using composite- Fe_{0.2}Al_{1.8}Zn₁O₄ catalyst

Entry	R	Product	Time (hr)	Yield (%) [@]	Melting point (°C)
1	Ph		0.75	98	156
2	4-MeOC ₆ H ₄		1.5	95	159
3	4-ClC ₆ H ₄		1.0	91	141-143

4	4-NO ₂ C ₆ H ₄		1.5	89	127
5	4-MeC ₆ H ₄		0.75	89	160-61
6	3-NO ₂ C ₆ H ₄		1.75	80	163
7	n-C ₉ H ₁₉		4.25	78	----
8	2-NO ₂ C ₆ H ₄		3.5	77	119
9	2-furyl		2.0	89	163-65
10	2-MeOC ₆ H ₄		2.5	87	148-149
11	2-FC ₆ H ₄		3.0	92	151

@Reactions were carried on aldehyde (1mmol), EAA (2 mmol) and NH₄OAc (1.5 mmol) using Fe_{0.2}Al_{1.8}Zn₁O₄ (10 mol %) in 2 drops HCl and CH₃CN

2. EXPERIMENTAL DETAILS

The chemicals used were SD fine made and were used without further purification. Melting points of the synthesized compounds were recorded in capillaries open at one end and were uncorrected. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra were recorded on Varian spectrophotometer using CDCl₃ solvent and TMS as the reference.

IR spectra of the samples were recorded on Perkin FTIR spectrometer using KBr discs. The catalyst was synthesized by reported literature method sol-gel method [21-23].

2.1 General procedure for synthesis of 1, 4-dihydropyridine derivatives:

A mixture of aldehyde (1 mmol), ethylacetoacetate (2 mmol), ammonium acetate (1.5 mmol) and the composite $\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ catalyst (10 mol %) in acetonitrile (2 mL) was refluxed for appropriate reaction time as specified in Table 1. The progress of reaction was monitored by Thin Layer Chromatography (25% EA: Hexane). After completion of reaction as indicated by TLC; the reaction mass was diluted with acetonitrile and filtered off to separate the catalyst. The filtrate was concentrated on rotary evaporator and purified by crystallization from ethanol to obtain the pure product.

All aldehydes reacted smoothly to afford the corresponding 1, 4-dihydropyridines in excellent yield (Table 1). The spectral data of representative compounds are mentioned below:

Diethyl 1, 4-dihydro-2, 6-dimethyl-4-phenylpyridine-3, 5-dicarboxylate (Entry 1, Table 1): Yield: 98%; Colorless solid; m.p. 156 °C; IR (KBr) : 500, 620, 1050, 1100, 1180, 1200, 1500, 1667, 1690, 2350, 2867, 3440 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.23 (t, J = 7.2 Hz, 6H), 2.35 (s, 6H), 4.09 (q, 4H), 4.97 (s, 1H), 5.7 (brs, 1H), 7.08 - 7.26 (m, 5H); ^{13}C NMR (CDCl_3): δ 14.28, 19.23, 39.59, 59.56, 103.92, 126.06, 127.89, 144.10, 147.61, 167.54.

Diethyl 1, 4-dihydro-2, 6-dimethyl-4-(4-methoxy-phenyl) pyridine-3, 5-dicarboxylate (Entry 2, Table 1): Yield: 92%; Colorless solid; m.p. 194 °C; IR (KBr) : 500, 700, 1000, 1050, 1200, 1250, 1300, 1500, 1700, 2400, 3300 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.25 (t, J = 7.1 Hz, 6H), 2.38 (s, 6H), 3.94 (s, 3H), 4.05 (q, J = 7.1 Hz, 4H), 4.87 (s, 1H), 5.92 (s, 1H), 6.88 (d, J = 8.7 Hz, 2H), 7.11 (d, J = 8.6 Hz, 2H); ^{13}C NMR (CDCl_3): δ 14.68, 19.55, 39.13, 55.30, 59.85, 104.53, 113.62, 129.19, 140.79, 144.17, 158.30, 167.98.

Dimethyl 1, 4-dihydro-2, 6-dimethyl-4-(2-nitro-phenyl) pyridine-3, 5-dicarboxylate (Entry 8, Table 1): Yield: 75%; colorless solid; m.p. 172-173 °C; IR (KBr): 757, 858, 1020, 1103, 1190, 1217, 1284, 1309, 1352, 1434, 1492, 1529, 1618, 1649, 1689, 3020, 3442 cm^{-1} ; ^1H NMR (CDCl_3): δ 2.34 (s, 6H), 3.79 (s, 6H), 5.73 (s, 1H), 5.87 (s, 1H), 7.21 - 7.30 (m, 1H), 7.41 - 7.54 (m, 2H), 7.66 - 7.70 (m, 1H); ^{13}C NMR (CDCl_3): δ 19.0, 34.29, 52.84, 103.0, 125.65, 126.88, 130.88, 132.70, 142.10, 145.42, 147.54, 167.62.

3. RESULTS AND DISCUSSION

Different aldehydes were used with ethyl acetoacetate and ammonium acetate and $\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ in acetonitrile to illustrate the generality of condensation reaction. The results are summarized in Table 1. The present protocol affords products in good yields. Products were obtained in short reaction time. This method tolerates a variety of functional groups such as methyl, methoxy, hydroxyl, halides etc under the reaction conditions. Importantly, aromatic aldehydes carrying either electron-donating or electron-withdrawing substituents all reacted very well, giving moderate to excellent yields of desired products. The substituent on the aromatic ring of aldehyde did not show much difference in the yield of products.

After completion of reaction, the catalyst was recovered by diluting the reaction mass with hot ethanol followed by filtration and the recovered catalyst was reused for few more cycles. During washing with the solvent, it is clearly evident that there was no leaching of catalyst and was confirmed by performing the reaction with the filtrate. The catalyst reusability study was carried out using the model reaction of benzaldehyde (1 mmol), EAA (2 mmol) and NH_4OAc (1.5 mmol) using $\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ (10 mol %) in 2 drops HCl for the synthesis of diethyl 1, 4-dihydro-2, 6-dimethyl-4-phenylpyridine-3, 5-dicarboxylate (Entry 1, Table 1) whose results are summarized in Table 2.

Table 2: Recycle study of the $\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ composite catalyst.

Run	Fresh	I	II	III
Yield (%)	98	96	95	94

From our investigations, we observed that the catalyst shows excellent to good reactivity with promising yields even for the next three cycles in the same reaction. Since, there was no observable loss in the percentage yield; there is no leaching of the catalyst. Table 3 shows the comparative study of the effect of $\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ composite with some literature protocols on the synthesis of 1,4-dihydropyridines.

Table 3: Comparative study of effect of $\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ composite on the synthesis of 1,4-dihydropyridines.

Entry	Catalyst	Conditions	Yield (%)	Reference
1	Sulfonic acid on silica gel (0.2 g/mmol)	hexane or neat, 60 °C, 4.5-7 h	85-95	[10]
2	Alumina- H_2SO_4 (0.2 g/3mmol)	MeOH, 70 °C, 2-5 h	82-95	[12]
3	HY-zeolite (10 mg/mmol)	EtOH, reflux, 2.5-3.5 h	70-90	[13]
4	I_2 (30 mol %),	EtOH, r.t., 2.5-5 h	80-95	[16]
5	$\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ (10 mol %)	ACN, 2-drops HCl, reflux, 0.75-4.25 h	77-98	[Present method]

4. CONCLUSION

In conclusion, we have developed a simple and efficient method for the synthesis of 1, 4-dihydropyridines by using the composite $\text{Fe}_{0.2}\text{Al}_{1.8}\text{Zn}_1\text{O}_4$ as a solid catalyst, which has many advantages such as ease of preparation, easy handling, reusability and insolubility in most organic solvents. The present protocol affords the 1, 4-dihydropyridines in moderate to excellent yield within short reaction time. We believe this catalyst as a valuable addition to the synthesis of heterocyclic compounds such as 1, 4-dihydropyridine derivatives.

ACKNOWLEDGEMENTS

The authors are greatly thankful to the Principal, Deogiri College, Aurangabad (MS), India for providing the laboratory facilities and encouragement during the work.

REFERENCES

1. a) Buhler FR, Kiowski WJ. Calcium antagonists in hypertension. *J. of Hyperten.* 1987; S3: 3-10, b) Reid JL, Meredith PA, Pasanisi FJ. *Cardio. Pharma.* 1985; S18:18-20.
2. Bahekar S, Shinde D. Synthesis and anti-inflammatory activity of 1, 4-dihydropyridines. *Acta pharmaceutica A.* 2002; 52: 281-287.
3. Subudhi BB, Panda PK. Synthesis antiulcer activity of 1, 4-dihydropyridines and their mannich bases with sulfanilamide. *Ind. J. Chem.* 2009; 48B: 725-728.

- 162 4. Pattan SR, Dighe NS, Musmade DS, Tambe SK, Kale SH, Gaware VM, Chavan PA.
163 Synthesis and evaluation of some new substituted 1, 4-dihydropyridine derivatives
164 and their anticonvulsant activity. *J. Chem. and Pharm. Res.* 2010; 2: 246-252.
- 165 5. a) Fassihi A, Azadpour Z, Delbari N. Synthesis and antitubercular activity of novel 4-
166 substituted imidazolyl-2, 6-dimethyl-N3, N5-bisaryl-1, 4-dihydropyridine-3, 5-
167 dicarboxamides. *Eur. J. Med. Chem.* 2009; 44: 3253–3258. b) Sirisha KG, Achaiah,
168 Reddy VM. Facile synthesis and antibacterial, antitubercular, and anticancer
169 activities of novel 1, 4-dihydropyridines. *Arch. Pharm. Chem. Life Sci.* 2010; 343:
170 342 – 352.
- 171 6. Prakash O, Hussain K, Kumar R, Wadhwa D, Sharma C, Kamal RA. Synthesis and
172 antimicrobial evaluation of new 1, 4-dihydro-4-pyrazolylpyridines and 4-
173 pyrazolylpyridines. *Org. & Med. Chem. Lett.* 2011; 1(5): 1-6.
- 174 7. Sunkel CE, De casa Juana MF, Santos L. 4- Alkyl-1, 4-dihydropyridines as specific
175 PAFacether antagonists. *J. Med. Chem.* 1990; 33: 3205-3210.
- 176 8. Kumar A, Maurya RA, Sharma S, Kumar M, Bhatia G. Synthesis and biological
177 evaluation of N-aryl-1, 4-dihydropyridines as novel antidyslipidemic and antioxidant
178 agents. *Eur. J. Med. Chem.* 2010; 45: 501–509.
- 179 9. Bossert F, Meyer H, Wehinger E. 4-Aryldihydropyridines, a new class of highly
180 active calcium antagonists. *Angew Chem. Int. Ed. Engl* 1981; 20:762.
- 181 10. Gupta R, Gupta R, Paul S, Loupy A. Covalently anchored sulfonic acid on silica gel
182 as an efficient and reusable heterogeneous catalyst for the one-pot synthesis of
183 Hantzsch 1,4-dihydropyridines under solvent-free conditions. *Synthesis* 2007; 2835-
184 2838.
- 185 11. Maiti S, Menendez JC. A mild protocol for the efficient synthesis of 5, 6-
186 unsubstituted 1, 4-dihydropyridines. *Synlett*, 2009; 2249-2252.
- 187 12. Mustafa A, Cuneyt F, Mustafa Z, Mustafa K, Ukislamo G, Hulya D. An efficient one
188 pot synthesis of 1, 4-dihydropyridines using alumina sulfuric acid (ASA) catalyst.
189 *Tur. J. Chem.* 2009; 33: 769-774/
- 190 13. Nikpassand M, Mamaghani M, Tabatabaeian K. An efficient one-pot three-
191 component synthesis of fused 1, 4-dihydropyridines using HY-zeolite. *Molecules*
192 2009; 14:1468-1474.
- 193 14. Bartoli G, Babiuch K, Bosco M, Carlone A, Galzerano P, Melchiorre P, Sambri L.
194 Magnesium perchlorate as efficient lewis acid: a simple and convenient route to 1, 4-
195 dihydropyridines. *Synlett*, 2007; 2897-2901.
- 196 15. Debache A, Boulcina R, Belfaitah A, Rhouati S, Carboni B. One-pot synthesis of 1,
197 4-dihydropyridines via a phenylboronic acid catalyzed Hantzsch three-component
198 reaction. *Synlett*, 2008; 509-512.
- 199 16. Akbari JD, Tala SD, Dhaduk MF, Joshi HS. Molecular iodine-catalyzed one-pot
200 synthesis of some new Hantzsch 1, 4-dihydropyridines at ambient temperature.
201 *Arkivoc*, vol. xii, pp. 126-135, 2008.
- 202 17. Wu XY. Facile and green synthesis of 1, 4-dihydropyridine derivatives in n-butyl
203 pyridinium tetrafluoroborate. *Syn. Comm.* 2012; 42: 454-459.
- 204 18. Wang S-X, Li Z-Y, Zhang J-C, Li J-T. The solvent-free synthesis of 1, 4-
205 dihydropyridines under ultrasound irradiation without catalyst. *Ultrason. Sonochem.*
206 2008; 15: 677–680.
- 207 19. Kotharkar SA, Shinde DB. Microwave assisted synthesis of 1, 4-dihydropyridines.
208 *Ukra Bioorganica Acta*, 2006; 1: 3-5.
- 209 20. Joshi VM, Pawar RP Microwave assisted expeditious synthesis of bioactive
210 polyhydroquinoline derivatives. *Eur Chem Bull.* 2013; 2(9): 679-682.

- 211 21. Stetanescu M, Stoia M, Stefanescu O, Barvinschi P Obtaining of $\text{Ni}_{0.65}\text{Zn}_{0.35}\text{Fe}_2\text{O}_4$
212 nanoparticles at low temperature starting from metallic nitrates and polyols. J. Ther
213 & Anal Calorim, 2010; 99:459-464.
- 214 22. Wang Q, Yang H, Morphology and magnetic properties of $\text{Fe}_x\text{Co}_{1-x}/\text{Co}_y\text{Fe}_{3-y}\text{O}_4$
215 nanocomposites prepared by surfactants-assisted-hydrothermal process. J.
216 Nanopart. Res. 2009; 11:1043-1051.
- 217 23. Baroti MR, Characterization and preparation of nanocrystalline MgCuZn ferrite
218 powders synthesized by sol-gel auto-combustion method. J. Sol-Gel Sci. Technol
219 2009; 52(3):171-178.
220