

Environmental parameters and *Biomphalaria* snail distribution along River Kochi, West Nile region, Uganda

ABSTRACT

Aims: To explore the abundance and distribution of the common fresh water mollusks in River Kochi, with a special focus on *Biomphalaria* species, a vector responsible for transmitting *Schistosoma mansoni* to humans.

Study design:

Place and Duration of Study: This study was conducted between October 2007 and March 2008 as a master's project along Kochi River in Koboko, Yumbe and Moyo in the West Nile region of Uganda.

Methodology: Five sites along the river approximately 20 km apart, were selected and data on snail abundance and various environmental variables thought to be influencing the distribution of snails along this river were collected. These variables included: altitude, season of the year, water flow velocity, water pH, water temperature and concentrations of total dissolved solids in the water.

Results: Findings indicate that numbers of *Biomphalaria* species of snails increased with decreasing altitude (mean numbers 0, 15.33, 19, 50 & 73.33 from highest to lowest altitude points) and no snails of this species were recorded during the wet season. Water flow velocity and pH were the main predictors of the presence of *Biomphalaria* snails (Pearson correlations -0.749 and 0.614 for flow velocity and pH respectively). *Biomphalaria* snail numbers increased when water velocity decreased and the reverse was true for pH.

Conclusion: Altitude influences the distribution *Biomphalaria* snails and hence potential

prevalence of schistosomiasis. Water users of Kochi River should therefore try to minimize contact with water in this river especially during the dry season. Local leaders should lobby to government for alternative sources of water during the dry season.

9

10 *Keywords: Biomphalaria, Schistosomamansoni, River Kochi, West Nile*

11

12 **1. INTRODUCTION**

13

14 Approximately 30 species of *Biomphalaria* are recognized and the genus is widely distributed
15 in South America and on the African continent [1]. *Biomphalaria* is an aquatic snail that acts as
16 a host for a human blood fluke *Schistosomamansoni* that cause the disease intestinal
17 schistosomiasis (bilharzia) in humans [2]. With its many lakes, rivers, streams, swamps and
18 ponds, Uganda has a diverse fresh water environment that offers numerous and suitable
19 habitats for the *Biomphalaria* species. Currently two species of *Biomphalaria* namely: *B.*
20 *stanleyi* and *B. sudanica* (hereby known as *Biomphalaria*) are the most common in the west Nile
21 region of Uganda [3].

22 Whereas Nelson [4] in 1950s pioneered research works on schistosomiasis in West Nile region
23 approaching the infections from ecological and geographical points of view and from both
24 human populations and snail vectors in water bodies, research that followed his works mainly
25 concentrated either in human communities that live close to the shores of lake Albert/ Albert
26 Nile [5, 6, 7, 8, 9 & 10] or from hospital records [5, 11], with the exception of findings of Kazibwe
27 et al [3]. Contrary to the aforementioned studies, Kazibwe [3] looked at the effect of
28 environmental factors on the distribution of *Biomphalaria* in Lake Albert, Western Uganda.
29 Findings from this study revealed that climatic conditions primarily air temperature, rainfall, lake
30 depth level, water temperature; water conductivity and water pH influence the distribution and
31 abundance of snails in Lake Albert. Similarly Appleton [11] and Thieltges et al. [13] showed that

32 climatic conditions primarily rainfall and temperature influence the distribution and abundance
33 of snails because they have an effect on their breeding and the rate of schistosomal
34 development.

35 It is clearly evident from the aforementioned studies that the studies on *Biomphalaria* snail
36 species ecology was restricted to large water bodies, with little or no attention given to small
37 ones, which are also a source of water and fish for the local communities. However, it is
38 important to mention that Odongo-Aginya and others [14] conducted a research on urban
39 *Schistosomamansoninear* Enyau River in Arua town, a small river in the highland areas of the
40 region further away from the Nile River but his focus was on infections in humans. In addition,
41 human populations in the township comprise of people from different origins and locations, and
42 may therefore not have given dependable results since all the *S. mansoni* cases registered may
43 not have been contracted from Enyau River. We therefore strongly believe that conducting
44 research on vector dynamics in smaller water bodies will result into better understanding of the
45 disease prevalence and its distribution in the region considering that a lower number of snails
46 mean a lower number of cercariae and therefore a diminished risk of infection. Studies on
47 diseases vectors are very important for evidence based mitigation and control measure. The
48 main objective of the study was therefore to investigate effects of altitude, season and water
49 environmental variables on the distribution of *Biomphalaria* snail species along the Kochi River,
50 West Nile region. Findings from this study will generate some current information on the
51 distribution of the disease vector in the different infection zones and altitudes in the study area.

52


53 **2. MATERIAL AND METHODS**

54 **2.1 Study sites**

55 The study took place in Kochi River located in Koboko, Yumbe and Moyo districts of West Nile
56 region. Generally, the study area was divided into three altitude zones i.e. Koboko with altitude
57 of above 1000 m; Yumbe (two sites: Yumbe 1 and Yumbe 2) with altitude range of 700 m to

1000 m and Moyo (two sites: Moyo 1 and Moyo 2) with altitude range of 600 m to 700 m above sea level. Kochi River has its origin in Koboko district near Uganda-DR Congo boarder at an altitude of above 1000 m where it starts as a small stream and gradually widens downstream as it passes through Yumbe district and finally joins the Albert Nile in Moyo at an altitude of about 600 m above sea level (Fig 1). This river stretches all the schistosomiasis infection belts of the region that Nelson [4] had established in 1958. The rainfall pattern in this region is bimodal peaking in late March to May (about 900 mm), and August to December (above 900 mm) each year. The rest of the months experience dry spells with sporadic rainfall which fluctuate the water levels of the rivers and its stream tributaries where some seasonal ones dry up completely.

Fig.1. Map showing the study sites



71

72 2.2 Estimating snail abundance and water parameters

73 *Biomphalaria* snail abundance was estimated from well-defined areas along the river. These
 74 areas measured 30 m along the bank and 3 m into the main body of the water. The corners of
 75 these rectangular sampling areas were marked by pegs so that successive samplings could be
 76 performed across the same area. These areas were searched for a period of 30mins and all
 77 snails found floating or attached to vegetation were collected using a scooping net with a long
 78 handle and placed on white plastic trays in order to be able to rapidly identify the different
 79 species based on the standard field identification key guide of the Danish bilharziasis
 80 laboratory [15]. Each site was visited weekly and snail samples collected over a period of six
 81 months between October 2007 and March 2008. Three of the months (October, November and

December) experienced heavy rains of above 900 mm and have been recorded as wet, whilst the other three months (January, February and March) experienced little or no rains and have been recorded as dry. Although our main focus was *Biomphalaria*, snail types like *Lymnae*, *Bulinus* and *Pila* species were collected because they coexist with the *Biomphalaria* snails and are intermediate hosts to other human and animal diseases. Snails were collected from 5 altitude belts spread across the study area at intervals of about 20 km apart.

We took measurements on water flow velocity, water pH, water temperature and concentration of total dissolved solids (TDS) in the water shortly before collecting the snail samples. Water flow velocity was obtained by sprinkling methyl orange dye from the upstream mark of the sampling area and recording the time taken for the dye colour to cover the 30 m distance to the downstream mark and velocities computed. Values for pH and temperature were obtained by using a pH meter integrated with a temperature probe (Model 3150/REV A/04-95). TDS concentration was determined using a conductivity meter (Model 4200/REV A/05-95).

2.3 Data analysis

Data were analysed using Genstat version 3. Firstly, we made a descriptive summary of the abundance of all snails in total, and then secondly we singled out *Biomphalaria* species and explored how its distribution is affected by the environmental factors considered in this study. Pearson-r Correlation Coefficient tests were done to establish associations between the different variables. Environmental variables that had strong associations with the abundance of *Biomphalaria* species were then used in Simple Regression models.


102

3. RESULTS AND DISCUSSION

Higher numbers of *Biomphalaria* and *Pila* were recorded in Yumbe and Moyo while that of *Lymnae* and *Bulinus* snail species were registered in Koboko (Fig 2). Considering that these locations are positioned at different altitudinal zones, these results already indicate an effect of

107 altitude on the distribution of these snail species. The results further show that no *Biomphalaria*
108 species of snails were recorded during the wet season, and the number of snails increased
109 with decreasing altitude during the dry season from none recorded at an altitude of 1189 m to a
110 mean of 62 snails recorded per month at an altitude of 638 m or 639 m (Table 1).

111 **Fig. 2. Abundance and distribution of common snail species at sites along River Kochi**



115 **Table 1: Altitudinal and seasonal variation in the number of *Biomphalaria* species**

Season	Mean	
	Dry	Wet
Altitude		
638.0	73.33	0.00
639.0	50.00	0.00
898.0	19.00	0.00
933.0	15.33	0.00
1189.0	0.00	0.00

116

117 A correlation analysis indicated that there was an association between the number of
 118 *Biomphalaria* snails and water flow velocity (-0.749) and between *Biomphalaria* snails and pH
 119 (0.614) (Table 2). Water flow velocity and pH were in turn highly correlated (-0.899) indicating a
 120 strong association between them (Table 2 and Fig 3). Weak associations existed between
 121 number of *Biomphalaria* snails and Temperature and Total dissolved solids (Table 2). In view
 122 of the result shown in table 1 that show that snail incidence varied with altitude and season we
 123 found it important to summarize mean values for water flow velocity and pH levels in the same
 124 way. Results for water flow velocity show that snails were found only in the dry season at the
 125 five sites where the velocity was in a range of 0.19 to 0.31 m/s. No snails were recorded at
 126 Koboko, which is the site at the highest altitude. The water velocity in the dry season was 0.48
 127 m/s, which is at the lower end of the range of values shown for the wet season and above the
 128 value of 0.4 m/s (the upper limit for presence of snails shown in Table 1). A similar trend can
 129 be seen for pH. Snails were found only when pH values were 7.1 or above (Table 3).

130 **Table 2: A correlation matrix between numbers of *Biomphalaria* and environmental**
131 **variables**


132

133

*** Correlation matrix ***					
NBiom	1.000				
TDS	-0.264	1.000			
Temp	0.012	0.746	1.000		
Velocity	-0.749	0.182	0.027	1.000	
pH	0.614	0.099	0.084	-0.899	1.000
	NBiom	TDS	Temp	Velocity	pH

*NBiom = number of *Biomphalaria* snails; TDS = Total Dissolved solids

134 **Fig. 3: Relationship between water flow velocity and water pH**



135

136

137 **Table 3:Altitudinal and seasonal variation in Water flow velocity and water pH**

Mean (Water flow velocity)			Mean (water pH)		
Season	Dry	Wet	Season	Wet	Wet
Altitude			Altitude		
638.0	0.1867	0.4733	638.0	7.200	6.947
639.0	0.2267	0.5400	639.0	7.187	6.837
898.0	0.2917	0.5683	898.0	7.183	6.893
933.0	0.3083	0.5600	933.0	7.080	6.900
1189.0	0.4450	0.7283	1189.0	6.813	6.533


138 * Altitude decreases from Koboko to Moyo

139 Considering that pH and flow velocity were strongly correlated with each other, we decided to
 140 use Simple linear regression models for each variable to predict *Biomphalaria* snail
 141 abundance. The results showed highly significant relationships between water pH and flow
 142 velocity ($P < .001$) and that pH accounted for 38% of the variation while water flow velocity
 143 accounted for 56% of the variation in *Biomphalaria* numbers. There was a positive association
 144 between *Biomphalaria* snails and pH when pH is 7.1 or above and no snails were found when
 145 pH was below 7.1 (Fig 4). In addition, the number of *Biomphalaria* snails increased when
 146 water flow velocity decreased below about 0.4 m/s and no snails were found when velocity was
 147 0.4 m/s or above (Fig 5).

148

149


150 **Fig. 4: Relationship between numbers of *Biomphalaria* snails and water pH**



151

152

153 **Fig. 5: Relationship between numbers of *Biomphalaria* snails and water flow velocity**



154

155

156 **DISCUSSION**

157 *Bulinus* and *Lymnae* snail species were distributed throughout the river length. On the other
 158 hand, *Biomphalaria* and *Pila* species were not however found at all in some of the sites
 159 especially those towards the source of the river in Koboko above 1000 m. They were only
 160 found in sites towards the Nile in Yumbe and Moyo at fairly lower altitudes. The presence of
 161 *Lymnae* species in the river poses a threat of *Fasciola hepatica* (liver fluke) transmissions in
 162 domestic animals that graze along the river in case some of the animals happen to be infected
 163 with the disease. In the same token *Schistosoma haematobium* (urinary schistosomiasis) could
 164 easily spread in this area as their intermediate snail vectors (*Bulinus* species) are readily
 165 available in the river and the fact that major roads that cross into South Sudan where *S.*
 166 *haematobium* exists also cross this river. With high human mobility across these two countries,
 167 existence of *S. haematobium* in the area if not the region is highly likely and therefore needs
 168 investigation.

169 *Biomphalaria* numbers in the dry season were associated with water flow velocity and pH
 170 levels. A possible explanation for the association is that the river becomes wider and so the
 171 flow speed of the water reduces further downstream. Stable water conditions downstream
 172 would be particularly prevalent during the dry season. Such conditions would enable the snails
 173 to anchor more easily on the water vegetation. Also, as the debris carried down the river settles
 174 and rots down, so the pH of the water gradually increases. This would explain why higher
 175 numbers of *Biomphalaria* species of snails are associated with lower water flow rates and
 176 higher pH levels (See also [3]). It is however important to note that not all the variation in
 177 *Biomphalaria* species numbers was explained by water velocity and pH. It is possible that this
 178 unexplained variation could be due to other factors such as amount of vegetation and snail
 179 prey present at the study sites, which were out of the scope of this study.

180 Numbers of *Biomphalaria* species of snails increased with decreasing altitude and no snails of
 181 this species were found during the wet season. The complete absence of snails in the wet
 182 season is very difficult to explain. However, we think that this is attributed to the fact that during
 183 this time of the year, the water flow velocity in river Kochi was very high thereby drifting the
 184 snails away. In addition the water table was also very high therefore submerging the vegetation
 185 onto which the snails attach. This seems to suggest that there is need for lowering the sweep
 186 net further deeper into the water to search the snails.

187 Furthermore Kabetereine, [16] recorded bigger numbers of *Biomphalaria stanleyi* in shallow
 188 waters along Lake Albert during dry season and this was mainly attributed to the effects of light
 189 penetration on the growth of *Vallisneria* weeds which serve as food for the snails in the river.
 190 Fewer numbers of snails were recorded when lake levels increased and light penetration to
 191 support growth of the weeds reduced. In addition, warmer and wetter conditions encouraged
 192 snails to lay more eggs thereby increasing the densities of young snails several weeks later. In
 193 this current study we attribute the big numbers of *Biomphalaria* snails collected during dry
 194 season to the preceding wetter and warmer months of wet season that resulted in mass egg
 195 laying and subsequent development of these eggs into the large number of adult snails
 196 registered later in the drier months of dry season.

197 There was no significant effect of temperature on the distribution and abundance of
 198 *Biomphalaria* snail species along Kochi river. This finding is rather contrary to studies
 199 conducted elsewhere in the world [16, 11, 18, 19 & 20]. These findings could be attributed to
 200 the fact that there was/is no severe fluctuations in temperature as compared to the extreme
 201 cold and hot temperatures experienced in the study sites considered in the above mentioned
 202 studies.

203

204 4. CONCLUSION

205 In conclusion, we want to acknowledge that although this study was limited in time scope
 206 compared to earlier studies, our findings are consistent as they seem to indicate that despite
 207 national schistosomiasis control efforts, *Biomphalaria* snail species are still present within the
 208 West Nile region. The continued presence of these snail species and other associated snail
 209 species in smaller water bodies could thwart the efforts to contain schistosomiasis in this
 210 region and pose an unforeseen threat to a number of snail transmitted diseases to humans,
 211 and domestic animals in communities along the rivers most especially in the lower altitudes.
 212 We recommend regular community sensitisation by the Ministry of health about the risks of
 213 getting into contact with the river water during the dry season and that the concerned local
 214 governments lobby to government to provide alternative sources of water e.g. boreholes that
 215 can be used in the dry season to minimise peoples' contact with the river water. Mass control
 216 interventions by the government to the schistosomiasis pandemic in this region following quick
 217 diseases surveys in human communities will yield little results if no focus is paid to the water
 218 sources where the disease is contracted. Further research may be directed towards scaling up
 219 the study along other rivers in the west Nile region and also incorporating other variables like
 220 the amount of vegetation and snail prey present at the sites.

221

222

223 REFERENCES

- 224 1. Brown, D. S. Freshwater snails of Africa and their medical importance. Taylor &
 225 Francis, London; 1994.
- 226 2. Jordan, P. & G. Webbe. Epidemiology. In: Human schistosomiasis(Eds P Jordan, G.
 227 Webbe, & R. F. Sturrock). Commonwealth Agricultural Bureau International,
 228 Wallingford;1993:87-158.
- 229 3. Kazibwe, F., B. Makanga, C. Rubaire-Akiiki, J. Ouma, C. Kariuki, N. B. Kabatereine, M.
 230 Booth, B. J. Vennervald, R. F. Sturrock, J. R. Stothard. Ecology of *Biomphalaria*

- 231 (Gastropoda: Planorbidae) in Lake Albert, Western Uganda: Snail distribution,
232 infections with schistosomes and temporal associations with environmental dynamics.
233 *Hydrobiologia*. 2006;568(1):433-444.
- 234 4. Nelson, G.S. Schistosomamansoni infection in West Nile District of Uganda. Part II. The
235 distribution of *S. mansoni* with a note on the probable vectors. *East African Medical*
236 *Journal*. 1958;35:335-344.
- 237 5. Ongom V.L. & Bradley D.J. 1972. The epidemiology and consequences of
238 Schistosomamansoni infection in West Nile. Part I. Field studies of a community at
239 Panyagoro. *Transaction of Royal Society of Tropical Medicine and Hygiene*.
240 1972;66:835-851
- 241 6. Ongom, V.L. *et al.* The epidemiology and Consequences of
242 Schistosomamansoni Infection in West Nile, Uganda. II. Hospital investigation of a
243 sample from the Panyagoro Community. *Transaction of Royal Society of Tropical*
244 *Medicine and Hygiene*. 1972;66:851-863.
- 245 7. Bukenya, G. & S. Andama. Circumstantial epidemiology of Schistosomamansoni in the
246 West Nile District of Uganda: Results of a cross-sectional study in the Rhino Camp
247 Area. *Journal of Tropical Medicine and Hygiene*. 1986;89:243-248.
- 248 8. Kabatereine N.B., C. Ariho, & N.O. Christensen. Schistosomamansoni in Pakwach, Nebbi
249 District, Uganda, 40 years after Nelson. *Tropical Medicine and Parasitology*.
250 1992;43:162-166.
- 251 9. Laroni-Lakwo T, Odongo-Aginya EI, Schweigmann U, Schickerling S, Linder D,
252 Doehring-Schwerdtfeger E. Transmission of *Schistosoma mansoni* in Rhino Camp,
253 Uganda. *E. Afr. Med. J.* 1994;71:165-166.
- 254 10. Kabatereine, N.B., F. Kazibwe, & J. Kemijumbi. Epidemiology of schistosomiasis in
255 Kampala, Uganda. *East African Medical Journal*. 1996;73:795-800.
- 256 11. Williams, E.H., R.J. Hayes, & P.G. Smith. Admissions to rural hospital in the West Nile
257 District of Uganda over a 27 year period. *Journal of Tropical Medicine and Hygiene*.

- 1986;89:193-211.
12. Appleton, C.C. Review of literature on abiotic factors influencing the distribution and life cycle of bilharziasis intermediate snail. *Malacological Review*. 1978;11:1-25.
13. Thieltges, D., K.Jensen & R. Poulin. 2008. The role of biotic factors in the transmission of free living endohelminth stages. *Parasitology*. 2008;135:407-426
14. Odongo-Aginya, E.I. Lakwo TL, Schweigmann U, Schickerling S, Lindner D, Mueller A, Doehring-Schwerdtfeger E. Urban Schistosomamansoninear Enyau River in Arua Town, Uganda. *East African Medical Journal*. 1994;71:604-606.
15. Danish Bilharziasis Laboratory. A Field Guide to African Freshwater Snails, (2nd edn.) East African species. World Health Organization Collaborating Centre, Copenhagen. 1987:29-32
16. Kabatereine, N.B., F.M. Flemming, U. Nyandindi, J.C.L. Mwanza, and B. Lynsey. The control of schistosomiasis and soil transmitted helminths in East Africa. *Trends in Parasitology*. 2006;7:332-339.
17. Shati, A. A. Factors affecting the prevalence of human schistosomiasis in Aseer regions, Saudi Arabia. *Journal of Biological Sciences*. 2009;9:815-819.
18. Martens, W.J.M. Modelling the effect of global warming on schistosomiasis on the prevalence of schistosomiasis. *GLOBO Report Service*. 1995;10:1-31.
19. Gryseels, B., K. Polman., J. Clerinx & L. Kestens. Human schistosomiasis. *Lancet*. 2006;368:1106-1118.
20. Brooker, S. Spatial epidemiology of human schistosomiasis in Africa risk models transmission dynamics and control. *Transaction of Royal Society of Tropical Medicine and Hygiene*. 2007;101:1-8.