

# **A Retrospective Study of *Cinara cupressivora* Damage on *Cupressus lusitanica* Clonal Seed Orchard in Malawi between 1997 and 2003**

## **ABSTRACT**

A retrospective study covering a period of seven years (1997-2003) was conducted to assess the extent of damage of *Cinara cupressivora* on *Cupressus lusitanica* Orchard and determine any correlation with climatic factors in Dedza, Malawi. Furthermore, the study was also aimed at determining the efficacy of *Pauesia juniperorum* as a biological control. The data was extracted from monthly reports of the Forestry Research Institute of Malawi (Centre), where a total of 319 trees were assessed. A statistically significant ( $X^2=13.97$ ,  $P<0.001$ ) association was found between the damage of the trees and increased number of *C. cupressivora*. The damage was found to be eighteen times (Odds Ratio=18.1) more likely to occur on trees attacked by *C. cupressivora* than those not attacked. The hot-dry season was found to be significantly ( $X^2=8.6$ ,  $P<0.001$ ) associated with the increased number of *C. cupressivora*, and the damage was found to be three times (Odds Ratio=3.4) more likely to occur in this season compared to cold-wet and warm-wet seasons. Consequently, the results further show a significant ( $X^2=26.37$ ,  $P<0.001$ ) association between the survival of trees and the presence of *P. juniperorum*. The trees attacked by *C. cupressivora* were found to be twenty-nine times (Odds Ratio=29.1) more likely to survive with the presence of *P. juniperorum* than those where the parasitic wasp were absent. It is, therefore, recommended that classical biological control is the most suitable and permanent solution for control. Hence, *P. juniperorum* is a potential agent for the biological control of *C. cupressivora*.

**Key words:** season; biological control; *Pauesia juniperorum*; odds ratio; survive.

## **1. INTRODUCTION**

The cypress aphid, *Cinara cupressivora* is a significant pest of Cupressaceae species and has caused serious damage to naturally regenerating and planted forests in Africa, Europe, Latin America and the Caribbean and the Near East [1, 2]. This indicates that the pest has great dispersal ability and adaptability to different climates and hosts [3]. It was first reported in Africa, from northern Malawi in 1986 [4, 5]. It was

then rapidly spread in East and Central African countries, including Burundi, Uganda, Kenya and Tanzania [5, 6]. While it is not a pest in its native Europe and North America, it has rapidly established itself in Africa as a devastating cypress pest [6-8]. It has been estimated that the aphids cause an annual loss of growth increment worth \$US13.5 [9], and has killed \$US 41 million worth of trees in Africa [10] and over \$US 2.4 million of losses in Malawi alone [4].

*Cinara cupressivora* has caused extensive dieback and mortality of *Cupressus lusitanica* Miller. *Cupressus lusitanica* is a tree native to Guatemala and Mexico [11-13]. It is widely planted in Southern, Eastern and Central African region in farmlands as hedges in urban and rural areas and it is grown in gazetted forests for production of timber [6]. *Cinara cupressivora* adults and nymphs suck the plant sap on terminal growth of trees, which causes retarded growth and desiccation of the stems. This may result in a progressive die bark on heavily infested trees. In addition, the aphid feeding is accompanied by copious production of honey dew which encourages the growth of sooty mould. The mould causes foliage discolouration and interferes with photosynthesis and gas exchange [1, 2, 12-16]. The presence of ants and the occurrence of lady bird beetle, which tends to feed on the honey dew is often an indicator of aphid infestation [12, 17, 18].

In view of this, between 1991 and 1993 surveys in Europe and North America were conducted by the International Institute and Biological Control (IIBC) to identify natural control agents to use in Africa. A parasitic wasp *Pauesia juniperorum*, native to Western Europe was finally selected as a potential control agent [19]. *Pauesia juniperorum* only attacks aphids belonging to the sub-family Lachnidae which are found exclusively on conifers. An adult *Pauesia* is about 10 mm long, almost twice the length of its host. It has a black head, brown-black thorax, yellow legs and a yellowish abdomen which becomes darker in older insects. After emerging from its cocoon, the adult female, seeks and lays eggs, inside cypress aphids. Within the parasitized aphid the egg hatches into a larva which feeds and develops rapidly through several instar stages. The parasitoid pupa remains inside the host cuticle which stretches, darkens and becomes hard and gets attached to the twigs. At this stage, the parasitized aphid is referred to as a mummy. The pupal period lasts for about six days. After full development, the adult parasitoid cuts an incomplete circular hole at the posterior end of the mummy to emerge. The total development time from egg to adult is about 14 days. Adults can live up to 7 days [19-24].

In mid and late 1990's consignments of the *P. juniperorum* were released in many *C. lusitanica* hedges and plantations in Malawi. The effectiveness of the parasitoid was observed in the field and a decline in the severity of damage has been observed [21]. However, the quantification of the efficacy of the parasitoid is scarce and the information on the damage caused by the *C. cupressivora* and any correlation with climatic factors is limited. Therefore, the objective of this study was to quantify the damage of *C. cupressivora* on *C. lusitanica* Orchard and determine any correlation with climatic factors in Dedza, Malawi. The study was also aimed at quantifying the efficacy of *P. juniperorum* as a biological control.

## 2. MATERIALS AND METHODS

### 2.1 Study site

The study was conducted in Malawi located in Southern Africa in the tropical savannah region in *C. lusitanica* clonal seed orchard (14° 19' S; 34° 15' E and about 1600 m above sea level) at Chongoni Forest Plantation, Dedza. Dedza receives 1200 mm to 1800 mm rainfall per annum, with annual temperature ranging from 7°C to 25°C. It is situated about 85 km southeast of the capital Lilongwe. The clonal seed orchard number TB71/2/5 was established in 1971 with twenty one clones from Kenya and each clone had twenty-one laments (*number of trees*). The clones were planted in a completely random design in three replicates at a spacing of 3m x 3m.

### 2.2 Data collection

Records of Biological Control Programme carried out by Forestry Research Institute of Malawi (FRIM) and International Institute and Biological Control for the period starting from January 1997 to September 2003 were used for this study. The records were collected monthly during the stated period and comprised of tree number, clone number, branch number, number of aphids found in the branch, number of *P. juniperorum* or mummy present in the branch, and damage category of the branch. Damage was categorized into two status; live and partially dead. The records comprised of a total of 319 trees.

### 2.3 Statistical analysis

As data covered a long period of time, care was taken to verify duplication before electronic entry. The collected data was captured onto Microsoft Excel software as a Dbase relational database for editing,

validation, verification, and generation of descriptive statistics. Seasonal analysis was examined by dividing the year into three seasons; hot-dry (September-October), warm-wet (November-April), and cold-wet (May-August) seasons. An association between damage and presence of aphids; damage and seasons; and survival of tree or branches and presence of *P. juniperorum* were evaluated by calculating the chi-square ( $X^2$ ) test for association and the Mantel-Haenszel common odds ratio (OR) at a confidence level of 95% using SPSS Version 17.

### 3. RESULTS

#### 3.1 Association between *Cinara cupressivora* and damage on *Cupressus lusitanica*

Summary of the results on association between *C. cupressivora* and damage on *C. lusitanica* are presented in Table 1. The results shows that there were statistically significant ( $X^2=13.97$ ,  $P<0.001$ ) association between the damage of the trees and increased number of *C. cupressivora*. The damage was found to be eighteen times (Odds Ratio=18.1) more likely to occur on trees attacked by *C. cupressivora* than those not attacked.

**Table 1** Association between presence of *Cinara cupressivora* and damage caused on *Cupressus lusitanica*

		Presence of <i>Cinara cupressivora</i> (%)		Total (%)
		Yes	No	
Damage	Partially dead	76.8	16.4	93.2
	Live	1.4	5.4	6.8
	Total	78.2	21.8	100

$X^2=13.97$ ,  $df=1$ ,  $P<0.001$ , odds ratio=18.1

#### 3.2 Season and presence of *Cinara cupressivora*

The hot-dry season was found to be significantly ( $X^2=8.6$ ,  $P<0.001$ ) associated with the increased number of *C. cupressivora*, and the damage was found to be three times (Odds Ratio=3.4) more likely to occur in this season compared to cold-wet and warm-wet seasons (Table 2).

**Table 2** Mean number of *Cinara cupressivora* present per tree branch during different seasons

Season	Mean number of <i>Cinara cupressivora</i> per tree branch $\pm$ s.e.	Percentage
Hot-dry	5.8 $\pm$ 0.4 <sup>a</sup>	50.4
Warm-wet	1.8 $\pm$ 0.3 <sup>c</sup>	15.7
Cold-wet	3.9 $\pm$ 0.4 <sup>b</sup>	33.9

Note: <sup>a,b,c</sup>Mean with different subscript within a column significantly differ ( $P<0.001$ )

### 3.3 Efficacy of *Pauesia juniperorum* as a biological control on survival of *Cupressus lusitanica* attacked by *Cinara cupressivora*

There were significant ( $X^2=26.37$ ,  $P<0.001$ ) association between the survival of trees/branches and the presence of *P. juniperorum* (Table 3). The trees/branches attacked by *C. cupressivora* were found to be twenty-nine times (Odds Ratio=29.1) more likely to survive with the presence of *P. juniperorum* than those where the parasitic wasp were absent.

**Table 3** Association between presence of *Pauesia juniperorum* and survival of *Cupressus lusitanica* attacked by *Cinara cupressivora*

		Presence of <i>Pauesia juniperorum</i> (%)		Total (%)
		Yes	No	
Damage	Live	85.2	7.4	92.6
	Partially dead	2.1	5.3	7.4
	Total	87.3	12.7	100

$X^2=26.37$ ,  $df=1$ ,  $P<0.001$ , odds ratio=29.1

## 4. DISCUSSION

Even though 16.4% of the partially dead trees and branches showed no presence of *C. cupressivora*, indicating that they may be other factors causing the death [13], the finding of this study indicates *C. cupressivora* as a major significant risk factor to the damage of *C. lusitanica*. This association is also consistent with the findings in Kenya [13, 23]. According to [13], the cypress aphid exploits a wide range of feeding sites varying from green branches to woody stems. Damage mainly occurs by sap feeding, as a result the foliage turns yellow and brown. The saliva produced is phototoxic and leads to necrosis in the phloem hence withering of twigs. Feeding retards new growth and causes desiccation of the stems with a progressive dieback of heavily infested trees [10, 12, 13], and this would explain the significant association found in this study. The findings by [25] also supports the present results. Owuor [25] reported that the overall effect ranges from partial damage to ultimate death of the entire tree and this depends on the infestation severity and in case of severe infestation the death of mature trees can occur within three months.

The study revealed that the peak season for numerous cypress aphids is hot-dry season (September-October) and the population starts build up in cold-wet season (May to August). However, in warm-wet season (November-April; rain season) the population of cypress aphids decline. The significant association of the high occurrence of the cypress aphids in hot-dry season found in this study is consistent with the

findings of [26] in Mauritius. According to [26] the cypress aphids have been reported to increase in number during hot-dry season and decrease in rainy season. For instance, no mummies were observed in a survey in Mauritius during rainy season since the population of the cypress aphid was low [26]. Hot-dry conditions seems to favour both cypress aphid activity and survival [27-32]. A similar pattern was observed in Columbia [33] and Uganda [34]. Hence, cypress aphid populations are strongly influenced by weather conditions, and they are present throughout the year [26, 28, 31]. Findings of a significantly higher number of cypress aphid in hot-dry season in Kenya [13, 23, 35] also support the results of the present study. Mwangi [35] state that, "Population density of cypress aphid is highest during the hot, dry season and lowest during the season of heavy rains. The decline in population density results in some recovery of damaged trees."

The study has revealed that the trees or branches of *C. lusitanica* attacked by *C. cupressivora* were found to be twenty-nine times more likely to survive with the presence of *P. juniperorum* than those where the parasitic wasp were absent. The present results are in agreement with the results in literature [21-23, 26, 27, 35-38]. According to [21], there is high efficacy of *P. juniperorum* as a biological control on survival of *C. lusitanica* attacked by *C. cupressivora* because the female wasp lays its eggs in live cypress aphids and the eggs hatch into larvae that feed on the pest's internal organs, eventually killing it. However, Kairo and Murphy [27] reported that the effectiveness is greater in older hosts, but in young hosts the cypress aphid escape parasitism. Hosts parasitized before they were 9-days-old became mummies without reproducing. Those that were 9 days or older achieved part of their reproductive capacity before dying. Thus, the impact of *P. juniperorum* is markedly reduced when attacking older hosts.

The best effective management option of the aphid is integrated pest management [23, 36]. However, basing on the finding of this study and also, since cypress aphid is an exotic pest, it is, therefore, recommended that classical biological control is the most suitable and permanent solution for control. Hence, *P. juniperorum* is a potential agent for the biological control of *C. cupressivora*.

## 5. CONCLUSION

The present study has revealed that there was statistically significant association between the damage of the trees and increased number of *C. cupressivora*. The damage was eighteen times more likely to occur

on trees attacked by *C. cupressivora* than those not attacked. The hot-dry season was significantly associated with the increased number of *C. cupressivora*, and the damage was three times more likely to occur in this season compared to cold-wet and warm-wet seasons. Consequently, the results further shows a significant association between the survival of trees and the presence of *P. juniperorum*. The trees attacked by *C. cupressivora* were twenty-nine times more likely to survive with the presence of *P. juniperorum* than those where the parasitic wasp was absent. It is, therefore, recommended that classical biological control is the most suitable and permanent solution for control. Hence, *P. juniperorum* is a potential agent for the biological control of *C. cupressivora*.

## REFERENCES

1. F.A.O. Forest Pest Species Profile: *Cinara cupressivora* Watson and Voegtlin 1999. FAO, Rome. 2007. (Available at <http://www.fao.org/forestry/13557-011835cf4f2791802c40723e174bd99d.pdf>)
2. Durak R, Borowiak-Sobkowiak B, Socha M. Bionomy and Ecology of *Cinara cupressi* (Bluckton) (Hemiptera: Aphididae). Polish Journal of Entomology. 2007; 76:107 – 113.
3. Ruiz C, Lanfranco D, Carrillo R, Parra L. Morphometric Variation in the Cypress Aphid *Cinara cupressi* (Bluckton) (Hemiptera: Aphididae) Associated to Urban Trees. Neotropical Entomology. 2014; 43:245 – 251.
4. Chilima CZ. The status and development of conifer aphid damage in Malawi. In: *Exotic Aphid Pests of Conifers. A crisis in African Forestry*. Proceedings of a Workshop at Muguga, Kenya, June 3 – 6, 1991. Food and Agriculture Organization, Rome, Italy. 1991; 64 – 67.
5. Ciesla WM. Cyprus aphid, *Cinara cupressi*, a new pest of conifers in eastern and southern Africa. FAO Plant Protection Bulletin. 1991; 39(2/3):82-93.
6. Kamunya SM, Day RK, Chagala E, Oling'otie PS. Variation and inheritance resistance to cypress aphid *Cinara cupressi* Bluckton in *Cupressus lusitanica* Miller. Annals of Applied Biology. 1997; 130:27 – 36.
7. Odera JA. Some opportunities for managing aphids of softwoods plantations in Malawi Assistance to Forestry Sector Malawi, MLW/086/020. FAO, Rome. 1991; pp. 135.
8. Obiri JF. Variation of cypress aphid (*Cinara cupressi*) (Bluckton) attack on the family Cupressaceae. Common Forestry Review. 1994; 73(1):43 – 46.

- 206 9. Murphy ST. Cypress severe damaged by the cypress aphid in a rural community in Kenya,  
207 Proceedings of a Workshop at Antalya, Turkey, October 13 – 22, 1997, Eleventh World Forestry  
208 Congress, 1997.
- 209 10. Day RK, Abraham IJ, Kfir R, Murphy ST, Mutitu KE, Chilima CZ. Biological control of homopteran  
210 pests of conifers in Africa, *CAB International 2003, Biological control in IPM systems in Africa*. Eds. P.  
211 Neuenschwander, C. Borgemeister and J. Langewald. 2003; pp. 101 – 112.
- 212 11. Farjon A. A monograph of Cupressaceae and Sciadopitys. Royal Botanical Gardens, Kew. 2005.
- 213 12. Memmott J, Day RK, Godfray HCJ. Intraspecific variation in host quality: the aphid *Cinara cupressi* on  
214 Mexican cypress, *Cupressus lusitanica*. *Ecological Entomology*. 1995; 20:153 – 158.
- 215 13. Orondo SBO, Day RK. Cypress aphid (*Cinara cupressi*) damage to cypress (*Cupressus lusitanica*)  
216 stand in Kenya. *International Journal of Pest Management*. 1994; 40(2):141 – 144.
- 217 14. Normark BB. Molecular systematics and evolution of the aphid family Lachnidae. *Molecular Phylogeny*.  
218 2000; 14:131-140.
- 219 15. Mujtar V, Covelli J, Delfino M, Grau O. Molecular identification of *Cinara cupressi* and *Cinara tujafilina*  
220 (Hemiptera, Aphididae). *Environmental Entomology*. 2009; 38(2):505-512.
- 221 16. Footitt RG, Maw HEL, Von Dohlen CD, Hebert PDN. Species identification of aphids (Insecta:  
222 Hemiptera: Aphididae) through DNA barcodes. *Molecular Ecology Resources*. 2008; 8(6):1189-1201.
- 223 17. Ciesla WM. *Cinara cupressivora*, NAFC-ExFor. 2003. (Available at  
224 [www.spfnic.fs.fed.us/exfor/data/pestreports.cfm?pestidval=161&langdisplay=english](http://www.spfnic.fs.fed.us/exfor/data/pestreports.cfm?pestidval=161&langdisplay=english))
- 225 18. Ortiz-Rivas B, Martínez-Torres D. Combination of molecular data support the existence of three main  
226 lineages in the phylogeny of aphids (Hemiptera: Aphididae) and the basal position of the subfamily  
227 Lachninae. *Molecular Phylogenetics and Evolution*. 2010; 55(1):305-317.
- 228 19. Chilima CZ. Cypress Aphid Control: First African Release of *Pauesia juniperorum*. FRIM Newsletter  
229 (Forestry Research Institute of Malawi). 1995; pp. 2.
- 230 20. Mutitu KE, Ogembo JO, Chagala E. Rearing *Pauesia juniperorum*: A biological control agent of  
231 cypress aphids. KEFRI, Kenya. 2000.
- 232 21. Chilima CZ, Murphy ST. The Introduction of *Pauesia juniperorum* (Stary) (Hymenoptera: Braconidae)  
233 into Malawi for the biological control of *Cinara* species (Homoptera: Aphidae). In: Proceedings of the  
234 XXIIUFRO World Congress, Kuala Lumpur, Malaysia, 2000.

- 235 22. Kairo MT, Murphy ST. Comparative Studies on populations of *Pauesia juniperorum* (Hymenoptera:  
236 Branidae), a biological control agent for *Cinara cupressivora* (Hemiptera: Aphididae). Bulletin of  
237 Entomological Research. 2005; 95(6):597 – 603.
- 238 23. Mutitu KE. Biological Control of Cypress aphid in Kenya. Tree Pest Management Network Newsletter.  
239 2002; pp. 2.
- 240 24. Durak R, Sadowska-Woda I, Machordom A, Borowiak-Sobkowiak B. Biological and genetic studies of  
241 Polish populations of *Cinara tujaefilina*. Bulletin of Insectology. 2008; 61:159-160.
- 242 25. Owuor AL. Exotic conifer aphid in Kenya, their current status and options for management. In: *Exotic*  
243 *Aphid Pests of Conifers. A crisis in African Forestry*. Proceedings of a Workshop at Muguga, Kenya,  
244 June 3 – 6, 1991. Food and Agriculture Organisation, Rome, Italy. 1991; pp. 58 – 63.
- 245 26. Alleck M, Seewooruthun SI, Ramlugun D. Cypress aphid status in Mauritius and trial releases of  
246 *Pauesia juniperorum* (Hymenoptera: Braconidae), a promising biocontrol agent. MAS 2005, Food and  
247 Agricultural Research Council, Reduit, Mauritius. 2005; pp. 313 – 317.
- 248 27. Kairo MTK, Murphy ST. Temperature and plant nutrient effects on development survival and  
249 reproduction of *Cinara* sp. Nov., an invasive pest of cypress trees in Africa. Entomologia  
250 Experimentalis et Applicata. 1999; 92:147-156.
- 251 28. Larsson S. Seasonal Changes in the within-crown distribution of the aphid *Cinara pini*. Oikos. 1985;  
252 45:215 – 222.
- 253 29. Robinet C, Roques A. Direct impacts of recent climate warming on insect populations. Integrative  
254 Zoology. 2010; 5:132-142.
- 255 30. Hulle M, d'Acier AC, Bankhead-Dronnet S, Harrington R. Aphids in the face of global changes.  
256 Comptes Rendus Biologies. 2010; 333:497–503.
- 257 31. Zhou X, Harrington R, Woiwod I, Perry JN, Bale J, Clark SJ. Effects of temperature on aphid  
258 phenology. Global Change Biology. 1995; 1:303-313.
- 259 32. Steinkraus D. Factors affecting transmission of fungal pathogens of aphids. Journal of Invertebrate  
260 Pathology. 2006; 92:125-131.
- 261 33. Watson GW, Voegtlin DJ, Murphy ST, Footitt RG. Biogeography of the *Cinara cupressi* (Hemiptera:  
262 Aphididae) on Cupressaceae, with description of a pest species introduced into Africa. Bulletin of  
263 Entomological Research. 1999; 89(3):271 – 283.
- 264 34. Kiwaso P. Cypress aphid in Uganda. Tree Pest Management Network Newsletter. 2002; pp. 3.

- 265 35. Mwangi JG. Integrated Pest Management Model for Kenya. National Cypress Aphid Project, Kenya.  
266 2002.
- 267 36. Mills NJ. Biological control of forest aphid pests in Africa. Bulletin of Entomological Research. 1990;  
268 80:31 – 36.
- 269 37. Wei JN, Bai BB, Yin TS, Wang Y, Yang Y, Zhao LH, Kuang RP, Xiang RJ. Development and use of  
270 parasitoids (Hymenoptera: Aphidiidae and Aphelinidae) for biological control of aphids in China.  
271 Biocontrol Science and Technology. 2005; 15(6):533-551.
- 272 38. Boettner GH, Elkinton JS, Boettner CJ. Effects of a biological control introduction on three non-target  
273 native species of saturniid moths. Conservation Biology. 2000; 14:1798–1806.