

# The Faroe, Oerkney and Sardinia islands are pointing the dielectrophoretic force in the etiology of multiple sclerosis

Cahit CANBAY

Yeditepe University

Department of Electrical and Electronics Engineering, 34755, Istanbul, TURKEY

*e-mail: canbay@yeditepe.edu.tr*

## Abstract

This study aims to explain the cause of prevalence of multiple sclerosis (MS) disease in the Faroe, Oerkney and Sardinia Islands, by pointing the dielectrophoretic force in the etiology of MS. Initially, the geographical and geophysical specifications of these places have been investigated and then, the results were interpreted by considering the effects of the dielectrophoretic force, which is the essential cause of MS disease in the human body. These islands have very suitable conditions in terms of the geographic environment and the constitutive relations to transmit the electromagnetic field lines, compared to the other places. As a result of this study, it has been found out that direct correlation between the electromagnetic fields on the islands and the prevalence of MS disease. In order to support these claims, It has been made use of several geographical and the geophysical data and the results that have been achieved before by the author.

**Keywords:** Causal factor, dielectrophoretic force, Faroe islands, multiple sclerosis, Oerkney islands, Sardinia island.

## 1. INTRODUCTION

Multiple sclerosis (MS) is one of the most common neurological disorders whose symptoms occur as a result of the inflammation and breakdown in myelin. Myelin is made of lipids of proteins, tau cells and protective layers of nerve fibers. It insulates the nerve fibers found in the central nervous system (CNS) and the peripheral nervous system(PNS) by surrounding them. This properties may be likened to the loss of insulating material around an electrical wire. Demyelination is a degenerative process that erodes away the myelin sheath. The myelin basic proteins (MBPs), other particles which are separated from myelin sheath accumulate in the plaque-like debris form in some places of the brain and the spinal cord (Canbay, 2010). The intensity of electrical signals, traveling along the nerves, decrease and become noisy. A patient affected from MS disease sometimes lives a progressive disorders with the functions that are controlled by the CNS such as vision, speech, walking, writing, attention and memory.

The dielectrophoretic force has been found as the essential cause of MS disease, which is one of the most common neurological disorders (Canbay, 2010). In the study, the correlation between MS and the electromagnetic fields has been shown with the seven different ways and arguments (Canbay, 2013). In a recent study, it was explained that both the radiologically isolated syndrome (RIS) which occurs following the magnetic resonance imaging (MRI) examinations and highly suggestive Multiple Sclerosis disease occurs as a result of the dielectrophoretic force (Canbay, 2014). In fact, the study (Canbay, 2014) about RIS was evidence that the RIS can not be a correct concept.

Frequent MRI applications, the improper selection of the operating frequency of MRI scanners could lead to occur of MS-like syndromes in brain. In a study in 1999 (Westland K W, Pollard J D, Sander S, Bonner J G, Linington C and McLeod JG, 1999) has very interesting results about the demyelination. In that study, the researchers, systemically, injected reactive T cells to a non-neuro antigen (ovalbumin) of Lewis rats and caused them to accumulate in thoracic dorsal column by a prior injection of ovalbumin. According to their findings and interpretations, a more profound conduction block and more plaque-like region of demyelination were observed in the animals. However, they were not able to explain completely of the main cause of their findings. If the researchers had taken the dielectrophoretic effect on the myelin into account, they would be able to explain both the main cause of occurred demyelination, plaque-like structure, and the motion of T cells in a dielectrophoretic force field, like in the studies (Canbay, 2015) (Heida T, Rutten WL, Marani E, 2001). Moreover, when the stimulator had not been applied on the mouse, the plaque-like formations would not be observed.


This study proves the correctness of the author's hypotheses, showing the relationship between MS and the electromagnetic environment issues in the world. Especially, the Faroe, Oerkey and Sardinia Islands, widely known as the places with higher prevalence probable of MS disease in the world, has been focused in terms of the relationship between the electromagnetic environment and MS disease (Canbay, 2010) (Canbay, 2013) (Canbay, 2014), (Canbay, 2015). The various causal factors, which possible to be related to the distinct increase in multiple sclerosis in the Faroe, Oerkey, and Sardinia islands, have been studied on by the researchers. Some researchers underline of the effect of industrial changes and occupation on the islands. For some researchers, birthplace is considered as an indicator of MS disease (Lauer, 1986). The Sardinian cohort has been observed for a long time to find out the relation between the early multiple sclerosis and retrovirus (Sotgiu, et al., 2006). In the recent years, researchers and the national and international organizations have focused on the effect of the genetics (Ramagopalan, McMahon, Dymment, Sadovnick, Ebers, & Wittkowski, 2009) (Wang, et al., 2011), the immune system (Bates, Feix, Boggs, & Harauz, 2004), the environmental factors (Canbay, 2010) on MS disease. Some researchers have put forward that the high intake of vitamin D, which is proportional to the solar radiation, is associated with lower prevalence rates of the multiple sclerosis (Schwarz & Leweling, 2005). Some have the against view (Beebe, Kurtzke, Kurland, Auth, & Nagler, 1967). In a study, it is proposed that multiple sclerosis may be transmitted chiefly by sexual contact. Higher prevalence rate of multiple sclerosis on the Oerkey islands has been described by the increase in the number of troops on the island during the Second World War (Hawkes, 2002). Despite the fact that these results achieved by various studies by other researchers, they give the similar results with my approach and findings'. Yet, their studies should have been supported with more powerful arguments, as in my studies (Canbay, 2010) (Canbay, 2013) (Canbay, 2014) (Canbay, 2015).

According to the author's comparative study between the keraunic map of the world prepared by IEEE and the common reports of the WHO (World Health Organization) with MSIF (Multiple Sclerosis International Federation) dated 2008 and 2013 (Canbay, 2010) (Canbay, 2014), it has been found the some clues as to the respond the question "why some places have higher risk with respect to the other places in terms of Multiple Sclerosis disease?".


## **2. SUMMARY OF AUTHOR'S HYPOTHESES ON THE ETIOLOGY OF MULTIPLE SCLEROSIS DISEASE**

In the author's previous studies, it has been shown that the dielectrophoretic force is the main cause of MS disease through various approaches. To contribute to a better understanding of this issue, these approaches can be summarized as follows:

1-The effect of the dielectrophoretic force on Myelin Basic Protein (MBP) was determined in white matter, grey matter and cerebrospinal fluid (CSF) by use of Clausius-Mossotti (CM) equation as shown in Figure 1, Figure 2. Since the MBP, has a molecular weight of 18.5 kDa, a radius of 1.525 nm and an adhesion force of 0.05 mN/m between MBP and the nerve cell, the dielectrophoretic force at some frequencies and under long term exposure conditions, pulls off the MBP from the nerve cell (Canbay, 2010) (Canbay, 2014). The Figure 1 has been calculated by CM equation, taking into account the dispersive properties of all constitutive relations of human brain. In this case, the results become parallel to possible experimental results. If the event is viewed in terms of MBP and dielectrophoretic force, this graph explains many things. The extra low frequency electromagnetic fields (ELF) are very important in terms of creating ground to MS disease. As seen from this graph, in the eras that absence of the artificial electromagnetic field sources, MS disease had and had to be (look Schumann resonance frequencies, history of MS disease). Today, living things live under the influence of both natural and artificial electromagnetic fields. Some of the important artificial electromagnetic fields sources are the electrical power distribution systems (50-60 Hz), MRI scanners (63 MHz, 125 MHz ..), base stations, mobile phones and others. As seen in Figure 1, the MRI scanners working in 1.5 Tesla (63 MHz) can cause the protein leakage into the CSF. However, for the brain MRI scanners working in 3 Tesla (125 MHz) does not seem a problem in terms of MS disease.



**Figure 1.** Variations of dielectrophoretic force, affecting MBP in white matter, gray matter and CSF with respect to frequency, ( $10^0$ - $10^{11}$ ) Hz (Canbay, 2014).



**Figure 2.** Dielectrophoretic field affecting an ion and a dielectric particle in schematic representation of human brain (Canbay, 2010).


2-Electromagnetic field distribution, which results from natural electromagnetic field resources, even when there is no artificial electromagnetic field resource, induces an electromagnetic field in the human body. It can be said that lightnings, which form the final structure of electrosphere, lead to an electromagnetic field distribution released at Schumann frequencies in the electrosphere. The Schumann Resonance is a set of spectrum peaks in the ELF portion of the Earth's electromagnetic field spectrum. The nominal average the Schumann Resonance frequencies are observed 7.8, 14, 20, 26, 33, 39, and 45 Hz with slight diurnal variations. The maximum dielectrophoretic force in the human brain occurs at around of the Schumann Resonance frequencies (Figure 1). When the Schumann Resonance frequencies are compared with the results of Clausius-Mossotti equation as shown in Figure 1, it is clear that the history of MS disease is as old as the history of humanity (Canbay, 2010) (Canbay, 2014).

3-After a lightning, the local electric field intensity approaches the fair air electric field intensity level. As a result of this, a relaxed medium occurs. The fair air conditions provide minimum electric field intensity in our environment. On the contrary, cloudy weather conditions provide higher electric field intensity in the medium.


In the comparative study between keraunic map of the world prepared by IEEE and the common report of the WHO (World Health Organization) with MSIF (Multiple Sclerosis International Federation) dated 2013 (Canbay, 2014), we can find some clues in answering the question "why some places have higher risk with respect to other places in terms of prevalence of Multiple Sclerosis disease". Figure 3 and Figure 4 show that the prevalence of Multiple Sclerosis disease inversely proportional to the number of annual lightning. Generally, I don't need to investigate the prevalence of MS disease in South America and Africa. According to my hypotheses, most of countries in these continents had to be painted in blue and dark blue colors (Canbay, 2010). The comparison of the keraunic map in Figure 3 with the epidemiological map in Figure 4 and the consideration of the aforementioned data support my hypotheses.

Distribution of the electromagnetic field in the earth's atmosphere is influenced from

solar winds. This interaction mechanism may be described by Faraday's law. Compared the annual variations of the sunspot numbers (C.T. Russell, J.G. Luhmann, L.K. Jian, RG2004/2010) with Kurtzke's results regarding the annual incidence rates per 100.000 population of MS disease in the Faroe and Oerkney islands (Kurtzke, 2006) occurs an interesting result.



**Figure 3.** The keraunic map (Canbay, 2010).




**Figure 4.** The distribution of MS in the world (Canbay, 2014).

4-The dielectrophoretic force is capable of separating lipids of proteins from the nerve cell and accumulating the separated protein particles at places where the gradients


of square of electric field reach their maximum values in the human body. MS symptoms also can be observed in the same places (Canbay, 2010). Since the gradients of square of electric field in the brain of animals with four legs or without legs is approximately equal to zero. MS disease is not observed in these animals. MS is a disease of human's disease and will remain so in the future as well. There will not be any advantage of animals studies concerning MS without taking into account of the author's hypotheses on the etiology of MS disease. The passive mice cannot be used in the MS disease-related experiments.

5-The connection between electromagnetic pollution and asthma, diabetes, multiple sclerosis, chronic fatigue, fibromyalgia has been investigated in the study (Havas, 2006). In this study, GS (Graham/Stetzer) filters, microsurge meters operating at ELF and RF frequencies were used to reduce and monitor the dirty electricity inside the building where the patient lived. The observation results show that the symptoms of these disorders were reduced.

6-The interaction between the neurons and electromagnetic fields can be investigated by in vitro experiments. For example, as shown in Figure 6 and Figure 7, the fetal cortical rat neurons were trapped in the center of an electrode arrangement operating at different amplitudes and frequency ranges (Canbay, 2014).



**Figure 5.** (A) The electrode structure represented by four point charges. The point charges are located at  $100 \mu\text{m}$  from the center and are given such a value that the potential at a distance  $R = 50 \mu\text{m}$  (a distance equal to the “diameter” of the electrode tips) from these points is  $5 \text{ V}$  (or  $-5 \text{ V}$ ). (B) The relative dielectrophoretic force in the  $xy$  plane at  $z = 100 \mu\text{m}$ . The square of the electric field (the solid lines, contours, give the iso-electric field lines) and the gradient of this field (arrows) in (C) the  $xy$  plane at  $z = 0 \mu\text{m}$ , and (D) the  $(x = y)z$  plane at  $z = 0 \mu\text{m}$  (Heida T, Rutten WL, Marani E, 2001).



**Figure 6.** Image nr. 180 of several experiments using different amplitudes (i.e., the situation after 30 min. of field application). (A) No input signal (reference experiment), (B) 1 V/12 MHz, (C) 3 V/12 MHz, and (D) 5 V/12 MHz (Heida T, Rutten WL, Marani E, 2001).

7-A volcano in Iceland erupted in March of 2010. The lavas, volcanic ash plumes and the highest concentration of aerosols (tiny airborne particles, sulphur dioxide) in the ash plume spread into the atmosphere. As shown in Figure 15, the sulphur dioxide plumes located at higher altitudes are less affected by winds and concentrate on a strip over the Faroe and Oerkney Islands. The reason for higher concentration of aerosols on this strip over the Faroe and Oerkney Islands is the dielectrophoretic force over this region, just as in the brain, as conceptually expected to be by the author (Canbay, 2013).

#### 8- SARR-MS-MRI-RIS relationship

In MRI scanners, time varying gradient magnetic fields may stimulate nerves by inducing gradient electric fields in human body. The movement of the dielectric particle within the liquid in any gradient electrical field depends on the particle diameter, the dielectric permittivity of the liquid and the particle, and the electric field intensity. The CM equation can be used to find the force that acts on a dielectric particle.

In MRI applications, it is very important to minimize the dielectrophoretic force for MS disease. When  $\text{Re}[K(\omega)] > 0$ , particles are attracted to regions of stronger electric field when their permittivity exceeds that of the suspension medium. This is called positive dielectrophoresis (p-DEP). When permittivity of the suspending medium is greater than that of the particles, this results in motion of particles to lesser electric field ( $\text{Re}[K(\omega)] < 0$ ), as observed in Figure 6C. This is called negative dielectrophoresis (n-DEP) (Canbay, 2010).

The general concept of the gradient system in MRI scanners is used to provide the image of a biological slice. The gradient electrical fields occur as a result of discontinuity along the interface between two biologic tissues and also as result of the gradient effect based on the electrical field polarization generated by the gradient and the radio frequency (RF) coils. The dielectrophoretic force generated by the gradient and the radio frequency (RF) coils stimulates the MBP and other dielectric particles in the media such




as CSF, GM and WM within the human brain and spinal cord (Figure 7). Moreover, this force can also separate these particles from nerve cells under appropriate conditions and provide collecting of them in certain fields. At the same time, the dielectrophoretic force is proportional to the gradient of the Specific Absorption Rate (SAR). The Specific Absorption Rate Ratios (SARR) has been given on the brain and spinal cord model of human (Canbay, 2014). Table. 1 denotes the findings of a study that identifies the risk factors for the development of the first clinical event in a multinational cohort of RIS subjects and provides updates to the original 2009 RIS criteria. As it is seen from their findings, the presence of a spinal cord lesion was observed in 38.7%, CSF profiles were abnormal in 40.9%, and a positive family history for MS observed in 9.9%.

| Total Subject | Female       | Male | The mean age | Standard Deviation | mean clinical follow-up | Standard Deviation | CSF profiles (abnormal) | Spinal Cord lesion | MS observed (Positive Family History) |
|---------------|--------------|------|--------------|--------------------|-------------------------|--------------------|-------------------------|--------------------|---------------------------------------|
| 375           | 298 (79.5 %) | 78   | 37.1         | 11                 | 4.6 years               | 4 %                | 40.9 %                  | 38.7 %             | 9.9 %                                 |

**Table 1.** The findings of a study to identify risk factors for the development of the first clinical event in a multinational cohort (US, IR, TR, ITALY) of RIS subjects (Okuda D T, Mowry E M, Beheshtian A, Waubant E, Baranzini S E, Goodin D S, Hauser S L, Pelletier D, 2009)

In the author's recent study (Canbay, 2014), the MRI-RIS-MS connections which demonstrate SARR-Dielectrophoretic force-frequency relationship, were given. In this study, the regions may be effected by the dielectrophoretic force were computed by using the dispersive medium parameters of the brain tissues, CM and the SARR equations. As shown in Figure 8, the possible lesion places after the MRI scans was determined as, firstly, the cerebrospinal fluid, secondly, the spinal cord. The Figure 8, denote quite parallel results with the findings of the study (Okuda D T, Mowry E M, Beheshtian A, Waubant E, Baranzini S E, Goodin D S, Hauser S L, Pelletier D, 2009).






**Figure 7.** Schematic representation of the brain and the spinal cord in the gradient electromagnetic fields of an MRI scanner (Canbay, 2014).

The results, as shown in Figure 8, can be found if SAR ratios are calculated considering the dispersive properties of two media. In case of  $SARR=1$ , there will not be any gradient electric field because of the nature of biological tissues. The gradient electric field in the laboratory experiment (Figure 5) and (Figure 6) and the MRI examinations has a very intensive effect on MBP. MBPs (like nerve cells polarized and collected within half an hour in the laboratory conditions) will accumulate in the direction where the gradient of square of the electrical field is at its maximum. When the brain and spinal cord are located in the natural or artificial electromagnetic fields (Figure 7), as explained in the approach 1, in the laboratory experiment (Figure 5) and (Figure 6) and in the MRI radio frequency coils (Figure 9), they can be understood to be under similar conditions in terms of the occurrence of dielectrophoretic force. The electric field intensity in the brain and spinal cord under normal conditions is lower when compared to the electrical field intensity under the brain and spinal cord in the gradient and radio frequency coils of MRI scanners. If suitable conditions occur for developing of MS disease, the accumulation of MBP will also occur over the years.


The idea of RIS that emerged as a result of MRI practices is neither a new symptom imitating MS nor an anticipated one. Clearly, RIS can be accepted as the laboratory results of the dielectrophoretic force.

In this test, as the Clausius-Mossotti factor was  $Re(K) < 0$ , nerve cells were collected in the direction of decreasing electric field; in other words, in the center of the electrode array. Figure 8 presents the electric field distribution and illustrates where nerves collect and shows the number of nerves based on the exposure time of the electrical field (Canbay, 2014). Similarly, considering that the brain and the spinal cord are placed within the electrical field of the gradient and RF coils of MRI, as shown schematically in Figure 9, an aggregation of MBPs within the cerebellum and spinal cord based on the gradient of the squared electric field will be ensured.

The force which creates MS over several years in direction of CSF-GM-WM, is the same dielectrophoretic force that also creates RIS. However, the electric field intensity and the gradient of the squared electric field in an MRI scanner are more powerful.



**Figure 8.** SARR versus frequency, ( $10^6$ - $10^9$ ) Hz (Canbay, 2014)



**Figure 9.** The contour of the induced electromagnetic fields produced by cylindrical Z-coils. (a)–(c) Magnetic fields in the  $X = 0:1$  m plane. (d) Electric field (E) in the  $X = 0$  plane (Liu F, H. Zhao H, Crozier S., 2003).

Therefore in the laboratory, the collection or orientation of the MBPs or any dielectric particles within half an hour, where the gradient of the squared intensive electric field is at its maximum occurred because of the same dielectrophoretic force.

9-The breakdown mechanism of the blood brain barrier (BBB) has been explained by the interaction of the myelin basic protein (MBP) with the particles separated from T cells by the dielectrophoretic force effect on the BBB, the myelin and the particles.

Whereas, within a human brain that is under the effect of weak dielectrophoretic force, the collection of MBPs occurs very slowly due to the same reasons, and the observation


of symptoms of MS take 20-30 years. No one can ever expect to prove such a long-term process by observation. RIS is an accelerated form of MS disease. There is no this kind of syndrome. RIS is to be an expecting event created by a little more intensive dielectrophoretic force effect of MRI scanner.

### 3. METHOD


The author's hypotheses related to the etiology of MS disease supported with the scientific evidences can be applied clearly to research of the prevalence of MS disease in everywhere on the world. The necessary and sufficient conditions for MS can be investigated by the appraisal of the etiology of MS disease in the light of the impact of the dielectrophoretic force (Canbay, 2010) (Canbay, 2013) (Canbay, 2014) (Canbay, 2015). If there are necessary and sufficient conditions in any place regarding the occurrence of MS disease, the prevalence of probable MS will be more with respect to the other places. For the study (Canbay, 2010), the necessary and sufficient conditions for MS disease to occur is directly related to the conditions of the place inhabited at growing ages, namely until the ages of 20. At any place, where the frequency of occurrence of MS disease is higher, the environmental conditions are as follows:

- a- The year average of the number of lightning at that residential area is too small (except of deserts).
- b- The average annual number of cloudy days at that residential area is too much.
- c- The residential area is on the high mountains and hills.
- d- The residential area is located far from the large-scale conductive structures.
- e- As mentioned previously, the residential area contains some kinds of natural and artificial electromagnetic field sources or very close to them.
- f- Stress and contraction of the muscles, as required the principle of piezoelectric effect, generates the internal electric field intensity in the human body. (This subject is related to the person's psychology and the person's social and physical environment.)
- g- The increase of the dense and frequent use of the Magnetic Resonance Imaging (MRI) device, which plays a significant role in the diagnosis of Multiple Sclerosis disease, has brought along new concerns (Canbay, 2014) (Canbay, 2015). Additionally, increasing of stress has also an affect (as a factor of inner dynamic, stress is placebo equivalent of mechanic stress) that can be assumed as the source of dielectrophoretic field in the human body on MS. Yawning-Stretching movements are a behavioral syndrome associated with trans cranial application of electromagnetic fields for MS (Sandyk, 1998) because of eliminating polarized charges by the electromagnetic fields in the human body.

Figure 10 denotes two persons standing on the homogeneous ground in the fair air condition. In this case, the electromagnetic field distribution in the ground and the air has the regular electrical field distribution. Therefore, two persons are under the same electromagnetic field effect. But still the induced electromagnetic fields in the bodies of the two persons may be different from each other.



**Figure 10.** Two persons in the regular electromagnetic field distribution.




**Figure 11.** Three persons in the irregular electromagnetic field distribution.

As shown in Figure 11, if there are conductive structures, such as trees, the buried large metal materials, and the metal towers close to human body, the lines of natural and artificial electromagnetic fields flow on these conductive structures. As a result of this interpretation, the amount of electromagnetic field lines flowing on the human bodies varies spatially and so does the risk of MS for some individuals. Especially, as shown in Figure 11, while the conductive structure provides advantageous for the persons located at point A and C, contrary disadvantageous for the person at point B. However, the tree provides an additional advantage for the person at point C in terms of the electromagnetic effect. This result shows us that everybody sharing the same environment can not be


regarded as a nominate for MS. Absolutely, a person who nominate for MS disease should have stayed under the necessary and sufficient conditions from angle of MS disease for a long time in his/her growing era. In the literature, there are many studies expressing the genetic, the immunological reasons, and the correlation between the occurrence of relapses and the seasonal environmental factors, infections (Hawkes, 2002) and serum vitamin D levels, and speculated convincingly, regarding the people with MS disease living on these islands. In fact, it should have been assumed that their findings are the results of electromagnetic fields and the effect of dielectrophoretic force. The essential cause of maximum prevalence of MS disease is the existence of the suitable conditions because of the natural and the artificial electromagnetic field distribution on these islands. This study aims, both to give a new material support for the previous studies and to find the answer to the question, why these islands have risk factors from angle of multiple sclerosis disease and to prove that MS disease is not only an old disease but also a modern day disease. Moreover, MS disease has increased depending on the uncontrolled electromagnetic technologies and the sources.

#### 4. GEOGRAPHICAL AND GEOPHYSICAL STRUCTURE OF THE ISLANDS


Faroe Islands (62° N, 7° W) have a total area of 1400 km<sup>2</sup>, an average height of 300 m above sea level and formed a part of the North Atlantic Brito-Arctic Cenozoic Igneous Province that extends from the British Isles to Greenland. Faroe and Oerkney islands share geographic, ethnic (is not taken into account in the present study), and environmental similarities with other North Sea and Canadian countries of high MS incidence. The prevalence of MS per year in Faroe and Oerkney Islands are 66(1998) and 193 (1983), respectively (Pugliatti, Sotgiu, & Rosati, 2002). **Faroe Islands are the most sunless region in the World, having an entirely cloudless sky only, averagely, five days in a year. As a consequence of that, there is very little summer heat. At the same time, the number of days with lightning (keraunic level) on each island is lower, approximately equals 1** (Canbay, 2010). Since the warm period is really short, the sprouts of the trees ripe really hard, and only a very few kinds of trees can survive the winter. The trees make little progress in their effort upwards. Faroe Islands lie on a continental fragment, which has a total thickness of about 10-15 km of igneous rock added as extrusive lavas at the top. In addition to the zeolites which have (0-2) S/m conductivity, native copper (at 20 C° 2.4 x10<sup>6</sup> S/m) and pyrite deposits (1-10<sup>5</sup>) S/m were found in different places on Faroe Islands (Larsen, Knudsen, Frei, Frei, Rasmussen, & Whitman, 2006) (White, 2008) (Jørgensen, 2006). Figure 12, Figure 13 and Figure 14 show the representative distribution of electric fields on an island under the three different conditions. Figure 12 denotes the distribution of electric fields on the island in the fair air condition. As shown in Figure 12, the electric field line density over the island changes depending on the topographical structure and electrical parameters of the island. However, as shown in Figure 13, the electric field line density over the mine deposits is higher than the first situation. As shown in Figure 14, the electric field intensities over some places on the island are very higher compared to the situations (Figure 12) and (Figure 13).



**Figure 12.** Schematic representation of the distribution of electromagnetic field on an island in case of fair air and homogeneous island.




**Figure 13.** Schematic representation of the distribution of the electromagnetic field on an island in case of the fair air and heterogene island.



**Figure 14.** Schematic representation of the distribution of electromagnetic field on an island in case of the cloudy day and heterogeneous island.

The causes of the variations of the electric field line densities over the island are; the effect of electrically charged cloud over the island and the properties of electrical parameters of the underground of the island.

The explanation of the causes of higher prevalence of MS disease on the islands is interesting. I was thinking about the accumulation of the dielectric particles in the form of the plaques in the human brain due to the irregular electromagnetic field distribution. As known, Eyjafjallajökull volcano in Iceland erupted in March of 2010. Suddenly, I wanted to expect dual relations between the brain and our globe. As I expected, the tiny particles spread into the atmosphere due to Eyjafjallajökull volcano eruption should be collected onto Faroe Islands.



**Figure 15.** Distribution of the sulfur dioxide plumes in the electrosphere (NOAA Office of Satellite and Product Operations)



I have investigated the behavior of the sulfur dioxide plumes (very tiny airborne particles compare to volcanic ash) from Eyjafjallajökull volcano eruption in black southeast of Iceland. The lavas, volcanic ash plumes and the highest concentration of aerosols (tiny airborne particles, sulfur dioxide) in the ash plume spread into the atmosphere. NASA's Goddard Space Flight Centre, Greenbelt, Md. has used data from the Ozone Monitoring Instrument (OMI) to create satellite image updated on volcanic ash and sulfur dioxide plumes. The image in Figure 15 from the Washington Volcanic Ash Advisory Center on May 6 (NOAA Office of Satellite and Product Operations), shows the highest concentration of tiny airborne particles in the ash plume from the Eyjafjallajökull volcano in black southeast of Iceland. As shown in Figure 15, the sulfur dioxide plumes located at higher altitudes are less affected by winds and concentrate on a strip over the Faroe and Oerkney Islands. The cause of higher concentration of aerosols on this strip over the Faroe and Oerkney Islands is dielectrophoretic force over this region, just as in the brain, as conceptually expected to be by the author.

Sardinia is the second greatest island of the Mediterranean: it measures 24.090 km<sup>2</sup>, it is located between 38° 51' 52 " and 41° 15' 42" of latitude North and 8° 8' and 9° 50' of East longitude. The coasts of Sardinia (1.849 km long) are generally high and rocky, rectilinear for kilometers with vast and deep bays and fjords surrounded by smaller atolls. **The number of lightning days per year (keraunic level), is relatively lower compared to other places in the same latitude, is between about 1-5. This situation is unexpected for Sardinia Island and very crucial.** The reasons for decreasing of the lightning occurrence over Sardinia island has been explained in a study (Dietrich, Casella, Paola, Formenton, Mugnai, & Sano, 2011). The main deposits among the ore deposits of the Cenozoic volcanics under the Sardinia Island are kaolinite, bentonite, and pyrite which have electrical conductivities (0.4-1) S/m, (2-5) S/m, and average conductivity (10<sup>3</sup>) S/m, respectively (Palomba, Padalino, & Marchin, 2006) (Stewart, 1937) (Cita, Santambrogio, Melillo, & Rogate, 1990) (Carozzo, et al., 1997) (Shuey, 1976). These constitutive relations vary depending on dispersive properties from their water and NaCl content, temperature, pressure, and frequency point of view. People living on the Sardinia Island can be considered to be living very close to the conductive ground due to the sea water and also due to the existence of the ore deposits under the ground. As a result, the electric field lines over the islands are especially concentrated on mine deposits. Therefore, this concentrated electric field lines on the Sardinia Island triggers MS disease in this region.

## 5-RESULT

The dielectrophoretic force has been found as the essential environmental cause of the most common neurological disorder Multiple Sclerosis (MS) (Canbay, 2010). Moreover, the locations of the maximum gradient of squared electrical field in the parts of human body have been determined as possible places where the MS symptoms can be seen. The Faroe, Sardinia and Oerkney islands have the special properties in terms of to provide of the background the gradient electromagnetic fields in/on/over their environmental conditions, compared to the other places. The places, such as sea, lake and river shores may be suitable locations for MS patients. In this study, it has been shown that the prevalence of MS disease on the Faroe, Oerkney and Sardinia Islands related to the dielectrophoretic force, by taking into account all the findings and the results of author's the previous studies. The RIS-MRI-MS connections are the most important quasi experimental evidences confirming the close connection between MS and the

dielectrophoretic force. The dielectrophoretic force is the crucial factor for the etiology of MS disease (Canbay, 2015). Considering the author's hypotheses, it can be understood why the prevalence of MS is higher over these islands.

## CONCLUSION

In any living environment, the necessary and sufficient conditions to cause MS disease are as follows:

- a- The residential area in the region has too much of number of cloudy days, too low number of lightning days, in a year.
- b- The ground surface and/or underground structure are conductive compared to adjacent places.
- c- To live a long time in a higher place, which has too much of number of cloudy days, too low of number of lightning days in a year.
- d- To live far from the conductive structures on the ground surface, such as trees, the large metal materials etc.
- e- To live close to the electromagnetic field sources, which operating at the aforementioned conditions.

a'- The Faroe Islands are the most sunless region in the world, having an entirely cloudless sky only, averagely, five days a year. The electric field intensity increases and remains approximately constant at high level during the cloudy days. The Oerkney islands have the geographical and environmental similarities with the Faroe Islands. The annual number of days with lightning (keraunic level) on each island is lower, approximately equals 1 (Canbay, 2010).

b'- As mentioned before, these islands have a quite better electrical conductive ground surface or underground structure.

c'- The islands have higher altitude with respect to the sea level.

d'- There are no large trees on Faroe and Oerkney islands.

e'- The Faroe, Oerkney and Sardinia islands are situated far away from the mainland. Therefore, all electromagnetic sources such as the base stations, the TV transmitters, the high voltage transmission lines have to be place around the residential area on the islands. In addition, the natural electromagnetic features of the islands in terms of the appropriateness of MS disease were tested. Also, the stress of a living person in a region like the Faroe, Oerkney islands which has sunshine only five days in a year, can be compared with others.

Not only for these islands, learning of the electromagnetic properties of anywhere in the world is as vital as determining the weather reports in the world we live on it. Therefore, the maps, showing the distribution of electromagnetic field, covering the entire electromagnetic spectrum, will be important for the protection of the health of the living things and the environment. After this approach, MS disease will not remain as an unsolved mystery for the Faroe, Oerkney and Sardinia Islands anymore.

**Acknowledgement**-The author would like to thank Mr. Yaman Akgün in Yeditepe University, Mrs. Ümit Canbay and Begimay Ida Schmidt for their contributions in the preparation of the paper.

The author declares no financial or commercial conflicts of interest.

## REFERENCES

- (n.d.). Retrieved June 09, 2011, from NOAA Office of Satellite and Product Operations: [www.ssd.noaa.gov/VAAC](http://www.ssd.noaa.gov/VAAC)
- Bates, I. R., Feix, J. B., Boggs, J. M., & Harauz, G. (2004). An immunodominant epitope of myelin basic protein is an amphipathic  $\alpha$ -helix. *J Biol Chem*, 279(7), 5757-5764.
- Beebe, G. W., Kurtzke, J. F., Kurland, L. T., Auth, T. L., & Nagler, B. (1967). Studies on the natural history of multiple sclerosis 3. Epidemiologic analysis of the army experience in world war II. *Neurol*, 17, 1-17.
- C.T. Russell, J.G. Luhmann, L.K. Jian. (RG2004/2010). How unprecedented a solar minimum. *Geophysics*, 48, s. 1-16.
- Canbay, C. (2010). The essential environmental cause of multiple sclerosis disease. *Prog Electromagn Res, PIER*, 101, 375-391.
- Canbay, C. (2013, April 11-14). The appraisal of the etiology of the multiple sclerosis disease in the light of the impact of the dielectrophoretic force. *7th World Congress on Controversies in Neurology (CONy)*.
- Canbay, C. (2014, July). The radiologically isolated syndrome is the last link of the chain for understanding the etiology of Multiple Sclerosis disease. *Eur Sci J*, 10(21), 20-35 (ISSN: 1857-7881(print) e-1857-7431).
- Canbay, C. (2015). Multiple Sclerosis is not a disease of the genetic and immune system origin. *SYLWAN*, 159(3), 1-8.
- Carozzo, M. T., Balia, R., Loddo, M., Luzio, D., Marqiotto, C., Quarta, T., et al. (1997). Gravity modelling of the lower crust in Sardinia(Italy). *Ann Geofis*, 40(5), 1083-1094.
- Cita, M. B., Santambrogio, S., Melillo, B., & Rogate, F. (1990). 14. Messinian paleoenvironments: New evidence from the Tyrrhenian sea (ODP LEG 107). *Proc Ocean Drill Program, Sci Results*, 107, 211-227.
- Dietrich, S., Casella, D., Paola, F. D., Formenton, M., Mugnai, A., & Sano, P. (2011). Lightning-based propagation of convective rain fields. *Nat Hazards Earth Syst Sci*, 11, 1571-1581.

Havas, M. (2006). Electromagnetic hypersensitivity: Biological effects of dirty electricity with emphasis on diabetes and multiple sclerosis. *Electromagn Biol and Med*, 25, s. 259-268.

Hawkes, C. H. (2002). Is multiple sclerosis a sexually transmitted infection? *J Neurol, Neurosurg Psychiatry*, 73, 439–443.

Heida T, Rutten WL, Marani E. (2001). Dielectrophoretic trapping of dissociated fetal cortical rat neurons. *IEEE Transaction on Biomedical Engineering*(48), s. 921-930.

Jørgensen, O. (2006). The regional distribution of zeolites in the basalts of the Faroe islands and the significance of zeolites as palaeotemperature indicators. *Geol Surv Den Greenl Bull, GEUS*, 9, 123–156.

Kurtzke, J. F. (2006). An epidemiologic view of the etiology of multiple sclerosis. *Arc Neuroscien (mex)*, 11(2), s. 90-99.

Larsen, M., Knudsen, C., Frei, D., Frei, M., Rasmussen, T., & Whitman, A. G. (2006). East Greenland and Faroe–Shetland sediment Provenance and Palaeogenen Sand Dispersal Systems. *Geol Surv Den Greenl Bull, GEUS*, 10, 29–32.

Lauer, K. (1986). Some comments on the occurrence of multiple sclerosis in the Faroe islands. *J Neurol*, 233, 171-173.

Liu F, H. Zhao H, Crozier S. (2003). On the induced electric field gradients in the human body for magnetic stimulation by gradient coils in MRI. *IEEE Trans Biomed Eng*, 50, s. 17-20.

Okuda D T, Mowry E M, Beheshtian A, Waubant E, Baranzini S E, Goodin D S, Hauser S L, Pelletier D. (2009). Incidental MRI anomalies suggestive of multiple sclerosis: the radiologically isolated syndrome . *Neurol*, 72, s. 800-805.

Palomba, M., Padalino, G., & Marchin, M. (2006). Industrial mineral occurrences associated with cenozoic volcanic rocks of Sardinia (Italy): Geological, mineralogical, geochemical features and genetic implications. *Ore Geol Rev J*, 29, 118–145.

Pugliatti, M., Sotgiu, S., & Rosati, G. (2002). The worldwide prevalence of multiple sclerosis. *Clinical Neurology & Neurosurgery*, 104, 182-191.

Ramagopalan, S. V., McMahon, R., Dymment, D. A., Sadovnick, A. D., Ebers, G. C., & Wittkowski, K. M. (2009). An extension to a statistical approach for family based association studies provides insights into genetic risk factors for multiple sclerosis in the HLA-DRB1 gene. *BMC Med Gen*, 10.

Sandyk, R. (1998). Yawning and stretching- A behavioural syndrome associated with transcranial application of electromagnetic fields in multiple sclerosis. *Int J Neurosci*, 95, 107-113.

- Schwarz, S., & Leweling, H. (2005). Multiple sclerosis and nutrition. *Mult Scler*, 11(1), 24-32.
- Shuey, R. T. (1976). The electrical resistivity of galena, pyrite, and chalcopyrite. *Am Min*, 61, 248-259.
- Sotgiu, S., Arru, G., Mameli, G., Serra, C., Pugliatti, M., Rosati, G., et al. (2006). Multiple sclerosis-associated retrovirus in early multiple sclerosis: a six-year follow-up of a Sardinian cohort. *Mult Scler*, 12, 698-703.
- Stewart, D. (1937). Petrography of some rocks from the south Orkney islands and the antarctic archipelago. *Am Min*, 22, 178-194.
- Wang, J. H., Pappas, D., Jager, P. L., Pelletier, D., Bakker, P. I., Kappos, L., et al. (2011). Modelling the cumulative genetic risk factors for Multiple Sclerosis from Genome-Wide association data. *Gen Med*, 3(1/3), 1-11.
- Westland K W, Pollard J D, Sander S, Bonner J G, Linington C and McLeod JG. (1999). Activated non-neurol specific T cells open the blood-brain barrier to circulating antibodies. *brain*, 122, s. 1283-1291.
- White, R. S. (2008). Lower-crustal intrusion on the north atlantic continental margin. *Nature*, 452, 460-464.

