NUTRITIVE POTENTIALS OF *Centrosema pubescens Mimosa invisa and Pueraria phaseoloides* LEAF MEALS FOR GROWTH RESPONSES OF CHICKEN

F.C. NWORGU and G.N. EGBUNIKE+
Federal College of Animal Health and Production Technology, Institute of Agricultural Research and Training. PMB 5029 Ibadan, Nigeria

+ Animal Physiology Laboratory, Department of Animal Science. University of Ibadan, Ibadan, Nigeria

E-mail: fcnworgu@yahoo.com

Abstract

Nutritive potentials of *Centrosema pubescens*, *Mimosa invisa* and*Pueraria phaseoloides* leaf meals (LM) were determined and evaluated using different types of chicken within the years 2004-2009. The diets were formulated to contain 0,20/25, 30/40, 60/75 and 80/90g LM/kg feed and 120 to 150 birds were used in each experiment. Each treatment was replicated three times. All ingredients in each experiment were of constant weight, except the soybean and groundnut cake which test ingredients replaced some percentages weight for weight. The experimental design used was completely randomized design for the chicks and completely randomized block design for the growing pullets and cockerels. Data on feed intake and weight gain were subjected to one-way ANOVA and comparisons were made using Duncan's Multiple Range Test. Results revealed that the LM are rich in crude protein (21.36-23.34%) and ash (4.25 – 9.14%). The most available mineral elements were potassium (0.45-1.85%) and calcium (0.60 – 1.726%). The concentration of tannin was highest in the LM (1.57-3.35g/100gDM) unlike oxalate (0.037-0.065g/100gDM). The poorest LM in terms of nutritive value and chicken performance was *Mimosa invisa* leaf meal (MLM), while the best was *Centrosema pubescens* leaf meal (CLM). Inclusion of 40 – 60gCLM/kg feed for Black Nera chicks and finisher pullets and 75gCLM/kg feed for broiler starters and finishers resulted to significant (P<0.05) reduction in feed intake and weight gain. Feed intake and weight gain of broiler starters and finishers and cockerel growers fed 20 – 60gMLM/kg feed supplements significantly (P<0.05) and progressively decreased with increased dietary concentration of MLM supplement. Broiler starters and finishers fed 30-90PLM/kg feed had increased feed intake, which was progressive, unlike their weight gain which depressed significantly (P<0.05) with increased concentration of *Pueraria phaseoloids* leaf meal (PLM).Hence, 20 and 25gCLM/kg feed is recommended for pullet chicks/growing pullets and broiler chicken, respectively, while PLM and MLM are not recommended, for chicken nutrition due to depression in growth rate.

Keywords: Nutritive potential, legume leaf meal, growth responses, chicken.

UNDER PEER REVIEW

Introduction

The rising cost of poultry feeds has continued to be a major problem in poultry industry in developing countries, as feed cost is about 65 to 70% of the total cost of production (1) compared to 50 to 65% in the developed countries (2). High cost of conventional protein sources has necessitated the use of leaf meal supplements in poultry production. These leaf meals are readily available and cheaper than the conventional protein sources. Leaf meal supplements have been included into the diets of poultry as a means of reducing the high cost of conventional protein sources and to improve profit margin (3,4,5,6). The importance of legume leaf meals in poultry has been recognized by farmers because of their relatively high content of proteins, some minerals and vitamins (3,7). D'Mello (8) indicated that a distinguished feature of tropical legumes in comparism with their temperate counterparts is the occurrence of a diverse arry of non-protein amino acids in sufficient concentrations to precipitate adverse effects in farm animals.

Different leaf meals have been incorporated into poultry diets, incuding those of *Leucaena leucocephal* (9), *Cnidoscolus aconitifolius* (10), *Manihot esculenta* (11,12), *Centrosema pubescens* (13, 7), *Microdemis puberula* (14) among others. Onwudike (15) recommended dietary addition of 5% of *Leucaena leucocephala* leaf meal (LLLM) for improvement of body weight gain in chicks. Scott *et al. (16)* reported that dietary inclusion of LLLM more that (50g/Kg) caused reduced growth of broiler chicken and low egg production. Nworgu and Fapohunda (6) reported that 2.5% MLM reduced weight gain, while 7.5% significantly ($P<0.05$) reduced feed intake. Ngodigha (13) reported reduction in feed intake when broiler chickens were fed 15% *Centrosema pubescens* leaf meal (493.88g/bird) compared to 529.00, 532.51 and 516.40g/bird at 0.0, 5.0 and 10.0%, respectively. Hence, the objective of this study was to access the growth responses of chickens fed different leaf meal supplements.

Materials and Methods

The experiments were carried out at Bora Poualtry Unit of Federal College of animal Health and Production Technology, Institute of Agricultural Research and Training (IAR&T)

Ibadan, Nigeria. The experiments were carried out between the years 2004 to 2009 with the mean annual rainfall of 1398mm and mean monthly temperature of 22.71^{o}C and average monthly relative humidity of 75.54%. A total of 120 to 150 birds were

UNDER PEER REVIEW

used in each experiment. Tender leaves of *Centrosema pubescent, Mimosa* invisa and *Pueraria phaseoloides* were harvested from and around the paddocks in IAR&T. The leaves were detached from the vines/stems and were air and sun-dried for 2-3 days to a moisture content of about 12% (4). The dried leaves were milled/ground using a hammer mill with a sieve/screen size of 2.0-3.0mm. Samples of the leaf meals were collected for proximate and chemical analyses, while some were incorporated into the diets. Diets A,B,C, D and E contained 0,20/25, 30/40, 60/75 and 80/90g/kg feed of each forage meal, respectively. The diets were formulated according to NCR (18) and ARC (19) and ingredients were constant, except the test ingredient (the leaf meal), soybean meal and groundnut cake. The test ingredient replaced some percentages of soybean and groundnut cake in each experiment weight for weight. Gross composition of control diets is presented in Table 1. The birds were randomly allotted to the aforementioned dietary treatments in each experiment. Each treatment was replicated three times in a completely randomized design, but for growing pullets/pullet finishers and cockerel finishers, completely randomized block design was used. Data on feed intake were taken on daily basis, while weight gains were taken on weekly basis. Routine management practices and medication were taken as and when due.

Proximate and mineral composition of the test ingredients were determined by the procedures of Boehringer (20) and proximate composition of the diets was determined by the methods of AOAC (21), while their metabolisable energy (ME) was determined by the methods of outlined by Carpenter and Clegg (22). Gross energy of the test ingredients was estimated by the method of AFRC (23). Phytate was determined by the technique of Igbedioh *et al*. (24), while tannin was evaluated by the method outlined Hagerman and Ler (25) and oxalate by the procedures of Talapatra and Prince (26).

Data on feed intake feed, weight gain and feed conversion ratio were subjected to one-way ANOVA analysis of variance procedure of SAS (27). Duncan's Multiple Range Test (28) was used in assessing the significant differences among the treatments.

UNDER PEER REVIEW

Results

The results revealed that the leaf meals (LM) studied are rich in crude protein (CP). The crude fibre (CF) contents of these LM are moderate, most especially for C. *pubescent* leaf meal (CLM), while *P. phaseoloides* leaf meal (PLM) is very high in CF (Table 2). The CLM and PLM are rich in ash, while M.*invisa* leaf meal (MLM) is very low in ash. The CLM and MLM are rich in calcium, potassium and phosphorus. The PLM had the least calcium and potassium. The LM have moderate concentrations of magnesium and sodium, while the concentration of phosphorus in the PLM is the lowest. The concentrations of zinc and iron of the LM are moderate. The concentrations of oxalate, phytate and tannin in these LM are presented in Table 2. The most available anti-nutritional factor was tannin.

Feed intake, weight gain and feed conversion ratios of Black Nera chicks, Black Nera growing pullets, broiler starters and finishers fed graded levels of *Centrosema pubescens* leaf meal were significant ($P<0.05$) (Table 3). Dietary inclusion of 2 to 6% CLM supplements for Black Nera chicks resulted to increased feed intake and depression in both weight gain and feed conversion ratio compared to the birds placed on control diet. However, the best performance for Black Nera growing pullets in terms of weight gain was observed on the birds placed on 2% CLM, which also had elevated feed intake. Dietary inclusion of 2.5 to 5.0% CLM for broiler starters and finishers resulted to elevated feed intake which depressed at 7.5% (Table 3).

Inclusion of more than 20g/kg *Mimosa invisa* leaf meal to the diets of cockerel chicks resulted to lower feed intake and depressed weight gain and feed conversion ratio (Table 4). Similar scenario was observed for cockerel growers fed graded levels of *Mimosa invisa*leaf meal. Dietary inclusion o*f Mimosa invasa* leaf meal in broiler starters and finishers diets resulted to significant ($P<0.05$) reduction in feed, weight gain and feed conversion ratio, which were progressive. Broiler starters and finisher fed graded levels of *Pueraria phaseoloides* leaf meal had reduced weight gain and feed conversion ratio which were significant ($P<0.05$) and progressive compared to control (Table 5).

Discussion

The leaf meals (LM) studied are moderate in crude fibre (CF), except PLM. The relatively high CP and low CF with high concentration of ash will necessitate the use of CLM and PLM as protein and mineral supplements in chicken production. Aletor

UNDER PEER REVIEW

147 and Omodara (29) revealed that some tropical legumes have higher nutritive value
148 and that the availability of the component amino acids in the small intestine would be
149 more when the leguimous forage materials are fed dried than when fresh. The CP
150 value of CLM in this study is higher than the report of Aletor and Omodara (29), but
151 similar to that of Nworgu *et al*. (30) (19.6%) and Raharjo *et al*. (31) (21.4%). Wilson
152 and Lansbury (32) reported that *C.pubescens* contained 20.0% CP, with CF of 30.0%,
153 while Nworgu and Ajayi (33) noted that *C. pubescens* harvested at 12 weeks after
154 planting contained 18.70% CP, 11.80%CF,6.98% ash and other extract of 4.42%.
155 Dougall and Bogdan (34) highlighted that *M. invisa* leaves contained 16.2 and 37.2%
156 of CP and CF, respectively. The gross energy (GE) of the LM was 4354, 4419 and
157 4628Kcal/kg, for CLM, PLM and MLM, respectively. These values are in harmony
158 with the reports of Skerman *et al. (36)* who reported that tropical legumes have GE of
159 *4326-4802*KcalKg. Raharjo *et al*. (31) reported that GE of CLM was 3885Kcal/kg,
160 while that of PLM was 3872Kcal/kg. The variations in the CP,CF GE and EE could
161 be attributed to the age of cutting, climatic conditions, methods of processing and
162 analyses. Concentrations of calcium and potassium which were highest in the CLM
163 and MLM are in line with the submission of Aletor and Omodara (29) who stated that
164 calcium, potassium and sodium were the most available mineral elements in forage
165 meals. Calcium and phosphorus concentrations for CLM in this study are higher than
166 that reported by Raharjo *et al.* (31) whose value were 0.72 and 0.23% for calcium and
167 phosphorus, respectively, but they are in harmony with the reports of Gohi (37) and
168 Leche *et al*. (38) who reported 0.78 – 1.36% for calcium and 0.13-0.45% for
169 phosphorus. Neme and Nery (39) reported that MLM contained 0.14% phosphorus
170 which is lower than reported in this study. The report of Dougall and Bogdan (34) on
171 the calcium content of MLM (1.10%) is slightly lower than reported here (1.26%).
172 The calcium content of PLM in this study is relatively lower than the results of Neme
173 and Nery (43), Skerman *et al*. (39) and Lowry *et al*. (40) (0.36 – 1.72%). The values
174 of phosphorus and magnesium reported in this study are lower than the values of
175 Lowry *et al.* (40).
176 The diets for the chickens met the nutritional requirements of the birds are in line with
177 standards of NRC(18). Dietary inclusion of *Centrosema pubscens* meal in the diets of
178 broilers, cockerel chicks/finishers and pullet chicks/growers significantly ($P<0.05$)
179 resulted to reduced feed intake unlike the broilers fed graded levels of *Pueraria*
180 *phaseoloides* leaf meal (PLM). At both phases, feed intake of broilers fed 30-

UNDER PEER REVIEW

90PLM/kg feed increased with increased dietary inclusion of PLM supplements. This observation for the broilers fed PLM supplements is contrary to the submission of Nworgu (7), Esonu *et al.* (14), Udedibe and Opara (46). The increased feed intake for the broiler fed PLM supplements is associated with the lightweight of the PLM leaf meal and its lower energy. However, the weight gain of the broilers fed PLM supplements significantly ($P<0.05$) and progressively decreased with increased dietary concentrions of PLM. This observation is not in line with the reports of Ngodigha (13), Omeje *et al*. (42) and Ogbonna and oredein (45).

Dietary inclusion of 20-60kg *Mimosa invisa* leaf meal (MLM) in cockerel finishers, broiler starters and finishers resulted to depressed weight gain and feed conversion ratio which were significant ($P<0.05$) and progressive. Poor performance of these birds is attributed to the presence of mimosine in the MLM (44), low acceptability (36) and poor digestibility (8). These results are in harmony with the reports of Udebibie and Opara (46), but contradict, the submission of Ogbonna and Oredein (45).

Black Nera pullets tolerated the dietary inclusion of CLM in their diets more than the Black Nera chicks, as they had improved growth rate. Inclusion of 2-6g CLM/kg feed in the diets of the Black Nera chicks resulted to 15.93 – 30.90% depression in growth over control. This result is contrary to the report of Omeje *et al* (42) who concluded that 50-100gCLM/kg feed resulted to elevation of weight gain. However, this result agrees with earlier observations that dietary inclusion of leaf meal of *L. leucocephala,* (4and 8) and Alchornia *cordifolic* (46) depressed feed intake, weight gain and feed conversion ratio. Dietary inclusion of 25g CLM/kg feed for broiler starters resulted to 26.34 and 9.60% increase in weight gain over control and 75g CLM/kg feed, respectively. However, for broiler finishers, the increase in weight gain for 25gCLM/kg feed was 56.12% due to prolonged feeding, as the feeding was continuous from starter to finisher phase (i.e. 7 weeks). These results are contrary to the reports of Omeje *et al*. (42), but agree with the submission of Odensi *et al*. (5) when broiler chicks were fed graded leaves of *Tithonia diversifolia* and Daghir (17) who recommended 2% of *L. leucocephala* leaf meal for broiler starters and 3% for chick starters. Donkoh *et al*. (10) recommended 25g *Cindoscolus aconitifolius* leaf meal per kilogramme of feed for optimal chickens performances, while Nworgu and Fapohunda (12) Nworgu *et al*. (43) concluded that 25gMLM/kg feed was adequate for broiler chicks and finisher maximum performance.

UNDER PEER REVIEW

Conclusion

The leaf meal (LM) studied are rich in crude protein (21.36 – 23.24%) and ash (4.25-9.14%). The most available mineral elements were potassium (0.45-1.8%) and calcium (0.60-1.26%). *Centrosema pubescens* leaf meal (CLM) is the most outstaniding in mineral profile, nutritive value and chickens performance, while *Mimosa invisa* leaf meal (MLM) is the poorest. Broiler starters/finishers, cockerel chicks/growers and pullet chicks/finishers tolerated the CLM, CLM and PLM supplements differently. Dietary inclusion of 40-75g CLM/kg feed and 30-60 gMLM/kg feed resulted to depressed feed intake in the chickens, unlike the inclusion of 30-90PLM/kg feed which improved feed intake. Dietary inclusion of 75g CLM/kg, 30-60MLM/kg and 30 – 90PLM/kg feed resulted to depressed growth performance. For improved performance and adequate growth rate, it is advisable to include 20gCLM/kg feed in the diets of pullet chicks and growing pullets, while 25g CLM/kg feed is adequate for broiler chicks and broiler finishers. However, it is not advisable to include MLM and PLM in the diets of pullet chicks/growers, cockerel chicks/finishers and broiler chicks/finishers due to poor growth rate.

UNDER PEER REVIEW

249 **Table 1: Gross composition of control diets**

Ingredients%	Pullet Chicks (0-6wks)	Growing pullets (8-16wks)	Broiler chicks (1-5wks)	Broiler finishers (5-8wks)	Cockerel chicks (0-9wks	Cockerel growers (10-18wks)	Broiler chicks (0-4wks)	Broiler finishers (4-8wks)	Broiler starters (0-4kws)	Broiler finishers (4-8wks)
	Centrosema pubescens			*Mimosa invasa*		*Pureria phaseoloides*				
Maize	31.6	20.3	48.0	51.5	30.0	24.3	44.0	45.3	46.8	48.8
Corn bran	21.0	23.4	14.0	17.2	26.6	22.0	13.0	13.8	10.0	11.0
Palm kernel cake	12.0	21.0	-	-	9.0	20.4	-	-	5.2	5.2
Coca pod husk	8.5	9.5	-	-	10.0	8.0	5.0	5.0	-	-
Soybean meal	14.6	11.0	21.2	17.5	13.0	11.1	20.6	20.0	20.4	19.4
Groundnut Cake	6.5	6.5	9.5	7.0	4.0	6.4	10.0	9.0	10.3	9.3
Leaf meal	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Fish meal	2.0	3.5	4.0	3.0	4.2	3.5	4.0	3.5	4.0	3.0
Bone meal	2.0	2.5	2.5	3.0	2.3	3.5	2.6	2.6	2.5	2.5
Oyster shell	1.0	1.5	-	-	-	-	-	-	-	-
Salt	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Vitamin-mineral premix*,+	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.3	0.3	0.3
Lysine	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Methionine	0.1	0.1	0.1	0.1	0.2	0.2	0.1	0.1	0.1	0.1
Calculated analysis										
Crude protein(%)	19.0	18.9	22.0	18.8	18.7	20.3	22.0	21.2	22.6	21.3
Crude fibre(%)	6.9	8.8	4.5	4.4	6.8	7.6	5.4	5.4	4.3	4.4
Metabolisable energy (Kcal/kg)	2640	2526	2936	2966	2646	2436	2882	2889	2909	2932
Determined analysis										
Crude protein (%)	20.5	20.7	22.6	20.1	19.2	18.9	22.2	20.0	23.6	23.6
Crude fibre (%)	8.2	6.9	4.6	4.6	6.8	7.6	5.0	4.6	3.8	3.9
Metabolisable energy (Kcal/kg)	2740	2700	3100	3195	2700	2500	3001	3004	3000	3012

250 * Vitamin-mineral premix provides per kg of diet: Vit A=10000in, Vit D3= 1500iu;Vit E=3iu,Vit
251 B12=0.08mg;Vit.K=2mg;Ribofl=3mgProduction acid=6mg; Niacin=15mg; chlorine=3mg;Folic acid=4mg,
252 Mn=8mg,Zn=0.5mg;iodine=1.0mg;Co=1.2mg;Cu=10mg and Fe=20mg.
253 + Vit.-mineral premix provides per kg diet: Vit A=12,500,000iu, Vit D3=2,500,000iu; Vit E=25,000iu; Vit.K = 2.00mg; Vit.
254 B1=2.00mg; Cu=6.00mg; Fe=40.00mg; iodine=1.00mg; Cobalt-0.2mg; Selenium =0.15mg; Choline chloride = 500.00mg.
255 • For cockerels and pullet chicks/growers
256 + For broiler starters/finishers.
257
258
259
260
261

UNDER PEER REVIEW

262 **Table 2: Proximate chemical composition of experimental test ingredients (% DM Basis)**
263

	Experiment Test Ingredients		
Fraction	***Centrosema* leaf meal**	***Mimosa* leaf meal**	***Pueraria Ileaf meal***
Dry mater	**88.99**	**89.99**	**89.90**
Crude protein	**23.24**	**23.34**	**21.36**
Crude fibre	**8.80**	**11.29**	**21.52**
Ether extract	**3.32**	**2.38**	**3.20**
Ash	**9.14**	**4.25**	**8.00**
Nitrogen free extract (NFE)	**55.50**	**58.74**	**45.92**
Calcium	**1.22**	**1.26**	**0.60**
Potassium	**1.85**	**1.60**	**0.45**
Phosphorus	**0.54**	**0.38**	**0.12**
Magnesium	**0.30**	**0.24**	**0.28**
Sodium	**0.17**	**0.15**	**0.11**
Zinc (mg/kg/DM)	**40.01**	**40.00**	**20.00**
Iron (mg/kgDM)	**20.01**	**10.11**	**80.10**
Oxalate (g/100g DM)	**0.037**	**0.065**	**0.060**
Phytate (g/100g DM)	**0.48**	**0.37**	**0.32**
Tannin (g/100g DM)	**2.81**	**1.57**	**3.35**
Gross energy (kcal/kg)*	**4354**	**4628**	**4419**

- **Estimated by AFRC**

264 **Table 3: Feed intake and growth responses of chickens fed graded levels of *Centrosema***
265 ***pubescens* leaf meal**

Dietary level of leaf meal (g/kg)									
Source of leaf meal	**Type of chicken**	**0**	**20**	**25**	**40**	**50**	**60**	**75**	**SEM**
		Feed intake and weight gain (g/bird)							
Centrosema Pubescens	Black Nera chicks (0-6 wks)	964^{d} 339^{a}	1010^{a} 285^{b}	-	984^{c} 234^{d}	-	998^{b} 248^{c}	-	3.3 4.1
	Black Nera growing pullets/finisher pullet (8-16wks)	6469^{b} 699^{c}	6658^{a} 740^{a}	-	6093^{d} 719^{b}	-	6199c 734ab	-	1.8 5.8
	Broiler chicks/starters (1-5 wks)	2300^{b} 1009^{c}	-	2420^{a} 1369a	-	2410^{a} 1285^{ab}	-	2260^{b} 1238^{b}	12.2 27.7
	Broiler Finishers (5-8 weeks)	3210a 792^{b}	-	3100^{a} 1001^{a}	-	3200^{a} 985^{a}	-	2860^{b} 562^{c}	30.9 24.0
Feed conversion ratio									
	Black Nera chicks (0-6wks)	2.8^{c}	3.5^{b}	-	4.2^{a}	-	4.0^{a}	-	0.11
	Black Nera growing pullets/finisher pullets (8-16wks)	9.3^{a}	9.0^{ab}	-	8.5^{b}	-	8.4^{b}	-	0.51
	Broiler chicks/starter (1-5 wks)	2.3^{a}	-	1.8^{b}	-	1.9^{b}	-	1.8^{b}	0.07
	Broiler finisher s (5-8 wks)	4.16^{b}	-	3.19^{d}	-	3.3^{c}	-	5.1^{a}	0.17

266 abcd: Means with different superscripts on the same horizontal row differ
267 significantly(P<0.05),WKS=weeks Numerator = Feed intake; Denominator = Weight gain
268

UNDER PEER REVIEW

269 **Table 4: Feed intake and growth responses of chickens fed graded levels of**
270 ***Mimosa invisa* leaf meal**

Source of leaf meal	Type of chicken	Dietary level of leaf meal (g/kg) 0	20	30	40	60	SEM
		Feed intake and weigh gain (g/kg)					
Mimosa invisa	Cockerel chicks (0-9wks)	2908[c] / 517[d]	3142[a] / 644[a]	-	2706[d] / 553[c]	3082[b] / 588[b]	3.0 / 3.5
	Cockerel growers/ finisher (10-18weeks)	8800[a] / 1058[a]	-	8783[b] / 853[b]	-	8166[c] / 820[c]	3.3 / 15.7
	Broiler chicks (0-4 wks)	1282[a] / 538[a]	1235[b] / 504[b]	-	1174[d] / 505[b]	1204[c] / 403[bc]	1.6 / 5.5
	Broiler finisher g/bird (4-8 wks)	3741[a] / 659[a]	3642[b] / 413[b]	-	3371[d] / 400[c]	-	2.0 / 3.0
		Feed conversion ratio					
	Cockerel chicks (0-9wks)	5.6[a]	4.9[c]	-	5.1[b]	5.2[b]	0.09
	Broiler chicks (0-4 wks)	8.3[b]	-	10.2[a]	-	10.0	0.82
	Broiler finisher (4-8wks)	2.4[b]	2.5[b]	-	2.3[b]	3.0[a]	0.03
	Broiler finisher (0-4 wks)	5.9[a]	8.8[b]	-	8.4[b]	8.6[b]	0.05

271 **Abcd: means with different superscripts on the same horizontal row differ significantly ($p<0.05$);**

272 **WKS = Weeks ; Numerator = Feed in take; Denomminator = Weight gain**

UNDER PEER REVIEW

Table 5: Feed intake and growth responses of chickens fed graded levels of *Pueraria phaseoloides* leaf meal.

Source of Lead meal	Type of chicken	Dietary Level of Leaf Meal (g/kg) 0	30	60	90	SEM
		Feed intake and weight gain				
Pueraria phaseloides	**Broiler starter(0-4 wks)**	1202[b] / 536[a]	1122[b] / 479[b]	1234[a] / 453[c]	1246[a] / 399[d]	0.7 / 0.7
	Broiler finisher (4-8 wks)	4306[d] / 1466[a]	4387[c] / 1370[b]	4492[a] / 1354[c]	4446[b] / 1341[d]	0.7 / 0.6
		Feed conversion ratio				
	Broiler starter (0-4 wks)	**2.2[c]**	**2.3[c]**	**2.7[b]**	**3.1[a]**	**0.22**
	Broiler finisher (4-8 wks)	**2.9[b]**	**3.2[a]**	**3.3[a]**	**3.3[a]**	**0.24**

abcd: Means with different superscripts on the same horizontal row differ significantly (P<0.05)

Numerator = Feed intake; Denominator = Weight gain; WKS = Weeks

References

1. AFRC., 1990. Agricultural and Food Research Council: 750.

2. AOAC., 1990. Association of Official Analyatical Chemists. Official Methods of Analyses. 15th Edition, Washington, D.C.

3. ARC., 1995. Agricultural Research Council. In: World Poultry Journal 57:225-243. The nutrient requirements of farm livestock, poultry, 2nd Edition (London, Agricultural Research Council).

4. Aletor V.A. and Omodara O.A., 1994. Studies on some leguminous browse plant, with particular reference to their proximate, mineral and some endogenous anti-nutritional constituents. Animal Feed Science and Technology 46:343-348.

5. Boehringer., 1979. Mannheim GmbH Diagnostica 6800. Mannheim Germany, Cat. No. 124974.

UNDER PEER REVIEW

6. Carpenter K.J. and Clegg K.M., 1956. The metabolisable energy of poultry feed stuff in relation to their chemical composition. Journal Science, Food and Agriculture 7:45-48.

7. Daghir N.J., 1995. Feedstuffs used in hot regions. In: Poultry production in hot climates. Daghir, N.J. (ed): 1st Edition CAB International Wallingford UK: 125-154.

8. Donkoh A., Atuahene C.C., Poku-Prempeh Y.B. and Twnm I.G., 1999. The Nutritive value of chaya leaf meal (*Cnidoscolus aconitifolius* (Mill) Johnston): Studies with broiler chickens. Animal Feed Science and Technology (Netherlands) 77 (1-2):163 – 172.

9. Dougall H.W. and Bogdan A.V., 1966. The chemical composition of some leguminous plants grown in the herbage nursery at Kitale, Kenya. East African Agricultural and Forestry Journal 32:45-49.

10. D'Mello J.P.E., 1995. Leguminous leaf meals in non-ruminant nutrition. In: Tropical legumes in animal nutrition. 1st Edition. A.B. International, Wallingfort, Oxon, U.K.247-280.

11. D'Mello J.P.F., 1992. Nutritional Potentialities of fodder trees and fodder shrubs as protein sources in mongastric nutrition. Legume trees and other fodder trees as protein sources for livestock. Food and Agriculture Organization, Rome. 1st Edition. Speedy, A and Puglies, P.H. (eds.) :115-127.

12. Esonu B.O., Iheukwumere F.C., Iwuji T.C., Akanu N and Nwugo O.H., 2003. Evaluation of *Microdesmis puberula* Leaf meal as feed ingredient in broiler starter diets. Nigerian Journal of Animal Production 30(1): 3-8.

13. FAO., 1992. Legume tree and other fodder trees as protein sources for livestock. FAO Animal Production and Health Paper 102: 14-18.

14. Gohi B., 1981. Tropical feeds FAO Animal Production and Health Series No. 12: 529.

15. Hagerman A.E. and Ler I.G., 1983. Precipitation method for the qualitative determination of tannins. Journal of Agriculture, Food and Chemistry 26: 809-812.

16. Igbedioh S.O., Olugbemi K.T.A. and Akpapunam M.A., 1994. Effect of processing methods on phytic acid level in bambara nut and pigeon pea. Chemistry 50:147-151.

17. Leche T.I., Groenendyk G.H., Westwood N.H. and Jones M.W., 1992. Composition of animal feedstuffs in Australia. 1st Edition. Division of Animal Production, CSIRO: 78-84.

UNDER PEER REVIEW

18. Lowry B.J., Petheram J.R. and Tangendjaja B., 1992. Plants fed to village ruminants in Indonesia, Canberra. aciartechnical reports, no. 22:60.

19. McDonald P., Edward R.A. and Greenhalgh J.P.D., 1994, Evaluation of food. In:Animal nutrition. 4th edition. Longman Scientists and Technical, England: 260-283.

20. **Natanman R. and Chandrasekaran D., 1966**. Subabul leaf meal (*Leucaena leucocephala)* as a protein supplement for broilers. Indian Vet. Journal 73 (10): 1042 – 1042.

21. Neme A.A. and Nery J.R., 1965, Influence of mineral fertilizers and lime on the yield and chemical composition of perennial forage legumes. Proceedings 9th Grassl, Congr. Sao Paulo, Brazil: 655-670.

22. Ngodigha E.M., 1994. Incorporation of *Centrosema pubescens* in broiler diets. Effects on performance characteristics. Bulletin Animal Health Africa 42:159-161.

23. Nworgu F.C., Adebowale E.A., Oredein O.A. and Oni A., 1999, Prospect and economics of broiler production using two plant protein sources. Tropical Journal of Animal Science 2 (1): 159-166.

24. Nworgu F.C. and Faphunda J.B., 2002, Performance of broiler chicks fed mimosa (*Mimosa invisa)* leaf meal supplements.Proceeding of 27th Ann. Conf. of Nigerian Society of Animal Production (NSAP), Held at the Federal University of Technology Akure, Nigeria on the 17th – 21st March, 2002. Aletor V.A. and Onibi G.E. (eds.): 128 – 131.

25. Nworgu F.C., 2004. Utilization of forage meal supplement in broiler production. Ph.D. Thesis University of Ibadan, Ibadan – Nigeria: 136-146.

26. Nworgu F.C., Egbunike G.N. and Osayomi O.J., 2001. Performance of growingrabbits fed a mixture of leaf meals and concentrates. Tropical Animal production Investigation 4:37-48.

27. Nworgu F.C. and Ajayi F.T., 2005, Biomass, dry matter yield, proximate and mineral composition of forage legumes grown as early season feeds. Livestock Research for Rural Development (17): http//www.utafoundation.org/irrd1711/nwor/7121.htm

28. Nworgu F.C., Egbunike G.N., Ononogbu C.E., Fapohunda J.B. and Ogbonna J.U., 2003. Effect of mimosa (*Mimosa invisa)* leaf meal supplements on broiler finishers performance. Proceedings of the 8th Annual Conf. of Animal Science Association of Nigeria. Held at Federal University of Technology, Minna, Niger

UNDER PEER REVIEW

State on September 16 – 18th. Olatunji, E.A., Ayanwale B.A., Shiawoya Ed. And Aremu A (eds): 36-39.

29. Nworgu F.C. and Egbunike G.N., 1999. Performances characteristics of broilers fed diets with soyabean meal and groundnut cake as protein sources. Tropical Animal Production Investigation 2 (1): 51-59.

30. Odunsi A.A., Farinu G.O., Akinola J.O and Togan V.A., 1999, Growth, carcass characteristics and body composition of broiler chicks fed wild sunflower (*Tithonia diversifolia)* forage meal. Tropical Animal Production investigation 2:205 – 211.

31. Ogbonna J.U. and Oredein A.O., 1998. Growth Performance of cockerel chicks fed cassava leaf meal. Nigeria Journal of Animal Production 25(2): 129-133.

32. Ogbonna J. U. and Adebowale, E.A., 1993. Effects of sun-dried cassava peel meal asreplacement for maize and wheat offals on performance and nutrient utilization of

cockerels. Nigerian Journal of Animal Production 20(1&2): 61-70

33. Onwudike O.C., 1995. Use of the legume tree crops: *Gliricidia sepium and Leucaena leucocephala* as green feeds for growing rabbits. Animal Feed Science and Technology 51:153-163.

34. Omeje S.I., Odo B.I. and Egwu P.O., 1997. The effects of *Centrosema pubesecens* asleaf meal concentrate for broiler birds. In Book of Abstract of the 22nd Annual Conference of NSAP, held at Abubakar Tafawa Balewa University, Bauchi, Nigeria: 136-137.

35.S.A., Opoku R.S. and Atuahene C., 1990. *Gliricidia speium* leaf meal as an ingredient in layers diets. Animal Feed Science and Technology 29:302-308.

36. Raharjo Y.C., Cheeke P.R., Patton N.M. and Supriyati K. 1986. Evaluation of tropical forages and by-product feeds for rabbit production. Nutrient digestibility and effects of heat treatment. Journal of Applied Rabbit Research 9:56-66.

37. Reyes B., 1955. The digestibility of *Centrosema pubescens* and *Purearia javanica.* Philiphine J. Agric. 30:27-27.

38. Ross E. and Enriquez F.O., 1969. The nutritive value of cassava leaf meal. Poult. Sci. 48:846-853.

39. SAS., 1989. Statistical Analysis System, Institute Inc. SAS/START user's guide version. 6th Edition, SAS Inst. Inc. Cary. NC. 846.

UNDER PEER REVIEW

40. Scott M.L., Nesheim M.C. and Young R.J., 1982. Nutrition of chicken. Scott, M.L. and Ithaca, N.Y. (eds.). 1st Edition: 25-50.

41. Steel R.G. and Torrie J.H., 1980. Principles and procedures of statistics. A biometrical approach. 2nd Edition, McGraw Hill Book Co. Inc., New York.

42. Silva H.O., Fonseca R.A-da, Guedes-Filho-R-des and da – Fonseca R.A., 2000. Productive traits and digestibility of cassava leaf meal in broiler diets without addition of enzymes. Revista-Brasillerra-de Zootecnia. 29(3): 823-829.

43. Skerman P.J., Cameroon D.G. and Riveros F., 1988. *Centrosema spp.* In: Tropical Forage legumes. 2nd Edition, FAO Plant production and protection Series, No. 2. Rome – Italy. Skerman (ed.): 237-257.

44. Talapatra E.L. and Price B.L., 1948. Oxalate: Its chemistry and nutritional significance. World Review of Nutrition and Dietetics, Krager, New York: 51-55.

45. Tackie A.M. and Flenscher J.E., 1995, Nutritive value of wild sorghum fortified with *Leucaena leucocephala* (Wh.Lam). Bulletin Animal Health Africa 47: 223-275.

46. Topps J.H., 1992. Potential composition and use of legume shrubs and trees as fodder for livestock in the tropics. Tropical Journal of Agricultural Science Cambridge 118:1-8.

47. Udedibie A.B.I. and Opara C.C. 1998. Responses of growing broilers and laying hens to the dietary inclusion of leaf meal from *Alchornia cordifolia.* Animal Feed Science and Technology 71: 157 – 164.

48. Wilson A.S.B. and Lansbury T.J., 1958. *Centrosema Pubescens:* Ground cover and forage crops in cleared rainforest of Ghana. Empire Journal of Experimental Agric. 26: 351-364.

UNDER PEER REVIEW