

NEEM (*Azadirachta indica*) SEED CAKE/KERNEL AS PROTEIN SOURCE IN RUMINANTS FEED

A. Aruwayo^{1*} and S. A. Maigandi²

1. Department of Animal Production and Health, Faculty of Agriculture and Agricultural Technology, Federal University, Dutsinma, Katsina State, Nigeria.

2. Department of Animal Science, Faculty of Agriculture, Usmanu Danfodiyo University, Sokoto, Nigeria.

E-mail: aaruwayo@fudutsinma.edu.ng

ABSTRACT

Ruminant production that is very popular in Nigeria livestock production faces series of problems, chief among them is the shortage of protein supply in adequate amount for optimum performance of the animals. The forages, which constitute the bulk of the source of feed for these animals, fluctuate in supply with poor nutrient content especially during the dry season of the year. The convectional protein concentrates which boost the protein supply faces intense competition from man. It has become imperative for intensive effort to be made at making the unconventional sources of protein available to the farmers. Neem seed cake/ kernel cake fits as one of the unconventional protein supply in view of the high level of the protein content and balancing of the amino acids. It is capable of offering a big relief to these protein shortages. The neem seed is readily available in the northern part of Nigeria that is the home to the largest of number of ruminants in the country. However, they are not readily acceptable to the animals due its pungent smell and bitterness caused by the bioactive principles present. This paper reviewed availability of neem seed in Nigeria, the antinutritional and other bioactive factors inherent in the cake as well as the methods of processing and removing them. The review also highlighted the nutritional quality of the cake, effect on haematological and biochemical characteristics; and the economics of use of the cake. The paper concluded that neem seed cake/kernel could readily serve as safe source of protein to the ruminants with appropriate treatment to remove active principles.

Keywords: Neem kernel cake, Alkali treated neem kernel, haematological, biochemical, Uda lambs and Uda Rams

1. INTRODUCTION

Ruminant production in Nigeria is popular among the rural farmers. They are widely distributed in among rural, urban, and peri-urban areas representing about 63.7% of the total grazing domestic animals in Nigeria (1). Ruminants account for major supply of national meat supply (2). However, the level of domestic livestock production falls short of demand (3). The average Nigerian consumes only 3.245g of animal protein per day, out of the 34g recommended (4). This is grossly inadequate for normal body activities. There have been resultant effects on the health of the citizens and their productivity. The most important factor responsible for the gap in the relationship between the demand and supply of ruminants in Nigeria is the supply of feed both in quantity and quality.

Comment [A1]: This is a quite strong affirmation and I would suggest you to remove it. You may experience problems with vegetarians for example.

The main feed resources for small ruminants are natural pastures consisting of legumes and browse species (5). These pastures depend on rainfall, which fluctuates especially in the northern part of the country where the largest percentage of the animals are raised. The scarcity of energy and protein feedstuffs during dry the season is a major setback to ruminant livestock production in the tropics (6). The increase in human population further worsens the situation because of the pressure placed on the available land for grazing by other agricultural and non agricultural activities. During this period, the available forages are dry, protein content of which is very low and there is marked decrease in voluntary intake and digestibility by the animal (7; 8). The nutritional problems of ruminants have been increased by competition between man and the animals for the scarce grains and the protein concentrates feed making it difficult to meet up with nutritional requirements of the animals at affordable cost. Other factors which have contributed to the increasing cost of feed are under-production of various ingredients used in feed formulation and high inflation rates.

The commonest protein supplement for livestock feed in Nigeria in periods of low yield and availability of poor quality herbage are groundnut cake (GNC) and cotton seed cake (CSC). Groundnut cake (GNC) is the major plant protein supplement used in Nigeria. Its availability depends on the production level of groundnut. The unextracted groundnut has about 26 - 30% protein and it mainly consumed by humans while Groundnut cake has a protein content of about 38 - 47% (9) and 40-45% (10). It is palatable but however, deficient in lysine, methionine and threonine but rich in vitamin E, K, B, thianine and niacin (11). Cotton seed cake (CSC) is obtained from cotton after the removal of the lint, followed by oil extraction from the seed. It has a protein content of 38-44% (12) and 41-45% depending on the efficiency of oil extraction but deficient in lysine, methionine, leucine and isoleucine (13).

The prices of GNC and CSC products have been rising thereby increasing the cost of production. Researchers therefore considered the use of alternative sources of feed ingredients in order to reduce the cost of production. One of the promising material is the neem(*Azardirachta indica*) kernel cake. Neem (*Azardirachta indica*) seed cake is obtained from neem seed. Information available on Neem plant (*Azardirachta indica*) indicated that it is the only species in the genus *Azardirachta*. It is native to India and Burma, growing in tropical and mitropical regions (13). A fullgrown tree can produce 30-100 kg fruit depending on rainfall, soil type and ecotype. 50 kg of fruits yield 30 kg of seed giving 6 kg of oil and 24 kg of seed cake (13). It has several uses and can

Comment [A2]: this information comprehend the next information.

Comment [A3]: It seems obvious!

Comment [A4]: ???

be grown under reforestation and social foresting programmes. The neem tree goes prefers well drained deep soil, sandy loams with ground water at 3.048 metremeter dept or more. It takes about five years to produce the first fruit crop but can produce a good yield in the third year (14). The tree has the advantage of growing on marginal lands, where it will not compete with food crops (15).

Over four millions of neem trees have been planted in the northern part of Nigeria especially in Borno, Kano, Katsina and Sokoto states and rich in protein (16). However, dDespite its availability and the derivable benefits from the use of the cake, its use is still beleaguered by series of problems. So, the

The main objective of this study is to create awareness on the potentials of the use of neem seed cake in Nigeria. The study dwelt on the nutritional and antinutritional components; and the different methods of removing the bioactives in the cake. This was done with data obtained from reputable peer reviewed journals, published and unpublished reports on the use of neem seed cake/kernel cake as an unconventional sources of protein in ruminants.

DISCUSSION

2.1 Neem (*Azadirachta indica*) Tree Distribution and Seed Availability

Neem tree is planted in many states in Nigeria (17). Over four millions of neem trees have been planted in the northern part of Nigeria especially in Borno, Kano, Katsina and Sokoto states (16). About 3,500 hectares of land is under cultivation in Kebbi, Sokoto and Zamfara states, with a density of about 1,200 trees per hectare (18). In Nigeria, neem form about 90% of forestry established in 12 states within the savanna zone under the aforestation programme (19). The neem tree does well in well drained deep soil, sandy loams with ground water level at 3.05 metre depth or more. The Neem tree (*Azadirachta indica*) is an evergreen of the tropics and subtropics. It is native to India but widely planted and naturalized throughout Asia, Africa, Australia, the Caribbean and several Central American countries (20). It may grow up to 15m tall under ideal conditions and is reported to live up to 200 years (20). The tree has the ability to withstand frost due to a very suberised back layer and can easily survive desert conditions of temperature above 50°C and takes about five years to produce the first fruit crop but can produce a good yield in the third year (14). Neem tree can perform well in dry soils that are poor in nutrients and have low rainfall per annum requirement (130 mm) and the fruit yield is extremely variable ranging from 10 to 50kg per tree with an average of 20kg (21). Neem tree flowers between February and May with

Formatted: Indent: First line: 0.5", Space After: 10 pt, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Comment [A5]: This information is in the lines 67 and 71.

Comment [A6]: This information is in line 72. So you have to decide because similar information was given in introduction and discussion. Maybe in introduction you can inform about neem origin, countries and conditions to grow up and in the discussion information about seed production, flowers and fruits.

profuse clusters of small white flowers. The fruits are drupes, turning golden yellow on maturity, which occurs during June to August in India. The tree begins bearing fruit after 3 to 5 years and produces about 50kg fruits annually when mature in India. The seeds are about 2cm long and 1.5cm diameter, kernel of about 1.5cm length containing about 30 to 40 percent oil and 5 to 6 years old tree can yield 20kg kernels (14). The yield of 20.5kg fresh fruit for a fully grown neem tree was reported in Nigeria (17) and an average yield of 17.71kg in Nigeria (Sokoto and its environ) (22). The large number of the neem trees and its yield in the northern part of Nigeria is an advantage for its use.

2.2 Anti-nutritional Factors of Neem Seed Cake as Feed Ingredient

The possibility of using neem seed cake (NSC) in livestock ration was explored by (23) using the feeding practice of local farmers in Southern India. However, NSC pungent odour and the bitter taste caused by the active principles isolated from different parts of the plant namely azadirachtin, meliacin, gedunin, salanin, nimbin, valassin and many other derivatives of these principles constitute the hindrances to its use. Despite the high CP content, its incorporation in animal diets was discouraged due to their adverse effect on production traits (characteristics of animals such as quantity or quality of milk, meat, draught, etc.) because of the presence of bitter and toxic triterpenoids mainly nimbin, nimbidin, azadirachtin and salanin(24).

Biologically active principles isolated from different parts of the plant include: azadirachtin, meliacin, gedunin, salanin, nimbin, valassin and many other derivatives of these principles. Meliacine forms the bitter principles of the neem oil; the seed also contain tignic acid (5-methyl-2-butanic acid) responsible for the distinctive odour of the oil (25, 26,27, 28). Neem seed cake is very unpalatable due to the presence of salanolide meliacin which has been found to be one of the active principles of neem seed oil (29). These compounds are natural products called triterpenoids or more specifically, linonoids and they are concentrated in the seed despite their presence in other parts of neem tree.

Poor palatability was accompanied by either poor weight gain or loss of body weight along with lowered nutrient digestibility in crossbred calves fed concentrate mixture containing NSC that contributed 12.5, 25, or 50% of CP requirement (30). The author suggested that NSC was unsuitable for animal feeding even for maintenance. There was gradual decline in growth rate of rams with higher levels of NSC in concentrate ration (31). This then makes the processing of the

Comment [A7]: I can't understand the connection. Please rewrite from Despite until salanin.

Comment [A8]: Are they actives against what?

Comment [A9]: again you didn't explained what means CP as recommended: crude protein (CP).

cake before being fed to animals very important. The processing of the neem seed into palatable seed cake poses a challenge to its use as animal feed.

2.3 Methods of Processing and Removal of the Active Principles of Neem Seed Cake

Single seeded mature neem fruit contains 23.8% skin, 47.5% pulp, 18.6% shell and 10.1% kernel. The decortications of depulped seed yields about 26% kernel, which gives 45 to 50% oil leaving the rest as neem kernel cake (NKC) (32). Before oil extraction, neem fruits were soaked in water for 3-4 days and depulped using depulper machine. The seeds were dried for 7 days before being decorticated using a winnowing machine and then crushed after further drying for 3 days (33). The ripe fruits were dried by spreading them in the sun for fifteen days. The dried fruits were soaked in water for three days and then depulped. The depulped seeds were washed and sun dried for a period of ten days. The dry seeds were decorticated, further dried for five days, crushed and the oil removed manually to produce the neem kernel cake (22). The collected seeds were sun dried to constant weight. The dried seeds were soaked in water in an open basin for 72 h. The seed were poured into a jute bag to drain the water and later sun-dried to constant weight. Then, the water soaked and untreated seeds were taken to the mill separately for oil extraction. Cake obtained from the oil-extraction was then ground in a hammer mill-(34).

We can observe different spaces through the manuscript (1.5, 2.0...why?).

The bitter taste and pungent odour resulting in poor palatability discourage animals from eating it. NSC has been found to be unpalatable in calves (35), cross-breed bulls(36) and sheep (31) and possess antiparasitic properties (37) hence the need to remove them. The active principles are slightly hydrophilic but freely lipophilic and highly soluble in organic solvents like hydrocarbon, alcohols, ketones and esters (26,25,38). There are various methods of removing the oil from the neem seed cake. For the expeller neem seed cake, the crushed kernel will be steamed and the oil is pressed out using expeller machine. The hydraulic press neem seed cake processing is devoid of heat. The milled kernel was cold-pressed using the hydraulic press machine until the oil content of the residue (cake) is minimal. To reduce the oil content of the cake, it is further defatted using hexane. The removal of these bioactive can be done using simple techniques as follows (38)

Water extraction

Comment [A10]: From here you describe a process without an introduction. Is it a preparation for oil extraction?

Comment [A11]: Is this another description from another author? The information in this topic 2.3 is lost without an explanation or discussion...

Comment [A12]: Is this the only information about the biological activity of neem? No discussion about it again in this version of the manuscript.

The simplest technique is to crush or grind the kernels and extract them with water. They may be steeped overnight in a cloth bag suspended in a barrel of water or water can be poured into the bag and extract can be collected as it emerges. By using water extraction, it has been estimated that 20-30 kg neem seeds can normally treat one hectare of land. Normally, the proportions employed are about 500 g of kernels steeped in as much as 10 litres of water as they have low solubility in water.

Hexane extraction

Kernels are grated and steeped in the solvent hexane to extract only oil. The residue left after the hexane extraction still contains limonoid active ingredients and also subsequent water or alcohol extraction yields clean limonoids uncontaminated by oil.

Alcohol extraction

It is the most direct process for producing neem based pesticidal materials in concentrated form. Limonoids are highly soluble in alcohol solvents. The grated kernels are usually soaked in ethanol, but sometimes in methanol. It extracts the active ingredients which ranges between 0.2-6.2 percent.

Other attempts have been made at removing the bioactive principles of the cake for improved palatability. The solvent extraction method have been adopted in which the oil from the crushed neem kernel or cake is extracted using organic solvent of high polarity mixed with water and filtered (39, 40). Others include alcohol treatment (41) and alkali treatment (0.8% NaOH, wt/wt) with boiling (1:2.5wt/vol) of cake followed by water washing and draining off after washing (42). Of the several methods used to detoxify NKC, for example water washing (43), hexane extraction (40) and alkali treatment of neem kernel cake (44), water washing was found to be most effective despite the loss of 22% dry matter (45). However, to avoid this dry matter loss from water washing, alkali treated (20g NaOH/kg cake, wt/wt) NKC without washing was tried and found to be palatable to adult cattle and buffalo (46), proved promising in the feeding of buffalo calves(47) and; Uda lambs and rams (22).

2.4 Nutrient Digestibility and Performance

Feeding neem seed cake/kernel without treatment has been shown to have negative effect on nutrient digestibility and performance of ruminants. 27% level of inclusion of neem seed

190 cake was fed as a replacement to groundnut cake with consequent poor growth in cattle (48).
 191 There was poor palatability accompanied by either depressed weight gain or loss of body weight
 192 along with lowered nutrient digestibility in crossbred calves fed concentrate mixture containing NSC
 193 that contributed 12.5, 25.0 and 50.0% of digestible crude protein requirement, suggesting that the
 194 NSC as such was unsuitable for animal feeding even for maintenance (35). Lowered rumen
 195 protozoan count along with reduced enzyme activity was reported in calves fed NSC (49).

196 We can observe different spaces through the manuscript (1,5, 2,0...why?),

197 Over night soaking of NKC in NaOH (0.8%, wt/wt) and water followed by repeated draining off with
 198 two or three times water washing was carried out (45). Such water washed neem kernel cake
 199 (WWNKC) when incorporated after sun drying at 45% level in the concentrate mixture for bull
 200 calves for 273 days resulted in significantly lower growth rate and comparable DM intake, nutrient
 201 digestibility, balance of Ca, P, N and TDN intake with the control. There was significantly higher
 202 N-balance and faster growth of buffalo calves fed 40% WWNKC than in those on control diet (45).
 203 It was observed that there was no significant variations in milk yield, milk fat content, and sensory
 204 evaluation of milk, dry matter intake and digestibility when WWNKC was incorporated at 40% level
 205 in the concentrate diet of dairy cow for 300 days (50).

206 There were reports of comparable crude protein digestibility and efficiency; and live weight
 207 changes in Uda lambs and rams when fed alkali treated neem kernel cake soaked overnight in
 208 water and dried in the sun as shown in tables 1 and 2 respectively. Crude protein digestibility and
 209 efficiency in the treatment diets in these two tables compared favourably with control diets.

210 **Table 1: Live Changes and Crude Protein Digestibility of Uda Rams fed Alkali Treated Neem**
 211 **Kernel Cake (ATNKC)**

Parameter	Treatments					±SE
	A	B	C	D	E	
ATNKC Inclusion	4	5	10	15	20	
Initial body Weight (kg)	18.50	18.45	18.50	18.50	18.50	1.83
Final body weight (kg)	24.63	27.25	26.25	26.75	24.17	2.20
Body weight gain (kg)	6.13 ^b	8.80 ^a	7.75 ^{ab}	8.25 ^{ab}	5.67 ^b	0.83
Av. daily gain (g/day)	72.76 ^b	104.76 ^a	92.26 ^{ab}	98.25 ^{ab}	67.50 ^b	9.88
Feed intake (g/day)	692.93 ^{ab}	856.24 ^a	572.86 ^b	645.23 ^{ab}	691.02 ^{ab}	66.48

Comment [A13]: Rewrite

Comment [A14]:

Comment [A15]: How much? As it was said before: When we really want to show that a treatment is better or not, it's important to quantify.

Formatted: Normal, Left, Line spacing: single

Formatted: Font: (Default) Calibri, 11 pt

Comment [A16]: How much?

Comment [A17]: How much?

Dry matter intake (g/day)	658.78 ^{ab}	813.43 ^a	538.49 ^b	609.45 ^{ab}	654.34 ^{ab}	62.71
Crude Protein Intake (g/day)	107.51 ^{ab}	133.38 ^a	89.12 ^b	100.61 ^{ab}	108.04 ^{ab}	10.51
Crude Protein digestibility	77.23 ^b	92.27 ^a	86.47 ^a	86.67 ^a	78.44 ^b	2.20
Protein Efficiency	0.68 ^b	0.82 ^{ab}	1.03 ^a	1.02 ^a	0.71 ^b	0.093
Feed gain ratio	9.23 ^a	8.12 ^{ab}	5.86 ^b	6.41 ^b	9.67 ^a	0.83

Means not followed by the same superscripts are significantly different (P<0.05) along the row.

Source: 51

Table 2: Live weight Changes and Crude Protein by the Fattening Uda Rams fed Alkali Neem Kernel Cake (ATNKC)

Parameter	Treatments				± SE
	A	B	C	D	
ATNKC Inclusion	-	8.5	16.5	25	
Initial Weight(kg)	26.38	26.25	26.25	26.38	1.51
Final Weight(kg)	38.25	40.25	37.38	36.38	1.20
Weight Gain(kg)	11.88 ^{ab}	14.0 ^a	11.13 ^b	10.0 ^b	0.80
Average Daily gain(g)	141.37 ^{ab}	166.67 ^a	132.44 ^b	119.05 ^b	9.55
Feed Intake(g)	1320.25 ^a	1132.75 ^b	1057.50 ^b	1063.75 ^b	46.30
Dry matter intake(g)	1271.25 ^a	1117.25 ^b	998.25 ^b	1026.50 ^b	41.74
Crude Protein intake(g)	165.51a	145.93b	130.08b	136.28b	5.97
Crude Protein digestibility	83.08 ^a	79.59 ^a	76.11 ^a	66.80 ^b	2.74
Protein efficiency	0.86 ^u	1.14 ^a	1.02 ^{ab}	0.87 ^b	0.06
Feed gain ratio	9.04 ^a	6.71 ^b	7.77 ^{ab}	8.77 ^a	0.56

Means not followed by the same superscripts are significantly different (P<0.05) along the row.

Source:22

2.5 Nutritional Quality of Neem Seed Cake

NSC, a by-product of neem oil industry, is a non-conventional feed ingredient with great potential for livestock feeding (39,52). It has been noted as a rich protein source with 34% - 38% CP (33) and; 33.20% and 32.90% for alkali treated neem kernel cake (ATNKC) and neem kernel cake (NKC) respectively (22). Minerals contained in NSC is shown in Table 3. It is balanced in Ca, P but exceptionally high kernel Fe.

Formatted: Highlight

Table 3: Mineral profile of certain neem products (DM)

Product	%					ppm						
	Ca Pb	P	Mg	Na	K	Cu	Zn	Fe	Co	Mn	Cr	
Neem leaves	0.72	0.27	0.77	0.56	1.85	30	15	777	8	59	.5	30
Neem fruits	0.29	0.36	0.54	0.36	0.74	10	16	775	3	12	0.2	7.8
Neem seed	0.77	0.31	0.37	0.47	1.94	18	26	970	1.3	34.5	-	14.3
Neem cake	0.96	0.30	0.44	0.40	0.98	19	19	2705	1.5	70	1.0	10.6
FFNSM	0.07	0.01	0.10	0.59	1.52	10	60	300	-	20	-	-

Source: 53

FFNSM: Full fat neem seed meal

The undecorticated NSC contained 6.5 to 11.6% digestible crude protein_(36). The following are the crude protein percentages of 23.19%, 22.69%, 23.06% and 22.5% for raw neem seed meal, hydraulic press neem seed cake, solvent extracted Neem seed cake and expeller neem seed cake respectively (33). The protein quality of NKNeem kernel cake is comparable to that of peanut meal (PNM) (54). Neem cake consists of all essential and non-essential amino acids including sulphur containing ones, but with fewer amounts of histidine, lysine and tyrosine (53).

Chemical composition of neem seed cake/kernel cake and neem seed meal is shown in Table 4 and varied considerably depending on the method of processing. CP varied from 12.35% in NSC to 40.91% in urea treated neem kernel cake (UANKC). The crude fibre was highest in full fat neem seed meal of 40.50% (55) while the lowest value was reported in neem kernel cake with 11.40% (45). The ether extract (EE) is 0.38% in deoiled neem seed cake (56) and 27% in full fat neem seed meal (55). The nitrogen free extract was the lowest in full fat neem seed meal (18) with 14%

240 and highest in **neem seed cake** NSC (35) with 52.52%. UANKC seemed to be the best because of
 241 high crude protein of 40.91% and relatively low crude fibre of 11.43% (57).

242 **ATNKC Alkali treated neem kernel cake** consisted of 33.76% crude protein (46). The authors also
 243 reported that **alkali treated neem kernel cake** can be a wholesome substitute for **PNMpea nut meal**
 244 in terms of performance.

245

246 **Table 4: Chemical composition of various types of neem cake/meal (%DM)**

Neem cake/meal	CP	EE	CF	NFE	Ash	Ca	P	Reference
NSC	12.35	3.3	17.9	52.52	13.93	-	-	(35)
NSC	17.85	3.62	25.90	46.20	5.5	0.72	0.58	(23)
Neem fruit lake	14.97	2.40	26.95	40.92	15.17	1.65	0.31	(58)
NKC	33.50	10.40	11.40	29.70	15.00	-	-	(45)
NSM	35.51	9.14	11.89	28.48	14.98	0.77	0.69	(59)
Hexane Exp. NSM	38.31	1.06	13.46	29.02	18.15	1.05	1.08	(59)
Alcohol NSM	Exp.	40.35	0.71	13.92	26.02	19.00	1.12	1.11 (59)
NSC	20.90	14.80	19.20	29.50	15.50	0.14	0.81	(54)
FFNSM	16.80	22.80	34.61	23.10	2.7	-	-	(18)
FFNSM	14.20	27.00	40.50	14.00	14.30	0.07	0.01	(55)
DNSC	18.37	0.381	30.12	34.97	16.16	0.97	0.16	(56)
ATNKC	33.76	5.39	13.77	27.28	19.90	-	-	(46)
UANKC	40.91	4.44	11.43	28.10	15.12	-	-	(57)
ATNKM	36.01	4.80	15.60	27.80	15.80	0.78	0.50	(53)
UANKM	40.06	4.90	15.30	24.12	15.62	0.74	0.45	(53)

247 NSC: Neem seed cake; NKC/NKM: Neem Kernel Cake/Meal; WWNKC: Water wash NKC; NSM: Neem
 248 seed meal; FFNSM: Full fat NSM; DNSC: Deoiled NSC; ATNKC/ATNKM: Alkali treated NKC/NKM;
 249 UANKC/UANKM: Urea-Ammoniated NKC/NKM; Exp: Expeller.

250

251 2.6 Palatability of Neem Seed Cake/Kernel Cake

Comment [A18]: As recommended before you didn't review all the manuscript. You have to decide if you will use the full name or the abbreviations!! Is the same situation:" you inform that neem seed meal is NSM and in line 216 you repeat neem seed meal. The same was highlighted for NSC in different parts of the text. You have to decide if you will use the full name or the abbreviation and make it homogeneous in the text. Other examples: Line 217: neem seed cake, Line 218: neem seed meal, and others."

Formatted: Highlight

The palatability of the NSC can be improved with appropriate treatment to remove the active principles. NSC palatability improved when NSC was fed to sheep along with barley, molasses and ~~PNM~~peanut meal but consumption of concentrate mixture was reduced from 79 to 39% when NSC levels increased from 59% to 90% respectively (60). Yearling sheep could completely consume concentrate mixture comprising 75 parts NSC and 25-parts maize but the consumption was reduced to one third when NSC was fed alone (31). Buffalo calves though continued to relish even after increasing NSC from 5 to 15 parts when fed along with 7 and 20 parts of molasses and peanut meal respectively, the consumption was reduced to half on withdrawal of molasses (61).

2.7 Effect on Haematological and Biochemical Characteristics

The use of treated ~~NSC~~neem seed cake/neem kernel cake in the feed of ruminants has been proven to be safe for their consumption. Feeding of ~~ATNKC~~ alkali treated neem kernel cake to lambs for up to 20% level has no significant effect on the haematological and biochemical characteristics (51). Neem bitterness and toxicity can be inactivated by alkali treatment (60). This was also supported by (53) in his report that ~~ATNKC~~ alkali treated neem kernel cake can convert ~~NSC~~neem kernel cake to a wholesome vegetable protein supplement for growing buffalo calves. Inclusion of water washed ~~NSC~~neem kernel cake in the concentrate mixture of male kids at 15 and 25 parts for 180 days of feeding led to comparable intake and utilization of nutrients, besides normal balance of nitrogen, urinary creatine, blood haemoglobin, cholesterol and activities of ~~g~~Glutamate oxalo acetate transaminase (GOT), ~~g~~Glutamate pyruvate transaminase (GPT) and alkaline phosphates with significantly lowered blood glucose, urea and total protein (61).

The ~~s~~Serum glutamate oxalo acetate transaminase (SGOT) and ~~s~~Serum glutamate pyruvate transaminase (SGPT) values in the control and test diets Uda lambs fed ~~ATNKC~~ alkali neem treated kernel cake were comparable (51). He also showed that the ~~b~~Bilirubin total, ~~b~~Bilirubin ~~c~~Conjugate and ~~b~~Bilirubin unconjugate values in the study were normal and conformed to the report of (62) and (63). These values were within the range of 14-123u/l for SGOT and 15-44u/l for SGPT reported by (64). These indicate that inclusion of ATNKC is not toxic to the liver since SGPT and SGOT were excellent markers of liver damage caused by exposure to toxic substances (64). Incorporation of NSC up to 20% did not alter GOT and GPT activities in the blood (65). The urea nitrogen level of lambs fed ~~ATNKC~~ alkali treated neem kernel cake and control were not significantly different from each other (51) and were all within the normal range reported for sheep by (63) and (62). The authors also showed that the ~~S~~sodium and ~~p~~Potassium level obtained the

blood of Uda lambs fed ~~ATNKC alkaline treated neem kernel cake~~ were normal and within the range of 142-160mmol/l for Sodium and 4.3-6.3mmol/l for potassium reported by (63) This indicate that ATNKC inclusion up to 20% level did not interfere with the renal functions of the animals.

2.8 Economics of use

~~NSCNeem seed cake~~ has proven to be economical in the feed of ruminants. This is shown in tables 5 and 6 respectively in which treated ~~NSCneem seed cake~~ inclusion in the feed of Uda lambs and rams compared ~~favourably~~favourably in cost of feed per kilogram, cost of feed consumed per day and cost of feed per kilogram live weight gain. It was reported that 20% lower cost per weight gain was reported when deoiled peanut was replaced with ~~with~~-urea ammoniated ~~NKMneem kernel meal~~ in goats (61). The cost of feed per kg liveweight gain was lower for diets with inclusion of detoxified ~~NSCneem seed cake~~ than with the conventional feed in Nellore sheep (66).

Table 5: Cost of feed/kg liveweight of Uda lambs

Parameter	Treatments					±SE
	A	B	C	D	E	
Cost of Feed (\$/kg)	0.28	0.276	0.273	0.276	41.72	-
Cost of feed consumed (\$/day)	0.193 ^{ab}	0.236 ^a	0.157 ^b	0.178 ^b	0.182 ^{ab}	2.72
Cost of feed/kg live weight gain (\$/kg)	2.650 ^a	2.254 ^{ab}	1.700 ^b	1.811 ^b	2.702 ^a	36.24

Means not followed by the same superscripts are significantly different (P<0.05) along the row.
Source:22

Table 6: Cost of feed/kg live weight of Uda ram

Parameter	Treatments				±SE
	A	B	C	D	
Cost of feed (\$/kg)	0.247	0.227	0.218	31.06	
Cost of feed consumed (\$/day)	0.315	0.254	0.218	0.201	1.38

Comment [A19]: again: What do treatments A to E mean?

Comment [A20]: again: What do treatments A to E mean?

Cost feed per live-weight gain (\$/kg)	2.225	1.524	1.643	1.694	19.51
--	-------	-------	-------	-------	-------

Means not followed by the same superscripts are significantly different ($P < 0.05$) along the row.
Source:22

3.0 CONCLUSION

Neem seed is readily available in Nigeria especially in the northern part. It can easily be converted to neem seed cake/kernel cake. The active principles present in the seed have resulted in bitterness and pungent smell which have impacted negatively on the palatability and the consumption. However, the various methods of preparing the cake and removing the active principles have resulted in improved palatability and consumption as shown in the review. It is shown that ruminants utilise feed incorporated with treated neem seed cake/ kernel cake comparably with the control diets without any adverse effect on the haematological and biochemical parameters. Treated neem seed cake/kernel cake can then readily fit in as replacement for the convectional protein sources.

Comment [A21]: rewrite

REFERENCE

1. Getup JO. Socio-economic Considerations in Small Ruminant Production. In: Lakpini,C.A.M.,Adamu,A.M., Edoche,O.W. and Gefu, J.O.(eds). Manual for Small Ruminant Production in Nigeria. National Animal Production Research Institute,Ahmadu Bello University, Zaria, Nigeria. 2002; Pp.8-11.
2. David-West KB. Role of Government in small Ruminant production. In: *Small Ruminant Production in Nigeria*. (Adu, I.F., Osinowo, O.A., Taiwo, B.B.A, and Alhassan, W.S. (eds.)). Proceedings of National Conference on Small Ruminant Production, NAPRI, Shika, A.B.U. Zaria, Nigeria. 6-10 October; 1985.
3. NAERLS. Prospects and problems of 1999 cropping season. Report submitted to Federal Ministry of Agricultural Extension and Research Liaison services, Ahmadu Bello University, Zaria, October 1999; Pp 105.
4. Shuaib BA, Aliyu A, Bakshi JS. Nigeria National Agricultural Research Strategy Plans, 1996-2010. Department of Agricultural Science, Federal Ministry of Agriculture and Natural Resources, Abuja; 1997.
5. Mubi A. The effect of supplementing legume crop residue with *Faidherbia albida* pods for Goat feeding in the North-East Nigeria. *J.Agric. Res. and Dev.* 2003; 2:25-31.
6. Adegbola TA. Forage resources and beef production in Nigeria. In: *Beef Production in Nigeria*. (Osinowo, O.A., Ikhatua, U.S. and Ehoche, W.O. (eds)). Proceedings of the National Conference on Beef Production, July 27-30, NAPRI, ABU, Zaria, Nigeria; 1982.
7. Oyenuga VA. Nigeria's food and feeding stuffs: Their Chemistry and Nutritive value. Ibadan University press, Ibadan, Nigeria; 1968.

8. Steinbach J. Alternative to crop residues as feed resources in mixed farming system. In: *Crop Residues in Sustainable Mixed Crop/Livestock farming* (Renard C. Eds), AB international, Netherland; 1997.
9. Onwuka CFI, Isah OA, Oni AO, Aderinboye RY. Ruminant animal nutrition. Assesed on 20th Jan. 2013. www.unaab.edu.ng/attachment/459_LectureNote-ANN503_1.PDF (Unpublished)
10. Davis OA, Ezenwa NC. Groundnut cake as alternative protein source in the diet of *Clarias Gariepinus* Fry. *Int. J. Sc. & Nat.* 2010; 1(1). 73-76.
11. FAO. Groundnut (*Arachis hypogaea*). *Encyclopedia of Agricultural Sciences*, 2000; Vol. 3.
12. Milo D. Nutritional values of cotton seed cake. *Livestrong.com*. 2010. Assesed on 22nd Feb. 2013. www.livestrong.com.
13. Ramesh CS. Neem for sustainable development and environmental conservation. *Indian Perspective*, 2000; Pp 39-51.
14. Panhwar MH, Panhwar F. Neem-A natural Pesticide; 2003. Accessed 20th July, 2010, Available: <http://www.panhwar.com/Article 40.htm>
15. Sokumbi OA, Egbunike GN. Physiological responses of growing rabbits to neem (*Azadirachta indica*) leaf meal based diets: Haematology and serum biochemistry. *Trop. Anim. Prod. Invest.*, 2000; 3:81-87.
16. Anon. The Neem Tree. 2010. Accessed 30 August, 2012, Available: http://www.theneem.co.uk/acatalog/The_Neem_Tree.html,
17. NARICT. National Research Institute for Chemical Technology. *Economic importance of neem tree*. Seminar Exposition: Neem for Economic Empowerment. Conference Hill, Basawa, Zaria; 2004.
18. Fajinmi AO, Adedeji SK, Hassan WA, Babatunde GM. Inclusion of non-conventional feedstuffs in rabbit concentrate ration, a case of neem seeds. *Appl. Rabbit Res.* 1989; 13:125-128.
19. Nwokeabia OD, Annual report. Federal Department of Forestry, Federal Ministry of Agriculture, Abuja, Nigeria, 1994; pp.46.
20. The Neem Team. 2003. Assesed on 29 March, 2010. Available: http://www.theneem.co.uk/acatalog/The_Neem_Tree.html; 2003.
21. Radwanski S. Neem tree 2. Uses and potential uses. *World Crops and Livestock*, July/August, 1977; Pp. 167-168.
22. Aruwayo A. Effect of Evaluation of alkali treated neem kernel cake fed to Uda Sheep in a Semi-Arid zone of Nigeria, Ph.D Thesis, 2011, pg 131. (Unpublished).
23. Christopher J. Neem seed cake is also good for cattle: *Indian Farming*. 1970; 4:38-40.
24. Paul BN, Haque N, Garg, A. K. Effect of feeding neem kernel cake (*Azadirachta indica*) on rumen enzyme of buffalo. *India Vet. J.* 1996; 73: 720-728.
25. Schmutterer H. *The neem Tree Source of Unique National Product for IPM, Medicine, Industry and Other Purposes*. VCH Publication, New York; 1995.
26. NRC. *Neem: A Tree for solving Global Problems*, National Academy Press, Washington D.C; 1992.

- 379 27. Uko OJ, Kamalu TN. The neem tree: uses and potentials. *Nig. J. Exptl. Appl. Biol.* 2001;
380 2(2): 223-229.
- 381 28. Lale NES. Bio-activity and Limitation against wide spread use of neem products for the
382 management of insect pests. *Nig. J. Appl. Biol.*, 2002; 115-124.
- 383 29. Garg HS, Bhakuni DS. *Phytochem* 23:2383. In: Handbook of African Medicinal Plants
384 (Iwu, M. M., ed.), CRC press Inc., London,1993; PP. 124-128.
- 385 30. Arora SP, Singhal SP, Ludri RS. Nutritive value of neem seed cake (*Melia indica*). *Ind.*
386 *Vet. J.* 1975; 52:867-870.
- 387 31. Gupta RS, Bhaid MV. A study on the effect of feeding different levels of deoiled neem
388 fruitcake in the concentrate mixture on water consumption in lambs. A review of Agro-
389 Animal Science and Health. p. 121; 1980.
- 390 32. Ketkar CM. Utilization of neem and its by-products. Final Technical Report, Khach and
391 Village Industries Commission, Bombay, India; 1976.
- 392 33. Bawa GS, Orunmuyi M, Agbaji AS, Landam Z, Okekeifi UO. Effect of different methods of
393 processing neem (*Azadirachta indica*) seeds on performance of young Rabbits. *Pak. J.*
394 *Nutr.* 2007; 6 (3): 213-216.
- 395 34. Odunsi AA, Adegbile AA, Akande TO, Olayemi TB. Neem (*Azadirachta indica*) Seeds
396 Cake in Diets of Cockrel Chickens. *Int. J. Poult. Sc.* 2009; 8(1): 47-49.
- 397 35. Bedi SPS, Vijjan VK, Ranjhan SK. Utilization of neem (*Azadirachta indica*) seed cake
398 and its influence on nutrients digestibility in buffaloes. *Ind. J. Sci.* 1975b; 28:104-107.
- 399 36. Ananthasubramainiam CR, Menacherry M, Devasia PA. Studies on the feeding value of
400 neem (*Azadirachta indica*) seed cake for cattle. *Kerala J. Vet. Sci.* 1979;10:182-185.
- 401 37 Ogbuewu IP, Odoemenam VU, Obikaonu HO, Opara MN, Emanaham OO, Uceagbu MC,
402 Okoli IC, Esonu BO, Illoeje MU. The growing impotence of neem (*Azadirachta indica*) in
403 Agriculture industry. *Res. J. Med. Report.* 2011; 5(3):230-235.
- 404
- 405 38 Chandy, KT. Neem Utilization. Booklet No. 470 Medicinal Plants: MPS-15 **Assessed on**
406 **20th Feb. 2013. Available:** [www.inseda.org/.../CD%20.../53.../Neem%20Utilisation-](http://www.inseda.org/.../CD%20.../53.../Neem%20Utilisation-470.doc)
407 [470.doc](http://www.inseda.org/.../CD%20.../53.../Neem%20Utilisation-470.doc).
- 408 39 Nath K, Bedi SPS, Vijjan UK, Ranjham SK. Nutritional studies with neem seed cake. In:
409 *Anim. Rep. Anim. Nutr. Div. IVRI, Izatnagar, India; 1974.*
- 410 40 Sadagopan VR, Johri TS, Reddy VR. Feeding value of neem seed meal in broiler and layer
411 diet. *Indian Poultry Gazzette*, 1981; 65:136-142.
- 412 41 Gupta GP, June S, Mukherjee PC, Mitra CR, Mistra PS. Processed neem (*Melia indica*)
413 meal in poultry. In: Ketkat, C. F. (1976). Utilization of neem and its by-products. Final
414 Technical Report, Khach and Village Industries Commission, Bombay, India; 1976.
- 415 42 Agrawal DK, Garg AK, Nath K. The use of water washed neem (*Azadirachta indica*) kernel
416 cake in the feeding of buffalo calves. *J. Agric. Sci. (Camb).* 1987; 108:497-499.
- 417 43 Choudhary KK, Saikan A, Baruah KK, Pathak NN. Nutritional significance of neem seed
418 cake. *Poult. Guide.* 1981; 18:43-46.
- 419 44 Reddy VR, Rao PV, Reddy CV. Utilization of chemically treated neem oil in broiler chicks.
420 *Indian J. Anim. Sci.* 1988a; 58:830-834.

Formatted: Portuguese (Brazil)

- 421 45 Nath K, Rajgopal S, Garg AK. Water washed neem (*Azadirachta indica*) seed kernel cake
422 as a cattle feed. *J. Agric. Sci. (Camb)*, 1983; 101:323-326.
- 423 46 Katiyar RC, Sastry VRB, Agrawal DK. Nutrient utilization from alkali treated neem seed
424 kernel cake by cattle and buffalo. *Indian J. Anim. Nutr.* 1993; 10(4): 223-226.
- 425 47 Reddy KCV. Performance of buffalo (*Bubalus bubalis*) calves fed alkali treated and urea
426 ammoniated neem (*Azadirachta indica*) seed kernel cake incorporated rations. M.V.Sc.
427 thesis, Deemed University, India; 1992, (Unpublished).
- 428 48 Pradhan K. Utilization of neem and its by-products. In: Final Report, khadi and village
429 industries commission (ketkat, C.M., ed.), Bombay, India; 1975.
- 430 49 Mondal DP, Garg AK, Mondal BC, Pathak NN. Effect of feeding neem kernel meal on
431 ruminal protozoan assay. Abst. No. 137, 7th Animal Nutrition Research Workers Conf.
432 Bombay, India; 1995.
- 433 50 Nath K, Agrawal, DK, Hassan QZ, Daniel SJ, Sastry VRB. Water washed neem
434 (*Azadirachta indica*) seed kernel cake in feeding of milk cows. *Anim. Prod.* 1989; 48:497-
435 502.
- 436 51 Aruwayo A, Maigandi SA, Malami BS, Daneji AI. Growth performance of growing Uda
437 lambs fed graded levels of alkali treated neem kernel cake. *J. Sci.Tech. Res.* 2010; 9(4):
438 119-122.
- 439 52 Bawa GS, Orunmuyi M, Onabanjo OA. Effect of dietary inclusion levels of mechanically
440 extracted neem seed cake on performance of young rabbits. *Nig. J. Anim. Prod.* 2005;
441 32:233-239.
- 442 53 Gowda SK, Sastry VRB. Neem (*Azadirachta indica*) seed cake in animal. Feeding-scope
443 and limitations review. *J. Anim. Sci.* 2000; 13(50): 720-728.
- 444 54 Reddy VR, Rao PV. Utilization of differently processed undecorticated neem cake in broiler
445 chicks. *Indian J. Anim. Sci.* 1988b;54: 840-42.
- 446 55 Salawu MB, Adedeji SK, Hassan WH. Performance of broilers and rabbits given diets
447 containing full fat neem (*Azadirachta indica*) seed meal. *Anim. Prod.* 1994;58:285-289.
- 448 56 Garg AK. Studies on deoiled neem (*Azadirachta indica*) seed cake as a cattle feed. PhD
449 thesis, IVRI, Izatnagar, India. 1989; Pp. 129-175. (Unpublished)
- 450 57 Katiyar RC, Sastry VRB, Agrawal DK. Urea and alkali treated neem seed kernel cake as
451 livestock feed. In: *Anim. Rep. Anim. Nutri. Div.*, IVRI, Izatnagar, India; 1991.
- 452
- 453 58 Rao PV, Nath KC. Chemical of neem fruit cake. In: Gowda, S. K. and Sastry, V. R. B.
454 (2000). Neem (*Azadirachta indica*) seed cake in animal feeding-scope and limitation. *J.*
455 *Anim. Sci.* 1979; 13(5): 720-728.
- 456 59 Chand S. Nutritional evaluation of neem seed meal in chicks. Ph.D, Thesis, Rohilkhand
457 University, Bareilly, U. P., India; 1987. (Unpublished)
- 458 60 Bhandari AS, Joshi MS. The effect of feeding deoiled neem cake on health of sheep. *Ind*
459 *Vet. J.* 51:659-660; 1974.

Formatted: Portuguese (Brazil)

- 460 61 Baxi MR. Deoiled neem cake trial. In: Kektar, C.M. (1976). *Utilization of neem and it's by*
461 *product*. Final Rechnical Report, Khadi and Village Industries Commission, Bombay, India;
462 1976.
- 463 62 Coles EH. *Veterinary Clinical Pathology* (4th edition). W.B.Sanders Company, Harcourt
464 Brace Jovanovidi, Inc; 1986.
465
- 466 63 Merck Veterinary Manual of Biochemical Reference Ranges.
467 Avalable:[http://www.merckvetmanual.com/mvm/servlet/CVMHighLight?File=htm/bc/treft.ht](http://www.merckvetmanual.com/mvm/servlet/CVMHighLight?File=htm/bc/treft.htm&w;)
468 [m&w](http://www.merckvetmanual.com/mvm/servlet/CVMHighLight?File=htm/bc/treft.htm&w;); 2008.
- 469 64 Ranjna C. Practical Clinical Biochemistry Methods and Interpretation. 2nd ed., pp 117;
470 1999.
- 471 65 Gangopadhyay P, Pyne AK, Moitra DN, Mazumdar S. Studies on the biochemical
472 constituents of blood with incorporation of neem seed cake in the ration of Milch Murrah
473 buffaloes. *Indian J. Anim Health*. 1981;20: 61-63.
- 474 66 Madhavi K, Reedy TJ, Reddy YR, Reddy GVN. Effect of feeding differently processed
475 detoxified neem (*Azadirachta Indica*) seed cake based complete diet on growth, nutrient
476 utilization and carcass characteristics in Nellore sheep. *Livestock for Rural Development*.
477 2006;18(10).
- 478