

INFLUENCE OF AGRICULTURAL LIME ON SOIL PROPERTIES AND WHEAT (*Triticum aestivum* L.) YIELD ON ACIDIC SOILS OF UASIN GISHU COUNTY, KENYA.

**Osundwa, M.A.^{1*}, Okalebo, J.R.², Ngetich, W.K.², Ochuodho, J.O.³,
Othieno, C.O.², B. Langat⁴, Omenya, V.S.²**

* Ministry of Agriculture, P.O Box 95, Eldoret, Kenya. Email-osundwamargaret@yahoo.com

² University of Eldoret, Department of Soil Science, P. O. Box 1125, Eldoret, Kenya

³ University of Eldoret, Department of Seed Crop and Horticultural Sciences, P. O. Box 1125, Eldoret, Kenya

⁴ Moi University, Department of Agribusiness, P. O. Box 3900, Eldoret, Kenya

ABSTRACT

A study was carried out to investigate the influence of agricultural lime (21% CaO) from Koru, Kisumu on soil properties and wheat yield on acidic soils of Uasin Gishu County. Field trials were conducted at Chepkoilel University College farm and in Kipsangui area of Uasin Gishu county. Soils were analyzed to determine their pH, available P and other nutrient levels before treatment application. The experiment was a split plot arrangement with two wheat varieties as the main plots and the lime treatments as the subplots. The two varieties compared were 'Njoro BW 2' and 'KS Mwamba' characterized as tolerant and moderate tolerant to soil acidity, respectively. Phosphorus and nitrogen were applied as a blanket treatment at the rates of 40 kg P₂O₅ ha⁻¹ and 46 kg N ha⁻¹ respectively. Lime was applied at the rates of 0.0, 0.5, 1.0, 1.5 and 2.0 t ha⁻¹. Soils from the two sites were acidic with low to moderate available P for Chepkoilel and Kipsangui sites respectively. Soil pH and soil available P increased with the increase in the rate of lime addition. Wheat grain yield increased significantly (p<0.05) due to soil acidity amendment above the control. There was a high positive correlation between wheat yields and soil available P at both sites at harvest. High cost of inorganic inputs, low wheat grain prices and the effects of the erratic rains made the majority of the treatments economically unviable for adaptation by farmers. However, the most profitable treatment was 40 kg P₂O₅ ha⁻¹ with 46 kg N ha⁻¹ and 2 t ha⁻¹ in Njoro BW 2 at Kipsangui site. There was no viable treatment at Chepkoilel site. Higher wheat yields may probably be achieved from rates of lime above 2 t ha⁻¹.

Keywords: Lime, nitrogen, phosphorus, soil acidity, wheat

* Corresponding author. Tel. +2540726457993

Email-osundwamargaret@yahoo.com

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is the first most important cereal cultivated in the world. It is the second most important cereal crop grown in Kenya after maize [1]. In contrast to sustained increases in wheat productivity in other parts of the developing world, per capita production in Kenya continues to stagnate while the consumption has been on the increase [2]. Uasin Gishu county is a major wheat producing county of Kenya, producing more than one-third of the total wheat produced in the country (3). However, over the recent years there has been a general decline of crop yields due to unpredictable weather particularly rainfall, declining soil fertility, diminishing land parcels as the consequence of rapid population growth, low and unsustainable market prices for the produce and poor crop husbandry [4]. High costs of inputs mainly Di-Ammonium Phosphate and Calcium of Ammonium nitrate, diseases, weeds and poor crop husbandry, also contribute significantly to low grain yields in the county. In addition to the above constraints, the wheat crop in this county is mainly grown on ferralsols which are characterized by low pH (soil acidity) and low nutrient levels [5]. These ferrous impervious soils also result in frequent water stagnation [6]. Although ferralsols have good internal drainage; water stagnation in this county is caused by soil compaction and crusting as a result of the use of farm machinery. Nitrogen, calcium and magnesium deficiencies and toxicities of aluminum and manganese, which characterize these acidic soils, also limit crop production in this district [7]. Further, continuous use of acidifying fertilizers like Di-Ammonium Phosphate and urea, worsen an already bad situation [8].

Soil acidity is attributed mainly to abundance of hydrogen (H^+), aluminum (Al^{3+}) and manganese (Mn^{2+}) cations in soils at levels that interfere with normal plant growth. Soil acidity has a negative effect on crops mainly through P unavailability from P fixation in soils whereby the Fe and Al soil components (sesquioxides) fix sizeable quantities of P. Excess Al^{3+} ions, from soil acidity, tend to accumulate in plant roots and thereby prevent P, Mo and other ions translocation to the tops from the roots, as evidenced by the inhibition of root elongation and overall retarded crop development [9,10,11]. The detrimental effect of H^+ ions is not as distinct as that of Al^{3+} cations, but excess of H^+ ions in acid soils affects plant root membrane permeability and therefore interferes with ion transport [11].

In acid soils and P deficient tropical soils where the plant capacity to scavenge the native or use added P with efficiency is critical [12], correcting soil acidity and P fertilizer addition are important. Lime is widely known as the effective way of correcting soil acidity or inputs recommended for amelioration of acid soils [13]. Its direct effect is soil pH increase [14]. In Kenya, management of soil acidity through liming is highly recommended [11, 15]. Lime reduces Al, H, Mn, and Fe toxicities and increases P, Mg, Ca and Mo availability in acidic soils [11, 16, and 17]. Therefore, in P fixing acid soils, combined lime and P application is necessary for increased availability of the applied P for plant uptake. Although, not permanent the direct effect of lime lasts longer than any other amendment, such as organic materials. The most popular varieties of wheat grown in Uasin Gishu county are Njoro BW 2 and KS Mwamba because of their outstanding characteristics. Both wheat varieties are high yielding and tolerant to diseases such leaf and stem rusts, however the yields in the county have remained low even with proper crop husbandry. Thus the objectives of this study were to (1) To investigate the effects of agricultural lime (21% CaO), on wheat production in acid soils of Uasin Gishu county. (2) To determine the response in terms of growth, changes in pH and available P (3) To evaluate the economic returns of the two wheat varieties from amelioration of depleted soils by application of lime.

2.MATERIAL AND METHODS

2.1 Uasin Gishu County

The county lies between longitudes 34° 50' and 35° 37' East and latitudes South and 0° 55' North. It is a highland plateau. Its terrain varies greatly with altitude which ranges between 1500 and metres above sea level. Eldoret Town, the capital of Uasin Gishu county, is at an altitude of 2085 metres, marks the boundary between the highest and the lowest altitudes of the county. The county's general landscape is that of an undulating plateau with no significant mountains or valleys. The average rainfall is between 900 mm -1200 mm. Due to high altitude in the county, temperatures are relatively low. The highest is 24°C and the lowest is about 8.8°C. Humidity is moderate, averaging 56%. The average temperatures in the county are 18°C during the wet season with a maximum of 26.1°C during the dry season [18].

2.1.1 Study sites

2.1.1.1 Chepkoiel University College

The field experiment was conducted at the Crop Seed and Horticultural Science Department Field, Chepkoiel University College, Moi University in Moiben division. The soils of Chepkoiel belong to a group of soils found on plateaus and high level structural plains. The soils are of igneous origin, acidic (pH: 4.5-5.0), and low in fertility and are underlain with murram. They are classified as Rhodic Ferralsols according to the FAO/UNESCO classification and Oxisols according to the USDA classification [5, 19].

2.1.1.2 Kipsangui

The experiment was also concurrently conducted at Mrs Selina Maswai's farm in Kongasis location, Kipsangui sub-location, Soy division of Uasin Gishu county. The area receives a unimodal rainfall distribution pattern. The average amount of rains received in the area for the last ten years is 1171 mm. According to National Agricultural and Livestock Extension Programme Broad Based Survey (NALEP-BBS) report of 2011, the soils are sandy loam. They are classified as Rhodic Ferralsols according to the FAO/UNESCO classification and as Oxisols according to the USDA classification [5, 19].

2.1.1.3 Experimental design and treatment

Lime was applied three months prior to planting at the rates of 0.0, 0.5, 1.0, 1.5, and 2.0 ton ha⁻¹ (quantity) as CaO from Koru, Kisumu. Lime was broadcast evenly within the plots on a fine seedbed and then incorporated into the soil using a hoe. DAP fertilizer was applied as a blanket treatment at rate of 40 kg P₂O₅ ha⁻¹ (17.6 P kg ha⁻¹). N- was applied at the rate of 46 kg N ha⁻¹. To supplement the N in the DAP, urea foliar applied at was the rate of 28 kg ha⁻¹ before boot stage. Experimental design was a split plot with varieties namely Njoro BW 2 and KS Mwamba forming the main plot and lime levels as the sub plot. Liming materials were obtained from Koru in Kisumu County, Kenya which contains about 21% CaO. fine particles of 100 mesh-. The sub plot size was 4 m *4 m.

2.1.1.4 Soil Sampling

Prior to treatment application surface (0-15) cm depth soil samples were randomly taken using a soil auger from several points at the experimental sites and thoroughly mixed together to make a composite sample. Six weeks after planting and during harvesting composite samples were again taken from each plot at the depth of 0-15 cm to monitor changes in soil available P and pH.

Comment [SD1]: when was lime applied? lime needs time to react with soil. Also, was it incorporated into the soil?

Formatted: Strong, Font: (Default) Helvetica

Comment [SD2]: what was the fine-ness of the lime?

2.1.1.5. Field procedures:

The seeds were drilled by hand at the recommended row spacing of 25 cm by drill. The placement within the row by hand was predetermined prior to planting by running a mechanical drill planter and observing /noting the closeness of the spacing. The depth of placement was established at 2.5 cm -3 cm deep. After sowing, seeds were covered with top soil and slightly compressed to ensure close seed-soil contact. This was done to ensure a rapid and even germination [20]. The seed rate was 125 kg ha⁻¹. Pesticide, O,O-Dimethyl S-(N-methylcarbamoylmethyl phosphorodithioate (56)(Dimethotate) was used to control pests. Herbicide known as 3,4-methylenedioxyamphetamine (Puma Complete) was sprayed a month after planting i.e. 4 leaf stage at the rate of 0.75 litres ha⁻¹. Fungicide, Tebuconazole - (RS)-1-p-chlorophenyl-4,4-dimethyl-3-(1H-1,2,4-triazol-methyl)pentan-3-ol (Nativo 300 SC) at 0.75 litres ha⁻¹ was also applied at 4 weeks after planting and urea foliar was sprayed just before booting stage. Harvesting of wheat was done at physiological maturity. It was done on centre rows of each plot at final harvest by discarding outer rows per sub plot. Thus the inner rows were harvested by hand giving an effective area of 14 m². The ends of plots plants were discarded. In the harvested area, total heads, fresh weights and sub samples weights were taken and recorded. The wheat was threshed manually (by beating using a stick). The sub-samples were dried in the greenhouse and their weights recorded for estimates of dry weights. The straw within the harvested area was cut at ground level at harvest and its weight taken. Sub-samples from the straw were taken randomly from each plot and cut into small pieces and mixed thoroughly. All samples were air dried (in the absence of oven) and their fresh (initial) and dry weights recorded and used to compute yields per plot (grain and straw). These samples were then ground (0.02 mm) for plant tissue analysis to determine N and P contents. Yield was calculated using the relationship:

$$\text{Yield/plot} = \frac{(\text{Total dry weight} \times \text{yield/ha}) \times \text{effective area}}{\text{Total fresh weight}}$$

Comment [SD3]: this equation needs to be reviewed. What you calculated is yield/ha, so this needs to be the output of the equation.

2.1.1.6. Laboratory analysis

All the soil samples were air-dried and sieved through 2 mm mesh. Soil samples were analysed for soil pH (Soil H₂O; 1:2.5), texture (hydrometer method), total carbon (C) % (Weakley-Black method), total N (%) (Kjeldildah distillation method) and available P (Olsen method). Samples taken after treatment application were also analysed for pH (1:2.5, water) available P, plant N and P (Kjeldildah distillation method) and their concentration determined by spectrophotometer.

Comment [SD4]: still need dispersion method

2.1.1.7 Economic analysis

The most economically acceptable were treatments were determined by partial budgeting analysis to estimate the gross value of grain by using the adjusted yield at the market value of grain inputs during the cropping year. In partial budgeting only costs that vary from the control are used referred to as total costs that vary (TCV). The prices of lime, DAP, urea foliar fungicide and pesticide, bags for storing wheat, transport and wheat grain were determined through market survey at each of the two sites during the research period. Labour wage rates for applying lime, fertilizers and shelling of the grain were also determined through market survey to estimate the labour costs that vary. Yield data were adjusted downward by 10% since research has found out that farmers using the same technologies would obtain 10% yield lower than those obtained by researchers. The discounted rate of capital was determined at the rate of 10 and 20% per season and year, respectively and was applied to cash costs only. The discounted rate reflects the farmer's preference to receive benefits as early as possible and to postpone costs. All costs and benefits were converted to monetary values in Kenya Shilling (Ksh.) and reported on a per hectare basis.

The net accrued net financial benefits (NFBs) and TCV were then compared across the treatments dominance analysis the formula shown below.

$$NFB = (Y \times P) - TCV$$

Where $Y \times P$ = Gross Field Benefit (GFB), Y = Yield per ha and P = Field price per unit of the crop.

Treatment with less than or equal to treatment with lower TCV are dominated and were marked a "D" while ones with higher NFB than the treatments and lower TCV are undominated. The marginal rate of return (MRR) analysis was carried out on the undominated treatments.

$$MRR(\%) = \frac{\text{Change in NFB}(NFB_b - NFB_a) \times 100}{\text{Change in TCV}(TCV_b - TCV_a)}$$

where NFB_a = NFB with the immediate lower TCV, NFB_b = NFB with the next higher TCV, TCV_a = the immediate lower TCV & TCV_b = the next highest TCV
Change in NFB and TCV also referred to marginal benefits and costs, respectively.

2.1.1.7 Statistical analysis

The generated data of wheat grain yield, soil and plant material across the experimental sites were subjected to analysis of variance (ANOVA) with standard error SE using General statistics. Means were separated using least significance difference (LSD) whenever treatment difference were significant ($p=0.05$).

3. RESULTS AND DISCUSSION

3.1 Soil physical and chemical characteristics of the study sites

Table 1 shows a summary of soil parameters of the surface soils (0-15 cm) of the experimental sites determined in the laboratory before treatment application. The soils pH in the study areas was 4.92 and 5.32 for Chepkoilel and Kipsangui sites respectively. The low to moderate soil pH in both sites indicates that the acidic levels require to be amended through liming if optimum crop yield is to be achieved. According to [21], liming can neutralize soil acidity and most field crops perform best at a soil pH level between 6.0 and 6.8. The soils from Chepkoilel had organic carbon of 1.69% while those from Kipsangui had organic carbon of 2.66%. The soil available P was 9.88 mg kg⁻¹ for Chepkoilel site and 10.1 mg kg⁻¹ for Kipsangui site. For the soil particle size analysis, the soils from both sites were classified as sandy loam. The moderate soil available P in these soils suggested the need for supplementary P addition for increased crop yields [22]

Comment [SD5]: please check - it is likely 5.5 for an Oxisol like this.

Table 1. Soil physical and chemical characteristics of surface (0-15 cm) soils taken before planting (2009 LR) at two study sites in Uasin Gishu district, Kenya

CHEPKOILEL SITE

PARTICULAR	VALUE	METHODS
SAND %	60	Hydrometer method [1]
SILT %	16	
CLAY %	24	
TEXTURAL CLASS	SANDY LOAM	
PH (1:2.5 SOIL: WATER)	4.92	Glass electrode pH meter [2]
%N	0.17	Kjeldahl distillation method [3]
%C	1.69	Walkley-Black method [4]
C:N	10.1	
OLSEN P (MG KG ⁻¹)	9.88	Olsen method [5]

KIPSANGUI SITE

PARTICULAR	VALUE	METHODS
SAND %	60	Hydrometer method [1]
SILT %	18	
CLAY %	22	
TEXTURAL CLASS	SANDY LOAM	
PH (1:2.5 SOIL: WATER)	5.32	Glass electrode pH meter [2]
%N	0.28	Kjeldahl distillation method [3]
%C	2.66	Walkley-Black method [4]
C:N	10.1	
OLSEN P (MG KG ⁻¹)	10.37	Olsen method [5]

3.2 Effects of lime additions on soil pH during wheat growth


The effect of application of lime on the pH values of the soils taken at different time intervals of 46 days after planting and at harvesting for the Chepkoilel and Kipsangui sites are given in Figure 1 and 2. There was an increase in soil pH at both site and sampling dates with lime additions of 1.0, 1.5 t ha⁻¹ and 2.0 t ha⁻¹ in both wheat varieties. This was because lime likely increased the Ca²⁺ ions, which it contains, displaced the Al³⁺, H⁺, Fe³⁺ ions, prevalent in acid soils such as described by [14,17, 23]. Increased soil pH, due to lime, nitrogen and phosphorus has been reported by others and attributed to the effect of applied [24, 25]. Soil amendment material.

Comment [SD6]: there was no pH increase at a rate of 1 ton/ha at Chepkoilel with Mwamba- why? I wonder if the lime was applied shortly before the trial, or that it was just surface applied, so that it didn't have enough time to react with the soil? or was the lime not fine enough?

Comment [SD7]: it is strange that higher lime rates didn't increase pH more than low lime rates - explanation?

Formatted: OMlist, Tab stops: Not at 1.56"

Chepkoilel site-Njoro BW 2 wheat variety



Chepkoilel site-KS Mwamba wheat variety





Figure 1: Soil pH changes during the 2009 LR season for Chepkoilel site as affected by treatment application Error bars indicate standard error of means.

Kipsangui site-Njoro BW 2



Kipsangui site-KS Mwamba




Figure 2: Soil pH changes during the 2009 LR season for Kipsangui site as affected by treatment application Error bars indicate standard error of means.

Comment [SD8]: it seems lime didn't change pH at this site

3.3. Effect of lime on soil available P taken at wheat harvest

The results of available P taken at harvesting for Chepkoilel and Kipsangui sites as affected by lime additions for the two wheat varieties are shown in the Figure 3. The data were analysed across both experimental sites. There was a significant difference ($p=0.05$) of soil available P from treatments above the control with increase in the lime rate applied in both varieties and sites. Lime application increased both the native soil P and P fertilizer availabilities which were taken up by plant possibly decreased soil sorption capacity (not measured). Lime has been reported to reduce the P sorption in acid soils resulting in increased available P for plants [26], [27]. In highly weathered soils of the tropical and subtropical acid soils, the applied P fertilizers readily react with Al and Fe sesquioxides to form sparingly soluble P forms. This normally results in very low soil available P for plant absorption [28], [29]. The soil available P at Kipsangui site was higher than that at Chepkoilel site in both wheat varieties with the increase of lime levels. This could be attributed to the initial soil fertility status. The soil available P (P mg kg^{-1}) at both sites was lower in the control treatment when compared to all levels of lime additions. The soil available P was highest with lime addition of 2 t ha^{-1} in both varieties and sites. This could be attributed to the initial soil fertility status, where Kipsangui site had more fertile soil as compared to Chepkoilel site.




Figure 3. Changes in soil available P (mg kg^{-1}) at harvesting due to lime and P application during the 2009 LR for Chepkoilel and Kipsangui sites 125 days after application. Error bars indicate standard error of means.

Comment [SD9]: the y-axis label states N-uptake - is that a mistake?

3.4 Effects of soil amendments on wheat yield

Yields obtained from the test crop as a result of application of soil amendment materials; P, N and lime during the year 2009 LR are given in Figure 4 for Chepkoilel and Kipsangui sites. Amendment of soil acidity with lime plus P and N addition, increased grain yield significantly ($p=0.05$) in Chepkoilel and Kipsangui. The lowest grain yield in Njoro BW 2 wheat variety of 1.18 t ha^{-1} in Chepkoilel site and 1.27 t ha^{-1} in Kipsangui site were found on control where

Comment [SD10]: i hope that only lime was the changing variable - otherwise your study cannot distinguish between lime and fertilizer effects.

there was only P and N additions without lime. The lowest grain yields in KS Mwamba variety of 0.78 t ha^{-1} at Chepkoilel and 0.84 t ha^{-1} at Kipsangui sites were found on the control treatment. This is in agreement with [30] who stated that combined application of lime with phosphorus significantly increased grain yield in Kenya. However, there was no significant difference in the grain yield between the two varieties with addition of 1.0, 1.5 and 2 t ha^{-1} at Chepkoilel. This trend also confirms the fact that if the growth of wheat at any stage is limited by a specific factor such as nutrient supply, water, light or temperature, then the grain yield is limited irreversibly unless it can be compensated by modifying a yield component occurring at a later stage of development [31]. During the year 2009 Kipsangui area received 1031 mm of rainfall while Chepkoilel received 817 mm. At the grain filling stage, rainfall received in 30 days at the two sites was 144.5 mm and 47.5 mm for Kipsangui and Chepkoilel respectively. The difference in the amounts of rain received at both sites is reflected in the performance in the grain yield.




Figure 4.Wheat grain yields (t ha^{-1}) per site as affected by treatment application for both sites during the long rains 2009. Error bars indicate LSD _(0.05).

3.4 Correlation between wheat yields and soil available P at wheat harvest

Correlation analysis was conducted between wheat yields and soil available P at harvesting and the results obtained are given in Figure 5 and 6. There was a highly positive and significant correlation between soil available P and grain yields i.e. $r = 0.97$ and $r = 0.94$ for KS Mwamba wheat variety at Chepkoilel and Kipsangui sites respectively. The correlation between soil available P and grain yields was also positive i.e. $r = 0.66$ and $r = 0.97$ for Njoro BW 2 wheat variety at Chepkoilel and Kipsangui sites respectively. The high positive correlation is probably because lime increased the overall pH which led to the enhanced availability of phosphorus. Soil reaction (pH) affects the physical, chemical and biological properties of soils and crop yields. Soil acidity has a negative effect on crop yields mainly through reduced P availability though Fe and Al fixation of P [32]. This is in line with the findings of [33] found that the increase of soil pH resulted in a significant increase in grain yields. Grain yields responded to lime addition in acid soil but at different magnitudes. Further, according to [34] findings better grain yield was obtained when lime was applied in combination with fertilizer.

Comment [SD11]: it seems to me the lime didn't change the pH - was this the reason for limited yield increases?

Comment [SD12]: are you claiming excessive drought?

Comment [SD13]: your data don't show increases in pH with liming







Figure 6. Relationship between soil available P (mg kg⁻¹) and wheat grain yields (t ha⁻¹) as observed at harvesting during 2009 LR for Chepkoilel site for Njoro BW 2 and KS Mwamba.



r=correlation coefficient

Figure 6. Relationships between soil available P(mg kg⁻¹) and wheat grain yields (t/ha-1) as observed at harvesting during 2009 LR for Kipsangui site for Njoro BW 2 and KS Mwamba.

3.4 Uptake of P and N in wheat

Figure 7 and 8 show the effect of lime on the wheat grain N and P uptake. Lime had a significant ($p=0.05$) effect on grain P and N uptake however, there was no significant

increase on P uptake. Soil amendment (lime) increased N uptake in both wheat grain at both sites. The mean P uptake for the wheat grain in Njoro BW 2 variety was 6.39 kg P ha^{-1} and 5.35 kg P ha^{-1} for Chepkoilel and Kipsangui sites respectively means from all treatments. Increased P uptakes with lime, P and N fertilizers have been reported by other workers [35]. This has been attributed to better soil amelioration by the amendments [36]. According to [37], acidic conditions reduce the rate of release of mineral-N from organic-N. Liming however, increases the rate of mineralization and hence improves the supply of mineral-N to the plants.




Figure 7 Nutrient P uptake (kg P ha^{-1}) for wheat straw as affected by treatment application for Chepkoilel and Kipsangui sites. Error bars indicate standard error of means.

means.





Figure 8. Nutrient N uptake (kg P ha^{-1}) for wheat grain as affected by treatment application for the Chepkoilel and Kipsangui sites. Error bars indicate standard error of means.

3.5 Economic Analysis

Table 2 and 3 show economic analysis data of benefits, costs and marginal rate of return of treatments using partial budget techniques. The GFBs, NFBs, and TVC to investment were different among the sites. Treatments that produced lower NFBs were not worth for investment. They are known as dominated and were marked "D".

Most of the treatments did not realize economically viable returns. However, at Kipsangui site, for Njoro BW 2 wheat variety with treatment combination of 40 P₂O₅ + 46 N + 2.0 tons KL/ha⁻¹ and NFBs of Ksh 34,010 was the only viable treatment option (Table 3). At Chepkoilel site there was no viable treatment. Based on economic returns, the best treatment in Kipsangui site was Njoro BW 2 variety with 2 t/ha⁻¹ Koru lime. In this study the majority of the farm inputs for example DAP did not realize economically viable returns to investment due to high cost of inorganic inputs, low wheat prices offered and poor distribution of rainfall in the year 2009.

Table 2: Economic analysis of wheat yields of the soil amendment materials incorporated into the soil during 2009 LR for Kipsangui site

Treatments	GFP(Ksh)	TVC (Ksh)	NFB (Ksh)	MRR(%)
Njoro BW 2 wheat variety				
40 P ₂ O ₅ + 46 N + 0.0 tons KL/ha	34,290	5,786	27,926	
40 P ₂ O ₅ + 46 N + 0.5 tons KL/ha	42,660	15,292	25,838	D
40 P ₂ O ₅ + 46 N + 1.0 tons KL/ha	51,570	20,183	29,369	9
40 P ₂ O ₅ + 46 N + 1.5 tons KL/ha	56,970	24,482	30,040	14
40 P₂O₅ + 46 N + 2.0 tons KL/ha	66,420	29,494	34,010	72
KS Mwamba wheat variety				
40 P ₂ O ₅ + 46 N + 0.0 tons KL/ha	22,680	3,827	18,471	
40 P ₂ O ₅ + 46 N + 0.5 tons KL/ha	27,000	12,650	13,085	D
40 P ₂ O ₅ + 46 N + 1.0 tons KL/ha	40,500	18,315	20,353	12
40 P ₂ O ₅ + 46 N + 1.5 tons KL/ha	48,060	22,978	22,784	47
40 P ₂ O ₅ + 46 N + 2.0 tons KL/ha	54,540	27,459	24,353	31

Table 3: Economic analysis of wheat yields of the soil amendment materials incorporated into the soil during 2009 LR for Chepkoilel site

Treatments	GFB(Ksh)	TVC (Ksh)	NFB (Ksh)	MRR(%)
Njoro BW 2 wheat variety				
40 P ₂ O ₅ + 46 N + 0.0 tons KL/ha	31,860	5,376	25,947	
40 P ₂ O ₅ + 46 N + 0.5 tons KL/ha	36,720	14,290	21,001	D
40 P ₂ O ₅ + 46 N + 1.0 tons KL/ha	44,280	18,953	23,432	D
40 P ₂ O ₅ + 46 N + 1.5 tons KL/ha	49,410	23,206	23,883	D
40 P ₂ O ₅ + 46 N + 2.0 tons KL/ha	46,980	26,184	18,178	D
KS Mwamba wheat variety				
40 P ₂ O ₅ + 46 N + 0.0 tons KL/ha	22,680	3,827	18,470	
40 P ₂ O ₅ + 46 N + 0.5 tons KL/ha	27,000	12,650	13,085	D
40 P ₂ O ₅ + 46 N + 1.0 tons KL/ha	32,400	16,949	13,756	D
40 P ₂ O ₅ + 46 N + 1.5 tons KL/ha	38,880	21,430	15,308	D
40 P ₂ O ₅ + 46 N + 2.0 tons KL/ha	46,980	26,184	18,178	D

GFB=Gross field benefits, TVC = Total variable cost, NFB = Net financial benefits, MRR = Marginal rate of return, KL = Koru Lime as CaO, D = dominated treatment (i.e. with less than or equal to treatment with lower TVC that were eliminated from further consideration since

no farmer choose a treatment(s) with higher TVC and receive lower NFB), bold and underlined indicate economically viable treatment.

4. CONCLUSION

Acidity in the soils is a serious health problem. Soil acidity may negatively affect nutrient availability and cause Al and Mn toxicity. Integration of lime applications and phosphorus improved soil conditions resulting in increase in soil pH, available P, yield and nutrient uptake (N and P). The combination between Njoro BW 2 variety and soil amendment at the rate of 2 t ha⁻¹ lime (plus P and N) gave the highest grain yield and thus viable economic returns. High production costs, particularly of inorganic inputs (P fertilizer, foliar feed and lime) and relatively low prices of wheat in year 2009, made the majority of the treatments not realize economically viable returns for adaptation by SHF. Further long-term studies in these soils to investigate the effects of lime and fertilizer on wheat yield, as a basis for fertilizer formulations and recommendation is necessary.

Comment [SD14]: i don't think your data show this.

ACKNOWLEDGEMENTS

The authors would like to thank Moi University Research Fund for partially funding research work. Special thanks to laboratory and field technicians for assisting in the laboratory and field work. Most appreciation to the farmer Mrs Selina Maswai and School of Agriculture and bio-technology, University of Eldoret for allowing the research to be conducted on their farms.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

Authors may use the following wordings for this section: " 'Author 1', study conception and design performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. 'Author 2' 'Author 3' and 'Author 4' managed the analyses of the study. 'Author 4' interpretation of data' Author 5' and 'Author 6' managed the literature searches..... All authors read and approved the final manuscript."

REFERENCES

1. Ministry of Agriculture. Grain production in Kenya. Kilimo house, Nairobi Kenya. 2005
2. Osongo, M. Kenya Grain and Feed Wheat Update. Information network grain report, Nairobi, Kenya. 2003.
- 3 Ministry of Agriculture. Crop and Food situation Annual Report, Uasin Gishu district, Eldoret. Kenya. 1996.
- 4 Ministry of Agriculture. Crop and Food situation Annual Report, Uasin Gishu district, Eldoret. Kenya. 2009
- 5 FUR. Fertilizer Use Recommendation Project – Fertilizer Recommendations vol. 1-22. Kenya, Agricultural Research Institute (KARI)/Ministry of Agriculture. 1994.
6. Kenya Roads Authority. Northern Corridor Rehabilitation Programme Phase 111. Environmental and Social Impact Assessment for Rehabilitation of

- Timboroa- Eldoret Road (ACOA). 2010.
7. Lwayo, M.K., Okalebo, J.R., Muasya, R.M. and Mongare, P.O. A diagnostic Survey on the Production Constraints and Utilization of phosphate fortified wheat straw and maize stover compost for increasing cereal production in Uasin Gishu district, Kenya A Report, 2001. 108 pp.
 8. Neil,C., Claudi, N, (Ed). Ground works1: Managing soil acidity. In; Claude Proceeding of Tropical Soils Workshop on soil acidity and liming. Raleigh, NC.1991. pp1-23
 9. Kochian, L.V., Hoekenga, O.A. and Pineros, M.A. How do crop plants tolerate acid soil Mechanisms of aluminum tolerance and phosphorus efficiency. *Ann. Rev. plant biol.*, 2004. 55: 459- 493
 - 10 Kanyanyua, S.M., Ireri, L., Wambua, S and S.M. Nandwa S.M. *Acidic soils in Kenya; Constraints and Remedial options*. KARI Technical Note No. 11 ;KARI Headquarters, Nairobi, Kenya.2002
 11. Ligeyo, D.O. and Gudu, S.O. Further laboratory screening of More Kenya Maize Inbred lines for Tolerance to Aluminum In: Third year Progress Report, (March 2004 to 28 February 2005). Mcknight Foundation USA Project,EMBBRAPA, Purdue and Cornell University (USA) and Moi University(Kenya) Phase1, 2003-2005
 12. Swift, M.J., Dvorak, K.A., Mulongoy, K., Musoka,M., Sanginga,N. and Tian,G. The role of soil organism in the sustainability of tropical cropping systems In; Syers,S.K. and Rimmer, D.I. (Eds).Soil science and sustainable land management in tropics.CAB International, Cambridge University Press U.K. 1994.pp 155-170.
 13. Anetor, M. O and Ezekie, A. A. "Lime effectiveness of some fertilizers in a tropical acid alfisol", University of Ibadan, Ibadan, Nigeria. 2007.
 14. The, C., Calba H., Zonkeng, C., Ngonkeu, E.I.M., Adetimirin, V. Response of maize grain to changes in acid soils characteristics after soil amendment. *Plant and soil*, 2006. 28:45-57.
 15. Kisinyo, P. Constraints of soil acidity and nutrient depletion on maize (*Zea may*) production in Kenya. D. Phil Thesis. Moi University, Eldoret, Kenya. 2011
 - 16.Kamprath, E.J.Crop response to lime on soils in the tropics, *Agronomy and soil Science Society of America*, Madison Wisconsin, USA In: Adams F. (Ed). *Soil Acidity and Liming*. 2nd Edition. Agronomy Monograph 12. 1984b.pp 349–368.
 17. Moody, P.W., Aitken,R., and Dickson,T.Field amelioration of acid soils in south-east Queensland 111. Relationships of maize yield response to lime un-amended soil properties. *Australian Journal of Agricultural Research*,1998. 49(4):649-656.
 18. Jaetzold, R. and H. Schimdt. Farm management Handbook of Kenya. Vol II. Natural conditions and Farm management information. Part A.Western Kenya (Nyanza and Western provinces). Kenya Ministry of Agriculture Team, Nairobi Kenya, 1983. p397

19. Jaetzold, R., Schmidt, H., Hornetz, B and Shisanya, C. Farm management handbook of Kenya- natural conditions and farm management information (2nd Edition). Volume II/A western Kenya. Ministry of Agriculture, Nairobi.2006.
20. Acland, J.D. *East African Crops*. Longman Group, London, U.K.1971.
21. Mamo.M, Wortmann .S.S, Shapiro C.A. Lime use for soil acidity management. Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln Extension, United States Department of Agriculture.2009.
22. Ndungu, K .W., J.R. Okalebo, C.O. Othieno, M.N. Kifuko, A.K. Kipkoech and L.N. Kimenye. Residual effectiveness of Minjingu Phosphate Rock and fallow Biomass on crop yields and financial returns in western Kenya. *Expt. Agric.* 2006. Vol 42, pp323-336.
23. Bado, B.V., Segedo, M.P. and Lompo, F. (2004). Long term effects of mineral fertilizers, phosphate rock, dolomite and manure on the characteristics on an ultisol and maize yield in Burkina Faso, pp 77-88. In: Bationo, A. (ed). Managing nutrient cycles to sustain soil fertility in Sub- Saharan Africa. Academic Science Publishers. Nairobi Kenya.
24. Ernani, P.R., Cimelio, B. and M. Leonardo, M. Corn yield as affected by liming and tillage system on an acid Brazilian Oxisol. *Odiat Agron. Journ.* 2002. 94: 305 – 309.
25. White, J.R, Bell M.J. and Marzies, N.W. . Effect of subsoil acidity treatments on the chemical properties of a ferrosol. *Proc. Agron. Conf. Aust.* 10th – 15th Sept. 2006 pp 10 -15.
26. Tisdale, S.L., Nelson, W.L., and Beaton, J.D. Soil fertility and fertilizers. 5th edition Macmillan, New York, U.S.A.1990
27. Van Straaten, P. Agro geology: the use of crops. Enviroquest Ltd, 352 River Road, Cambridge, Ontario N3C 2B7, Canada.2007.
28. Keerthisinghe G., Zapta, F., Chalk, P. and Hocking, P. Integrated approach for improved P nutrition of plants in tropical acid soils, 2001. pp 974-975. In: Horst, W, J.Schenk, M.K., URKERT, A.B., Claaseen, N., Flessa, H., Frommer, W.B., Gildbach, H., Olf, H.-W., omheld, V.R., Sattlmacher, B., Schmidhalter, U., Schubert, S., Wiren, N.V. and Wittmayer, L. (Eds). Plant Nutrition-Food Security and Sustainability of Agr- ecosystems. Dordrecht: Springer.
29. Kochian, L.V. Cellular mechanism of aluminum toxicity and resistance in plants. *Ann. Rev. Plant Physiol.*, 1995. 46, 237 – 260.
30. Okalebo, J.R., Gathua, K.W., Woomer, P.L. Laboratory methods of soil and plant analysis: A working Manual. (2nd Ed). TSBF-CIAT and SACRED Africa, Nairobi, Kenya.2002.
31. Heyland, A. Werner; Lehrstuhl fuer Speziellen P flauenbav Und P flanzewzuechtung,. Wheat (*Triticum aestivum* L., T. durum Desf.

- University of Bonn, Germany..1999
32. Okalebo, J.R. Recognizing *the Constraint of Soil Fertility Depletion and Technologies to Reverse it in Kenya Agriculture*. Inaugural Lecture 6 Series NO.1, Moi University, Kenya.2009.
 33. He,Z.I., Baligar, V.C., Martens, D.C., Ritchey, K.D. and Kemper, W.D. . Factors affecting phosphate rock dissolution in acid soil amended with liming material and cellulose. *Soil Science Society of America journal*, 1996. 60: 1596-1601
 34. Anetor, M. O and Ezekie, A. A. "Lime effectiveness of some fertilizers in a tropical acid alfisol", University of Ibadan, Ibadan, Nigeria. 2007.
 35. Sharma, P.K, N. San kavan, S.K. Bhardwaj and S.P. Dixit. Studies on root characteristics, biomas productivity of wheat (Aradhana) in phosphorus, farm yard manure and mulch amended mountain acidic lands of western himalaya's India. *Proc. 17th WCSS*, 14 – 21 August, 2002. No. 22. Paper no. 2016
 36. Busari, M.A, Salako, F.K., Sobulo, R.A., Adetunji, M.A. and Bello, N.J. *Variation in Soil pH and Maize Yield as Affected by the application of Poultry Manure and Lime*. In *Managing Soil Resources for Food and Sustainable Environment*. Proc. of 29th Annual Conf. Soil Sci. Soc. Nig. 2000. Pp 139 – 142.
 37. Miranda, L.H.De and D.L. Rowell. *The effects of lime and phosphorus on the function of wheat roots in acid tropical soils and subsoils*. *Plant and Soil* 1987. 104 p 253 -262.