

DIETARY INFLUENCES OF *Aspilia africana* ON LITTER TRAITS OF BREEDING RABBIT DOES

Comment [i1]: It is written FEMALE RABBITS instead of RABBIT DOES

Comment [i2]: In the title should not be abbreviations

NseAbasi N. Etim^{1*}, Joseph S. Ekpo¹ and Glory E. Enyenihi²

¹Department of Animal Science, Akwa Ibom State University.
P.M.B. 1167, Uyo, Akwa Ibom State, Nigeria.

²Department of Animal Science, University of Uyo,
P.M.B. 1017, Uyo, Akwa Ibom State, Nigeria.

ABSTRACT

The study examined the effect of feeding *Aspilia africana* on litter traits of breeding does in a replicated completely randomized design at College of Animal Science and Animal Production, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria, between March 2009 and June, 2009. Thirty dutch breed rabbit does aged 5 to 6 months were used for the experiment. The treatments consisted of mixed forages (Centrosema pubescens (200g)), Ipomea batatas leaves (100g) and Panicum maximum (200g) without *Aspilia africana* (T₁; control), fresh *Aspilia africana* (500g/doe/day) and wilted *Aspilia africana* (500g/doe/day). The rabbits were fed the same concentrate diet (300g/animal/day) throughout the study and mixed forages from the commencement of the experiment till the does kindled. After parturition, fresh and wilted *Aspilia africana* forages were introduced in T₂ and T₃ respectively while the control continued on mixed forages throughout the study. The result of the study revealed no significant differences ($P > .05$) in the litter sizes at birth and at weaning among the various treatment groups, though T₁ had the highest numerical mean value at birth (5.60) and lowest at weaning (3.70). Litter weight at birth revealed significant differences ($P < .05$) in which T₁ recorded the highest mean litter weight (261.50g) while T₂ and T₃ weighed 116.50g and 175.00g respectively. At weaning, T₂ recorded significantly higher ($P < .05$) litter weight (736.30g), followed by T₃ (621.30g) while the control group (T₁) weighed the least (410g). The weekly litter weights from birth to weaning showed no significant differences ($P > .05$) in week 1 but from weeks 2 to 4 there were significant differences ($P < .05$) in which litters in T₂ had the highest values, 534.38g, 690.60g and 736.30g while T₁ recorded the lowest mean values of 297.50, 405g and 410g respectively indicating the *Aspilia Africana* groups had superior growth rate to the control group. This study revealed that *Aspilia africana* has the potential for increasing body weights of litters during lactation.

Comment [i3]: The meaning of T₂ and T₃ should be written

Keywords: *Aspilia africana*, rabbit does, litter traits, forage process

1. INTRODUCTION

One of the problems facing Nigeria and most developing countries is scarcity of food for the teeming population and feed for the dwindling livestock industry [1]. The relevance of animal protein in human and animal nutrition in Nigeria cannot be overemphasized. In recent times, there has been a significant short fall between the production and supply of animal protein to feed the ever increasing population [2]. Animal protein content in the diets of low income earners that constitute the majority of Nigerian populace is very low [3]. Average consumption of animal protein in this country is estimated at 4.5g/head/day as against a minimum requirement of 35g/head/day recommended by the Food and Agriculture Organization of the United Nations [4]. The level of nutrition greatly influences productivity in livestock, when adequate; it increases animal performance and makes animal products available for human use. Conversely inadequate quantity and quality of feed is known to be largely responsible for low livestock productivity in the tropics [5]. Various plans and agricultural policies have been formulated at one time or the other to combat the acute shortage of animal protein supply [6]. The production of the conventional protein and energy sources is still grossly inadequate in most developing countries of the world and often times demand exceeds supply [7]. This is reflected in the competition between man, livestock and industries for existing conventional feedstuffs which has led to high costs of these ingredients which consequently has contributed to the existing high prices of animal protein for the average Nigerian [8]; [9]. This situation warrants the evaluation and use of neglected novel plants that are cheap and locally available in feeding animals [10];[11]. One of such plants are forages. Proper processing of forages is yet another link in the chain of controlling quality of feedstuffs. [12] One of the forages suitable for feeding rabbits is *Aspilia africana*. *Aspilia Africana* is high in protein, vitamins and minerals. It is reported that *Aspilia Africana* contains ascorbic acid, riboflavin and thiamin. It is a good source of minerals such as Ca, P, K, Mg, Na, Fe and Zn [13]. The plant is also reported to be high in anti-nutritional factors such as alkaloids, saponins, flavonoids, phenols and tannins [13], some of which might be reduced through drying/wilting.

Moreover, to meet the increasing demand for animal protein, emphasis needs to be given to non-conventional sources as against the conventional sources such as cattle, sheep, goat, pig and poultry that would require more capital, space and time [3]. Efforts have to be directed towards boosting the micro-livestock sector [2]. There is need to focus attention on the production of livestock whose nutritional requirement does not put much strains on the limited sources of feed ingredients to which men also subscribe. An example of such livestock is rabbit. [14] stated that if efforts are intensified in encouraging and motivating rabbit keepers to increase their stock and adapt more rabbit innovations, this will undoubtedly increase rabbit production and ensure adequate nutrition for every household through animal protein consumption. This will enable developing countries like Nigeria attain household food and nutritional sufficiency. The means through which the aforementioned can be achieved lies with an improvement on litter traits of breeding does.

Nutritional status, parity, and environment among other factor were identified by [15] as influencing litter traits of breeding does. The litter traits of rabbits are very important features that determine the efficiency of rabbit production [16]; [17]; [18]. According to [19] breeder could influence the litter size and litter weight at weaning by the food provided for the nursing doe. Increased feed utilization is reported to result in a higher litter birth weight brought by high milk production of does [20]. [21] reported that *Aspilia africana* is used as lactation stimulant. [15] reported that does fed *A. Africana* had higher milk production. [19] observed rapid growth of kits during the peak of lactation coincides with three weeks of lactation. [22] reported that after the third week, milk production gradually declines. [23] reported that the mean daily weight gain of rabbits fed *A. africana* at four weeks of age was significantly greater than others. [23] further observed that feeding *A. africana* or *Centrosema pubescens* to rabbits at four weeks will enhance their performance.

There is a paucity of information on the effects of *Aspilia africana* on litter traits of breeding does. Therefore, this study was carried out to evaluate the feeding value of *Aspilia africana* on performance and growth of rabbit litters.

2. MATERIALS AND METHODS

2.1 Experimental Location

The study was conducted in the Rabbitry Unit of the Teaching and Research Farm of the College of Animal Science and Animal Production, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.

2.2 Experimental Animals and Management

Thirty sexually mature nulliparous dutch breed rabbit does and four bucks aged 5 to 6 months were used for the study. The rabbits were certified healthy and were identified with ear tagged for ease of identification. Both sexes were separated. Two weeks pre-experimental period was allowed to enable the animals adjust to the new environment. They were housed singly in rabbit hutches made of wooden frame and wire mesh. The animals were fed concentrate diet 300g/animal/day (Table 1) through out the period of the trial and mixed forages (*Centrosema pubescens* (200g/animal/day), *Ipomea batatas* leaves (100g/animal/day) and *Panicum maximum* (200g/animal/day) at the ratio of 2:1:2 until the start of the experiment. The does were mated two weeks after commencement of the experiment. The mating ratio was 1 buck: 10 does. The does were taken to the bucks' pen for mating and were all mated within an interval of 3 days. The does were palpated after 14 days of mating to confirm pregnancy. Following kindling, *Aspilia africana* forage was introduced as the experimental diet at the rate of 500g per animal per day.

Table 1. Percent composition of concentrate ration

Ingredients	T ₁	T ₂	T ₃
Maize offal	45.5	45.5	45.5
Palm kernel cake	30	30	30
Soybean meal	20	20	20
Blood meal	2.0	2.0	2.0
Bone meal	2.0	2.0	2.0
Vitamin-mineral Premix	0.25	0.25	0.25
Salt	0.25	0.25	0.25
	100	100	100

Comment [i4]: It should be given the content of the premix below the table.

Metabolizable Energy = 2620Kcal/kg Calcium = 1.10%
Crude protein = 18.0% Phosphorus = 0.80%

2.3 Experimental Design and Data Collection

The experiment was a completely randomized design with three treatments. The treatments consisted of mixed forages (*Centrosema pubescens*, *Ipomea batatas* leaves and *Panicum maximum*) without *A. africana* (control; T₁), fresh *Aspilia africana* (T₂), wilted *Aspilia africana* (T₃). The proximate composition of *Aspilia africana* is presented in Table 2 and that of the control forages (*Centrosema pubescens*, *Ipomea batatas* leaves and *Panicum maximum*) in Table 3. Concentrate diet (300g/animal/day) was offered to does in all the treatment groups during the trial. Ten does were randomly assigned to each treatment, each of which was replicated five times with two does per replicate.

Following kindling, data were collected on the litter size, litter birth weight and weekly weight gain of the litters measured using kitten scale. Then at weaning; litter size and litter weight were measured.

Table 2. Proximate Composition of *Aspilia africana*

Components (%)	Percent Composition
Crude Protein	23.0
Nitrogen Free Extract	43
Crude fibre	20.2
Ether Extract	3.0
Ash	8
Metabolizable Energy (Kcal/kg)	2920

Adapted from: Ugwuene and Ojewola (2009)

Comment [i5]: It should be added.

Table 3. Proximate Composition of the Control forages (*Centrosema pubescens*, *Ipomea batatas* leaves and *Panicum maximum*)

Components(%)	<i>Centrosema pubescens</i>	<i>Ipomea batatas</i> leaves	<i>Panicum maximum</i>
Crude Protein	22.8	13.2	8.5
Nitrogen Free Extract	46	45.5	39.2
Crude fibre	21	8.41	25.1
Ether Extract	2.3	2.8	1.0
Ash	7.0	11.8	8.0
Metabolizable Energy (Kcal/kg)	2856	2672.4	2810

Comment [i6]: It should be added.

2.4 Data Analysis

The data that were generated were subjected to the Analysis of Variance (ANOVA) [24] for a completely randomized design (CRD). Significant means were separated using Least Significant Difference (LSD) according to the methods of [24].

Comment [i7]: It should be deleted

3. RESULTS AND DISCUSSION

The results of the dietary effects of *Aspilia africana* on litter traits of experimental breeding rabbit does are presented in Table 4. Mean litter size for the group assigned to T_1 was higher than, the others followed by that of the group assigned to T_2 . The group placed on T_3 had the lowest litter size. Nevertheless, there was no statistically significant difference ($P > .05$) in litter size between all the treatment groups. The study recorded no still birth.

The litter sizes recorded in this study was lower than 7 to 8 kits reported by [25] and within the range of 2 to 9 kits observed by [17]. The low litter size of the study could be as a result of the does being in their first parity, since it has been shown that litter size at birth increase with successive kindling [19]; [26].

The mean litter size at weaning was statistically not different ($P > .05$), but varied numerically. The mortality rate in T_1 (control) was higher than in T_2 while T_3 (treated with *Aspilia africana*) recorded no mortality. Thus, there was a relatively high decrease in litter size at weaning of the control compared with the treated groups. This could be attributed to high milk production from does in T_2 and T_3 which were fed fresh and wilted *Aspilia africana* respectively. This is in line with the report by [15] that does fed *Aspilia africana* had higher milk production. The test plant (*A. africana*) helped stimulate milk let down and increase milk yield which thereby increased the survival of the litter and reduced preweaning mortality. This in turn resulted in higher litter size at weaning for T_2 while T_1 had the least although at birth T_1 had the highest litter size.

Litter weight at birth as shown in Table 4 were significantly different ($P < .05$) among the various treatment groups, which T_2 had the lowest mean body weight at birth and T_1 had the highest. This could be due to litter size at birth since T_1 had the highest litter size at birth. This agrees with the findings of [27] who reported that litter size exerts a positive effect on litter weight. The result of this study is also in line with the findings of [17] who stated that litter weight at birth depends on the breed and a mean birth weight per individual. However, the litter birth weight for T_1 may also be due to the ability of the rabbits in that group to effectively utilize the feed which enhanced better embryonic development as observed by [20] that increased feed utilization will result in higher litter birth weight. The overall litter birth weight obtained in this study is lower than 223.3g obtained by [17] and also lower than 246.77g obtained by [20]. The differences could be attributed to higher litter sizes the authors obtained at birth.

The mean litter weaning weight was highest in T_2 followed by T_3 while T_1 had the lowest. The mean weaning weights were significantly different ($P < .05$). Higher weaning weights in treatments 2 and 3 may be due to the role of the experimental plant as lactation stimulant as reported by [21] and the ability of the does in these treatment groups to effectively utilize minerals and other nutrients in the feed to increase the quantity/quality of milk for proper nursing of the kittens which thereby resulted in the higher litter weaning weight. The difference in mean weaning weights of litters in T_3 compared to T_2 could be due to level of utilization, differences in palatability, nutrient content and other unknown factors. While on the other hand, low litter weaning weights observed for the control group (T_1) could be as a result of low milk let-down from the does. This agrees with the findings of [19] that the breeder could influence the litter size and litter weight at weaning by the feed provided for the nursing doe.

199 **Table 4.** Effects of *Aspilia africana* on litter traits

Parameters	T ₁	T ₂	T ₃	SEM
Average litter size at birth	5.60	5.20	4.00	0.40
Average litter size at weaning	3.70	4.60	4.00	0.45
Average litter weight at birth (g)	261.50 ^a	116.50 ^b	175.00 ^{ab}	23.70
Average litter weight at weaning (g)	410.00 ^b	736.30 ^a	621.25 ^a	47.17

200 a,b,c, means in same row with different superscripts are significantly different ($P < .05$).

Comment [i8]: It should be deleted

201
 202 The results of the effect of *Aspilia africana* on weekly litter weights from birth to weaning are
 203 presented in Table 5. The body weights of the kits in week 1 showed no significant differences ($P >$
 204 $.05$) between the different treatments. Kits in T₂ had the highest mean value, followed by T₃ while T₁
 205 had the lowest mean value. By the second week, the weight gain of litters in T₁ was significantly lower
 206 than those of T₂ and T₃. However, the rate of gain in all the treatment groups in weeks 3 and 4 were
 207 higher than those in weeks 1 and 2 and T₂ had the highest mean value. The sharp increase in weights
 208 of kits at three weeks postpartum agrees with the findings of [20] who observed that rapid growth rate
 209 of kits during the peak of lactation coincides with three weeks of age from birth. Furthermore, it could
 210 be observed that T₂ and T₃ showed superior increase compared to T₁ (control) in week 3 which may
 211 be as a result of high milk yield/let-down from the mammary gland of does which was stimulated by
 212 the test plant (*Aspilia africana*) as reported by [25]. The mean weekly body weight gain for week 4
 213 also showed significant difference ($P < .05$) between the different treatments in which T₂ had the
 214 highest mean value, followed by T₃ and T₁ (control) still had the lowest value. The slight difference in
 215 the weights of kits in T₃ whose dam were fed wilted *A. africana* compared with T₂ which received fresh
 216 *A. africana* could be as a result of differences in palatability and nutrient loss during wilting. The higher
 217 weight gains for T₂ and T₃ groups could be due to better nutritive value of the test plant (*Aspilia*
 218 *africana*) since the does in all the treatment groups consumed all the concentrate and forages offered
 219 them without remnants and thus had similar feed intake/consumption. The results of this study agrees
 220 with the findings of [26] who reported that the mean daily weight gain of rabbits fed *A. africana* at four
 221 weeks of age was significantly greater ($P < .05$) than others. [26] further stated that feeding *A.*
 222 *africana* to rabbit at four weeks will enhance their performance.

223
 224
 225

Table 5. Effect of *Aspilia africana* on weekly litter weights from birth to weaning (g)

Weeks	T ₁	T ₂	T ₃	SEM
Week 1	290.00	380.63	316.25	30.94
Week 2	297.50 ^b	534.38 ^a	445.00 ^a	41.25
Week 3	405.00 ^b	690.60 ^a	584.40 ^a	43.43
Week 4	410.00 ^b	736.30 ^a	621.30 ^a	47.15

226 a,b,c, means in same row with different superscripts are significantly different ($P < .05$).

Comment [i9]: It should be deleted

227 4. CONCLUSION

228 The various parameters (litter sizes and weights at birth and at weaning and weekly litter
 229 weights from birth to weaning) studied to observe the effect of *A. Africana* on litter traits of breeding
 230 rabbit does showed that litters from does fed *A. Africana* showed superiority in values for all the
 231 parameters investigated. Thus, both fresh and wilted *Aspilia africana* improved the litter performance
 232 at weaning in rabbits. Therefore, *Aspilia africana* is favourable in lactating rabbit does feeding and the
 233 nutritive value of it is better than that of the mixed forages (*Centrosema pubescens*, *Ipomea batatas*
 234 *leaves* and *Panicum maximum*) used as control. Therefore, *Aspilia Africana* is strongly recommended
 235 for feeding breeding/lactating rabbit does to enhance the growth performance of their litters.

AUTHOR'S CONTRIBUTIONS

'Author A' (NseAbasi N. Etim) designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. 'Author B' (Joseph S. Ekpo) and 'Author C' (Glory E. Enyenihi) managed the analyses of the study. 'Author C' managed the literature searches and typesetting of the manuscript. All authors read and approved the final manuscript.

Comment [i10]: It should be deleted

REFERENCES

1. Carlini CR and Udedibie AB. Comparative Effects of Processing Methods on Hemagglutinating and Antitryptic activities of *Canavalia ensiformis* and *Carnavalia brasiliensis* seeds. J. Agric. Food Chemistry, 1997; 45, 4372-4377.
2. Akpan IA, Iwogor GS, Effiong OO and Akpanenua EJ. Growth performance of grasscutters fed diets treated with urea and urine solutions. Proc. of 34th Annual Conf. of Nig. Soc. for Anim. Prod., Calabar, March 18 – 21, 2009; 163-164.
3. Yusuf AM, Garba MH, Olafedehan OA and Okafor CI. Evaluation of the feeding potentials of *Vitellaria paradoxum*, *Nancteau latifolia* and *Terminalia macroptera* forage as supplements to concentrate feed for weaner rabbits. Proc. 34th Annual Conf. of Nig. Soc. for Anim. Prod., Calabar, March 18 – 21, 2009; 117-119.
4. Atsu DW. Contributing role of Animal Production in National Development. 7th Annual Conference of Animal Science Association of Nigeria (ASAN). University of Agriculture, Abeokuta. September, 16 - 17, 2002; 13-23.
5. Paterson RT. Use of trees by livestock: Calliandra, Catham, UK: National Resources Institute (NRI), 1994; 3-7.
6. Adediji OS, Amao SR, Ameen SA, Olayemi TB, Ojedapo LO and Aandiran TA. The Effects of Graded Levels of Wild Sunflower Meal on Growth and Blood Parameters of Weaned Rabbits. Proc. of the 44th Annual Conf. of ASN, "LAUTECH 2010". 2010; 737-740.
7. Matthew UD, Tegbe TS, Omege JJ and Adeyinka IA. The effect of duration of cooking pigeon pea (*Cajanus cajan*) seeds on the performance and carcass characteristics of broiler chicks. Nig. J. Anim. Prod. 2010; 37 (1):13-24.
8. Ukachukwu SN. Chemical and Nutritional Education of *Mucuna coclunhinensis* (Lyon's bean) as an alternative protein ingredient in Broiler Diets. Ph.D. Thesis, Dept. of Animal Science, Faculty of Agriculture, UNN, 2000; 1.
9. Akinmutimi AH. Effect of Cooking periods on the nutrient composition of velvet beans (*Mucuna puriens*). Proc. of 32nd Annual Conference of Nig. Soc. for Animal. Prod. Calabar, March, 18-21. 2007; 232.
10. Ahamefule FO, Ibeawuchi JA and Ahinan UG. Intake, nutrient digestibility and nitrogen balance of WAD goat fed cassava peel-leaf meals: In the new millennium; March, Umudike, Abia State, 2000; 25:71-72.
11. Etuk EB, Udedibie ABI and Obikiaonu HO. Replacement value of cooked pigeon pea (*Cajanus cajan*) seed meal for soybean meal and maize in broiler finisher diets. Proc. 7th Ann. Conf. Anim. Sci. Assoc. of Nig. (ASAN), September, 16 – 19, University of Agriculture, Abeokuta, Nig., 2002; 157-159.
12. Canadian Cattlemen's Association. Forage Processing – The National Voice of Canada's Cattlemen Association. Taking Action, August 14, 2013.
13. Etim NN, Williams ME and Ekpo JS. Carcass characteristics of rabbit does fed *Aspilia africana*. International Journal of Agriculture Innovations and Research. Volume 2, Issue 1, ISSN (Online) 2013; 2319-1473, 72-75.
14. Adisa BO and Oladoja MA. Factors associated with adoption of improved rabbit management practices and its implication on household food and nutrition security in Osun State. African Journal of Livestock Extension, 2003; 2:1-2.
15. Etim NN. Physiological and Reproductive responses of rabbit does to *Aspilia africana*. M. Sc Thesis. Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria. 2010; 30.

- 305 16. Cheeke PR. Potentials of rabbit production in tropical and sub-tropical agricultural systems.
306 Journal of Animal Science, 1986; 63:1581-1856.
- 307
- 308 17. Odubote IK and Akinokun JO. Reproduction and body weight performance of the New Zealand
309 white rabbit in humid tropics. Nig. J. Anim. Prod., 1991; 18:61-65.
- 310
- 311 18. Henry AJ, Udoh AE and Effiong OO. Reproductive performance of breeder rabbits fed diets with
312 different fibre sources. Proc. 34th Annual Conf. of Nig. Soc. for Anim. Prod., Calabar, March 18-21,
313 2009; 37-40.
- 314
- 315 19. Lebas FP, Coudert R, Rouvier and H. de Roichambeau. Food and Agriculture Organization (FAO)
316 of the United National Animal Production and Health Series, 1986; No. 21.
- 317
- 318 20. Asuquo BO. "Response of breeder rabbits to varying levels of pellet in a mixed feeding regime".
319 Nig. Journal of Anim. Prod, 1996; 23(1).
- 320
- 321 21. Burkil HM. The useful plants of West Tropical Africa, Royal Botanic garden, Kew, Vol. 1, 2nd
322 edition, Families A – D, 1985; 446-447.
- 323
- 324 22. Peace Corps. A Complete Handbook on Backyard and Commercial Rabbit Production, 1982.
325 Available: <http://www.apprpedia.org/Backyardandcommercialrabbitproductionll>
- 326
- 327 23. Ugwuene MC and Ojewola GS. The performance of weaner rabbits as influenced by time of
328 commencement of feeding different forages. Proc. of 34th Annual Conf. of NSAP, Uyo, 2009; 195-
329 197.
- 330
- 331 24. Steel RGS and Torrie JH. Principles and procedures of statistics. A biometric approach, 2nd ed.
332 McGraw Hill Book Co. Inc. New York. 1980.
- 333
- 334 25. Lukefahr SD and Cheeke PR. Rabbit Project Development Strategies in Subsistence Farming
335 System: Research Application. Wld. Anim. Rev., 1990; 68:26-35.
- 336
- 337 26. Oguike MA and Ohaja A. Effect of different protein sources on the reproduction performance of
338 rabbit does. Proc. 34th Annual Conf. of Nig. Soc. for Anim. Prod., Calabar, March, 18-21, 2009; 192-
339 193.
- 340
- 341 27. Szendro Z and Maertens L. Maternal effect during pregnancy and lactation in rabbit (a review)
342 Acta Agraria Kaposvariensis, 2001; 5(2): 1-21.
- 343
- 344 28. The American Heritage. Dictionary of the English Language, Fourth Edition. Published by
345 Houghton Mifflin Company. 2009.
- 346
- 347 29. Agromisa. Backyard rabbit farming in the tropic. Agrodo K-series No. 20. Third revised edition.
348 1999; 70.
- 349
- 350
- 351
- 352

DEFINITIONS, ABBREVIATIONS

Doe: Female rabbit

Kindling: The act of giving birth to young ones by a rabbit doe

Litter: All the young animals produced in one gestation

Litter size: The offspring produced at one birth by a multiparous mammal. E.g., rabbit, pig etc [28]

Litter weight: The (total) weight of all the off springs produced at one birth by a multiparous animal.
E.g., rabbit, pig etc

Parturition: The act of giving birth to young ones by a female animal

Weaning: The act of separating the young from their mother permanently; stopping them from
drinking the mother's milk [29]

A. africana: *Aspilota africana*

g: grams

Comment [i11]: Definitions and abbreviations should not be at the end of the text. They should be their first occurrence in the text .