

Seed and Seedling Performance of Bread Wheat (*Triticum aestivum* L.) as Influenced by Rate and In-season Nitrogen Application

Haile Deressa^{1*} and Nigussie-Dechassa R.¹

^{1*}Office of Research and Extension, Haramaya University, Dire-Dawa, Ethiopia

¹School of Plant Sciences, Haramaya University, Dire-Dawa, Ethiopia

Authors' contributions

The research experiment was designed by joint effort of the two authors. Field and laboratory activities and compilation of research results were done by the corresponding author. Both authors thoroughly evaluated the research results, and performed the statistical analysis and literature search. Both authors produced first draft of the manuscript which was thoroughly reviewed, corrected and improved by an author Nigussie-Dechassa. Both authors read and approved the final version of the manuscript.

ABSTRACT

Field and laboratory experiments were conducted in the wheat growing belt of south-eastern Ethiopia to assess effects of rate and time of N application on seed and seedling performance of local and improved bread wheat varieties. For the field experiments, a factorial combination of four N levels, two bread wheat varieties, and three times of N application were laid out in a RCBD with three replicates. Laboratory tests were conducted in a CRD with four replicates to evaluate seed germination and seedling vigor. The rate and timing of N application had significant ($P = .01$) effects on seed germination capacity and the seedling vigor index. A three times split application of N at 120 kg ha⁻¹ resulted in significantly ($P = .01$) higher percentages of normal seedlings, seed germination speed, seedling dry weight and vigor index compared to the other treatments. The results revealed that application of 120 kg N ha⁻¹ in three split doses at sowing, mid-tillering, and anthesis led to enhanced seed and seedling performance of the crop.

Key words: germination, nitrogen rate, seedlings, variety

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is one of the most important world cereal crops and is a staple food for about one third of the world's population [1]. In Ethiopia, it is grown annually on about 1.68 million ha of land with a total production of about 3.08 million t which makes the country the largest wheat producer in sub-Saharan Africa [2]. However, the average yield of wheat in the country is low (1.83 t ha^{-1}), which is only about half of the world average of 2.5 t ha^{-1} . The low crop productivity in the country may be attributed, among other factors, to low inherent soil fertility as well as meager and ill-timed N applications [3, 4]. Use of low rates of nitrogen (N) fertilizer is one of the major factors leading to low crop yields in developing countries [5]. Nitrogen deficiency has often constrained wheat production in wet areas [6, 7]. Availability of good quality seed is also an important factor limiting wheat productivity [8]. Application of N fertilizer resulted in a higher net assimilation rate [9], increased yield [10] and a higher seed protein content [11]. Timing of N application is also important for improving crop yield. Two to three equal split applications of N produced a higher grain yield of lowland rice compared with applying the whole dose of the nutrient at sowing [12]. The response of wheat to N fertilizer also varies with rate and timing of application relative to plant development [13].

Previous studies have revealed that seed quality is affected by N application with a profound influence on seed germination capacity [8, 14, 15]. The amount and timing of N application had a significant effect on seed germination speed and early vigor, with the application of N at high levels (120 and 180 kg N ha^{-1}) just before anthesis resulting in a markedly increased seed germination capacity [8]. Thus, a well-balanced supply of N throughout the growing season of the crop is essential for seed production as it has an influence, not only on yield, but also on seed quality, which further affects seed germination capacity, seedling vigor, and the productivity of the succeeding crop.

In Ethiopia, wheat seed quality is low due to genetic and management factors [16]. Therefore, it is important to enhance seed quality of the crop targeting germination capacity and seedling vigor through proper plant nutrition. Little research has been done in the country to study the effects of N application on wheat seed planting values.

Such a study is important to elucidate the influence of rate and timing of N application on seed quality of the crop. The results may lead to optimizing the quantity and timing of application of the nutrient for producing wheat seed with enhanced planting values for higher crop productivity. Therefore, this paper presents the results of a study conducted with the objective of evaluating the effect of N rate and timing of application on seed germination capacity and seedling vigor of bread wheat.

2. MATERIALS AND METHODS

2.1. Field Experiment

2.1.1. Description of the experimental sites

Field experiments were conducted in two different environments in the Bale and West Arsi Zones of Ethiopia during the 2010 main cropping season. The experiments were carried out at Sinana (located at 7°7' N, 40° 10' E) in Bale and Adaba (located at 7°01'N, 39°24'E) in the West Arsi Zones of Ethiopia. The two experimental sites have elevations of 2,400 and 2,300 m above sea level, respectively.

Sinana is characterized by a bi-modal rainfall pattern with two growing seasons. Annual rainfall, mean minimum and maximum temperatures of the site during the experimental year were 1,464 mm, 10.1°C and 20.4°C, respectively. On the other hand, Adaba is characterized by a mono-modal rainfall pattern, and received rainfall of 890.5 mm during the experimental year. The mean minimum and maximum temperatures at Adaba, during the cropping year were 8.15°C and 23.82°C, respectively.

Analysis of soil samples, before sowing, indicated that the soil texture of each experimental site was dominated by clay. Available Olsen P values were 13.3 ppm for Adaba and 15.4 ppm for Sinana. The available soil K content was 897.2 ppm at Adaba and 965.6 ppm at Sinana. Soil pH of the experimental sites was nearly neutral (6.9 at Adaba and 7.1 at Sinana). These values are in the pH range suitable for N availability. The soil of Adaba has an organic matter content of 2.2% whereas that of Sinana has 1.9%. The experimental soils had low levels of total N (0.11% and 0.14%, respectively, for Adaba and Sinana). According to Food and Agricultural Organization of the United Nations (FAO) [17], the P content of the soil is medium. The K content, at both sites, was very high [17, 18]. The organic C as and the total soil N content of both sites were

low [18]. The low organic matter and total N content of the soils signify that the release of NO_3^{-1} from mineralization of organic matter [19] would be low during the growing season of crops and application of external sources of N are required to maximize crop yields.

2.1.2. Experimental materials

Nitrogen fertilizer in the form of urea (46% N) and phosphate fertilizer in the form of triple super phosphate (TSP) (20% P) were used for as fertilizer material for the study. Two bread wheat varieties, Madda Walabu (HAR-1480) and Hollandi (local) with different morphological characteristics and yield potential were used as test crops. Madda Walabu is a CIMMYT origin semi-dwarf improved wheat variety, which was released by Sinana Agricultural Research Centre (SARC) while Hollandi is a local variety and is used for crossing in SARC breeding programs for its merits (adaptability and bread-making quality). The improved variety, Madda Walabu, was chosen because it is high yielding, well adapted to the area, and is widely grown by smallholder farmers as well as large scale commercial wheat producers in the region.

2.1.3. Treatments and experimental design

The treatments consisted of four levels of N (30, 60, 90 and 120 kg N ha⁻¹), two bread wheat varieties (Madda Walabu and Holandi) and three times of N application ($\frac{1}{2}$ at sowing and $\frac{1}{2}$ at mid-tillering; $\frac{1}{4}$ at sowing, $\frac{1}{2}$ at mid-tillering and $\frac{1}{4}$ at anthesis; $\frac{1}{3}$ at sowing, $\frac{1}{3}$ at mid-tillering and $\frac{1}{3}$ at anthesis). The experiment, at each site, was laid out in a randomized complete block design (RCBD) with a factorial arrangement of 4 x 2 x 3 = 24 treatment combinations and was replicated three times. The times of N application were adjusted according to Zadoks decimal growth stage for wheat at the time when moisture was available for nutrient dissolution and absorption.

2.2. Laboratory Experiments

2.2.1. Determination of seed germination capacity and vigor

After harvest standard germination tests for seeds obtained from control and treated plots were carried out between two layers of moist filter paper according to International Seed Testing Association [20] rules to evaluate treatment effects on seed germination capacity. The working samples, consisting of 400 pure seeds from each treatment, counted using an electronic seed counter and tested in a factorial CRD in

four replicates of 100 seeds. The seeds were placed at uniform spacing between layers of filter paper moistened with distilled water and rolled as a substratum. The prepared rolls were placed in a slanting position in plastic containers and incubated in a growth chamber at a temperature of 20°C and 80% relative humidity under illumination. Samples were monitored daily and supplied with deionized water when needed. Seedling vigor was determined based on seedling dry weight, which is often measured as an indicator of seedling vigor. Seedling dry weight was measured after the final count of the standard germination test. Ten normal seedlings were randomly sampled from each replicate and dried in an oven at 60°C to constant weight.

2.2.2. Determination of Seed Germination Speed

Seed germination speed, which also expresses seedling vigor and germination rate, was measured to determine treatment effects. For this purpose, another laboratory experiment was carried out using 100 seeds from each treatment. The seeds were sown on top of moist double filter paper arranged in a factorial CRD with four replicates of 25 seeds. The prepared samples were placed in Petri dishes and incubated at 20°C in a growth chamber in the dark. Subsequently, seeds were evaluated for the presence of normal seedlings from the fourth to the eighth day of germination. Seedlings were counted and removed daily.

2.3. Data Measurement

Germination capacity (%):- on the final day of the test period, germination components were categorized into normal and abnormal seedlings, fresh ungerminated and dead seeds according to international seed testing rules [20]. The respective percentages of each component were calculated based on the number of seeds used in the test.

Seedling dry weight (mg dry weight/germinal seedling):- this was measured as the average oven dry weight of ten randomly selected seedlings by weighing on an electric balance. *Seed germination speed*:- this was calculated using the following formula [21].

$$\text{Germination Speed} = \frac{\sum n}{\sum (n \times Dn)} \times 100$$

Where, n = number of normal seedlings germinated on day (D) and Dn = number of days from sowing corresponding to n.

Seedling vigor index:- was calculated by multiplying the standard germination percentage by mean seedling dry weight [22].

$$\text{Value Cost Ratio (VCR)} = \frac{\text{Value of increased yield obtained}}{\text{Cost of fertilizer used}}$$

2.4. Statistical Analysis

Analyses of variance, for the data recorded, were conducted using the SAS GLM procedure. Bartlett's homogeneity test of error variance was performed for all data of the two environments using Minitab software before performing a combined analysis. For seed germination components, variance analysis was done on the data transformed by the following function. $y = \arcsin \sqrt{p/100}$, where p is the percentage of the germination component. Significant differences among treatments were computed using Duncan's Multiple Range Tests.

3. RESULTS AND DISCUSSION

All measured and analyzed data were subjected to a homogeneity test of error variances using Bartlett's χ^2 test. Homogeneity of error variances was observed for the two test environments for all data. Hence, all through the text, the results of the combined analysis for the homogenous data are presented and discussed.

3.1. Seed Germination Capacity

The laboratory experiment showed normal germination of seed samples obtained from each treatment i.e. initially no seed germination was observed from the start of incubation until the 3rd day. Small numbers of seeds were germinated on the 4th day of incubation period. A rapid increase in the number of germinating seeds was observed on the 5th and 6th days. This was followed by a decrease in their appearance after the 7th day of incubation (Figure 1). Application of the highest N rate (120 kg N ha⁻¹) on the variety Madda Walabu resulted in a significantly higher percentage of total number of normal seedlings and proportion of germinating seeds in the early incubation period (4th to 5th days). However, for the local variety, the proportion of germinating seeds in the same period did not differ among treatments. For the local variety, application of 120 kg N ha⁻¹ resulted in the highest proportion of germinated seeds but on the 6th day of incubation (Figure 2). Generally, the percentage of seed that germinated during the

incubation period more or less followed a normal distribution curve that predicts the time to seedling emergence under different N application treatments.




Figure 1. Effect of N application rate on germination of normal seedlings of improved and local bread wheat varieties




Figure 2. Effect of N application rate on germination of normal seedlings per day of improved and local bread wheat varieties

Analysis of variance, for the data obtained at the end of the incubation period, revealed a significant influence of N rate and time of N application on the percentages of normal and abnormal seedlings, and dead seeds. The analysis showed there was no effect of time of N application on the proportion of fresh ungerminated seeds. Across all N rates and times of application. However, the varieties differed significantly ($P = .01$) in terms of the percentage of normal and abnormal seedlings. However, the varieties did not

interact with N rate and time of applications to influence all the germination components.

The highest percentage of normal seedlings (93.3%) was obtained in response to the application of 120 kg N ha⁻¹ and the lowest percentage (88.6%) was obtained at 30 kg N ha⁻¹. Conversely, the highest percentage of abnormal seedlings (5.8%) was observed at the lowest N application rate and the lowest (3.6%) was obtained at the highest N application rate. The percentage of dead and fresh ungerminated seed decreased significantly in response to increased rates of N application. Averaged over all N supply treatments, the variety Madda Walabu produced a higher percentage of normal (92.2%) and a lower percentage of abnormal (4.1%) seedlings. This indicates the superiority of the improved wheat variety over the local variety in terms of germination capacity.

The three times of N application treatments gave the highest percentages of normal seedlings while the two times of N application gave the lowest percentage. These results are consistent with other findings [8, 15] in which significantly higher percentage of normal seedlings reported in response to application of higher rates of N. The higher percentage of normal and lower percentage of abnormal seedlings in response to the application of the highest N rate in the three times split might reflect the significant role played by N in enhancing assimilate accumulation in the seeds, which may have promoted seedling growth in this study. These results evidently accentuate the importance of N for improved wheat seed germination, and hence improved crop productivity.

The varieties significantly differed in the percentages of fresh ungerminated seeds and dead seeds, in which variety Madda Walabu had the lowest proportions. The result indicates the importance of field application of higher N fertilizer rates on a wheat crop to minimize non-germinating seeds. Seeds produced in response to N application of ½ at sowing and ½ at mid-tillering (½P and ½MT) had a 28.6% higher percentage of dead seeds than seeds produced with three times of application. The superiority of the three times of N application in lowering the number of dead seeds may be attributed to the importance of applying N at the three critical growth stages for the synthesis and

accumulation of assimilate in seed during grain filling that promoted not only seed yield, as shown in the preceding sections, but also seed planting values.

3.2. Seed Germination Speed, Seedling Dry Weight and Vigor Index

Analysis of variance indicated a significant ($P = .01$) effect of field N application and timing on seed germination speed, seedling dry weight, and seedling vigor index. There were also significant variations between the varieties for the measured parameters. However, the varieties did not interact with N rate and time of application.

Seed germination calculated from the 4th to the 8th (final) days of the standard germination test revealed higher speed of seed from plants supplied with 120 kg N ha⁻¹. Field application of 30 kg N ha⁻¹, on the other hand, gave a significantly lower seed germination speed (Figure 3). Nitrogen application of $\frac{1}{4}$ at sowing, $\frac{1}{2}$ at mid-tillering and $\frac{1}{4}$ at anthesis ($\frac{1}{4}$ P, $\frac{1}{2}$ MT and $\frac{1}{4}$ A) resulted in a significantly higher speed of seed germination.

Nitrogen application at 120 kg N ha⁻¹ also resulted in higher germinal seedling dry weight (38.2 mg seedling⁻¹). Each increase in N application resulted in a significant increase in germinal seedling dry weight. Mean comparison of the varieties indicated a higher seedling dry weight of Madda Walabu compared to Hollandi, by 9.7%. Seeds obtained in response to the application of N in two equal splits at sowing and mid-tillering ($\frac{1}{2}$ P and $\frac{1}{2}$ MT) resulted in lower seedling dry weights compared with the three times of N application treatments (Figure 4).

Seedling vigor index was also significantly increased with increased rates of N application in which the highest (35.7) and the lowest (27.9) indices were obtained at 120 and 30 kg N ha⁻¹, respectively (Figure 3). The difference between the varieties in vigor index was 11.9% higher for the improved variety. The vigor index of seed produced by plants treated with $\frac{1}{2}$ P and $\frac{1}{2}$ MT was significantly lower than the vigor index of seeds from plants treated with three times split N applications (Figure 4).

The higher germination speed, seedling dry weight and vigor index of seed from plants supplied with higher N rates and three times of application may be ascribed to increased assimilate synthesis and translocations to the grain which may have resulted

in higher contents of protein and other food reserves. Corroborating these results, other authors [8, 15] also reported vigorous seedling growth for seeds obtained from wheat plants supplied with high N rates.




Figure 3. Seed germination speed, seedling dry weight and vigor index of bread wheat seeds obtained from crop supplied with different N rates




Figure 4. Seed germination speed, seedling dry weight, and vigor index of bread wheat seeds obtained from crop supplied with N at different growth stages

4. CONCLUSION

Low available soil N and low rates, as well as ill-timing of N application are major constraints limiting the sowing value of seed produced in major wheat growing areas of Ethiopia. The results of this study indicate that the capacity of seed germination and seedling growth was significantly enhanced in response to increased application of N in splits of three doses at three distinct growth stages of a wheat crop. The highest values for seed germination speed, seedling dry weight and vigor index were obtained at the application of 120 kg N ha⁻¹. In-season N applications of 1/4P, 1/2MT and 1/4A and 1/3P, 1/3MT and 1/3A also resulted in the highest seed germination capacity and seedling vigor. The improved wheat variety showed higher performance for these variables than the local variety. It could, therefore, be concluded that application of 120 kg N ha⁻¹ in three splits at sowing, mid-tillering and anthesis resulted in optimum seed germination capacity and seedling performance of the variety in the study areas. Therefore, farmers should cultivate the improved Madda Walabu variety by applying 120 kg N ha⁻¹ in three splits during the three distinct growth stages of the plant for optimizing the planting values of seed produced and productivity of the crop in the region.

ACKNOWLEDGEMENTS

We would like to thank the Local Seed Business (LSB) Project hosted by Haramaya University, which is sponsored by the Netherland Government through Wageningen University and the Royal Netherlands Embassy in Ethiopia, for funding the research. Sinana Agricultural Research Centre and Haramaya University are acknowledged for facilitating field and laboratory research activities. We appreciate the valuable assistance rendered by Firehiwot Getachew, Habtamu Tadesse, and Debele Tsegaye in field activities, laboratory work, and data measurement.

COMPETING INTERESTS

We, the authors, declare that there is no competing interest from other people or organizations which inappropriately influence this work.

REFERENCES

1. Hussain MI, Shah SH. Growth, yield and quality response of three wheat (*Triticum aestivum* L.) varieties to different levels of N, P and K. Int. J. of Agri. and Biol. 2002; 4(3): 362-364.

2. Central Statistical Agency (CSA). Agricultural sample survey 2009/10: Report on area production of crops. The FDRE statistical bulletin Vol. 391, Addis Ababa, Ethiopia; 2010.
3. Woldeab A, Mamo T, Bekele M, Ajema T. Soil fertility management studies on wheat in Ethiopia. In: Hailu G, Tanner DG, Mengistu H, editors. Wheat Research in Ethiopia: A Historical Perspective. Ethiopia: IAR/CIMMYT,; 1991.
4. Bekele H, Verkuil H, Mwangi W, Tanner D. Adoption of Improved Wheat Technologies in Adaba and Dodola Woredas of the Bale Highlands, Ethiopia. Proceedings of the Second National Maize Workshop. Ethiopia: IAR/CIMMYT; 2000.
5. Fageria NK. Plant tissue test for determination of optimum concentration and uptake of nitrogen at different growth stages in lowland rice. Comm. in Soil Sci. and Plant Anal. 2003; 34: 259-270.
6. Tanner DG, Amanuel G, Asefa T. Fertilizer effect on sustainability in the Wheat based smallholder-farming systems of southeastern Ethiopia. Field Crops Res. 1993; 33: 235-248.
7. Teklu E, Hailemariam T. Agronomic and economic efficiency of manure and urea fertilizers use on vertisols in Ethiopian highlands. J. Agri. Sci. 2009; 8(3): 352-360.
8. Warraich EA, Basra SM, Ahmad N, Ahmed R, Muhammad A. Effect of Nitrogen on Grain Quality and Vigor in Wheat (*Triticum aestivum* L.). Int. J. of Agri. and Biol. 2002; 4 (4): 517-520.
9. Sage RF, Percy RW. The nitrogen use efficiency of C₃ and C₄ plants. Plant Phy. 1987; 84: 954-958.
10. Al-Abdulsalam MA. Influence of nitrogen fertilization rates and residual effect of organic manure rates on the growth and yield of wheat. Arab Gulf J. of Sci. Res. 1997; 15: 647-60.
11. Ugalde TD. A physiological basis for genetic improvement to nitrogen harvest index in wheat. In: Randall PJ, editor. Genetic aspects of plant mineral nutrition. Dordrecht: Kluwer Academic Publishers; 1993.
12. Fageria NK, Prabhu AS. Response of lowland rice to nitrogen application and seed treatment with fungicide doses to blast control. Pesquisa Agropecuária Brasileira. 2003; 39:123-129.

- 493 13. Otteson BN, Mergoum M, Ransom JK, Schatz B. Tiller contribution to spring
494 wheat yield under varying seeding and nitrogen management. *Agron. J.* 2008;
495 100: 406-413.
- 496 14. Bulisani EA, Warner RL. Seed protein and nitrogen effect on seedling vigor in
497 wheat. *Agron. J.* 1980; 72: 657-662.
- 498 15. Seadh SE, El-Abady MI, El-Ghamry AM, Farouk S. Influence of micronutrients
499 foliar application and nitrogen fertilization on wheat yield and quality of grain and
500 seed. *J. of Biol. Sci.* 2010; 9: 851-858.
- 501 16. Bekele A. Biochemical aspects of wheat in human nutrition. Proceedings of ninth
502 Regional Wheat Workshop for Eastern Central and Southern Africa. Ethiopia:
503 IAR/CIMMYT; 1991.
- 504 17. Food and Agricultural Organization of the United Nations (FAO). Soil and plant
505 testing as a basis of fertilizer recommendations. *Soils Bulletin No. 38/2.* Rome,
506 Italy; 2008.
- 507 18. Landon JR. Booker tropical soil manual. A handbook for soil survey and
508 agricultural land evaluation in the tropics and subtropics. London: Booker Tate
509 Limited; 1991.
- 510 19. Roy RN, Finck A, Blair GJ, Tandon HLS. Plant nutrition for food security. A guide
511 for integrated nutrient management. Food and Agricultural Organization of United
512 Nations. *FAO Fertilizer and Plant Nutrition Bulletin No. 16.* Rome, Italy; 2006.
- 513 20. International Seed Testing Association (ISTA). International rules for seed testing.
514 *Seed Sci. and Techno.* 2002; 37: 54-59.
- 515 21. Kotowski F. Temperature relations to germination of vegetable seeds. *American*
516 *Society of Hort. Sci.* 1926; 23: 176-84.
- 517 22. Agrawal PK. Seed vigor. In: Srivastava, JP, Simarski, LT, editors. *Seed*
518 *Production Tech.* Syria: Aleppo, Simarski, ICARDA; 1986.
- 519