

Awareness of the parasitic weed *Alectra vogelii* (Benth) amongst extension officers in three districts in Malawi

V.H. Kabambe^{1*}, Y.L.B. Tembo¹, E. Kazira²

¹Lilongwe University of Agriculture and Natural Resources, PO Box 219, Lilongwe, Malawi

²Lilongwe Agricultural Development Division, PO Box 259, Lilongwe, Malawi

ABSTRACT

Aims: To determine awareness of *A. vogelii* amongst government extension officers in selected districts in Malawi.

Study design: Survey

Place and Duration of Study: Department of Crop and Soil Sciences Kasungu, Mchinji, Lilongwe and Balaka districts in Malawi

Methodology: A structured questionnaire was administered to 118 extension personnel in the study areas. The data were subjected to cross tabulation in SPSS to obtain frequencies of the various study parameters.

Results: Of the 118 officers 91 % were aware of *Striga asiatica* witchweed compared to 36 % that were aware of *A. vogelii*, while *Striga forbesii* and *S. gesnoiroides* were hardly known (<2 %). Predominant sources of information for *S. asiatica* were meetings (55 %), brochures (32 %), college (21 %), and radio (16 %). Field days, newspapers, books, internet and projects were seldom sources (< 6%). For *A. vogelii* the predominant source was meetings (24 %), followed by brochures (17%) and college (3%) while the rest were less than 2 %. The control measures known for both species were resistant variety, sanitation, rotation and manure application. More personnel were aware of these as control measures for *S. asiatica* than against *A. vogelii*. Generally supervisory staff were more aware of control measures for both species than frontline staff.

Conclusion: Results of this study have shown that little awareness exists on the parasitic weed *A. vogelii* compared to *S. asiatica*. The current extension methods involving meetings and brochures are operational at very low rates, while there is much less effort for dissemination with media and field days. Meetings are predominant means of information source, and that information flow between extension personnel in supervisory positions and frontline staff is minimum. Knowledge on control methods is poor, particularly for frontline staff.

Keywords: witchweeds, technology transfer

1. INTRODUCTION

1.1 Importance of legumes in Malawi

In Malawi grain legumes are an important component of the daily diet, farm income and cropping systems. They compliment maize (*Zea mays*), the staple cereal, in that they are

* Vernon H. Kabambe. Tel.: +265 999343465; fax: +265 (0)1277287
E-mail address: kabambev@yahoo.com.

rich in protein, and also contribute soil fertility improvement through biological nitrogen fixation. In the 2011/12 growing season legumes were grown on a total of 978,582 hectares, 66% of which were pulses including soybeans (*Glycine max*), common bean (*Phaseolus vulgaris*) and cowpeas (*Vigna unguiculata*), while 33% of the land had groundnuts (*Arachis hypogea*). On the other hand, maize was grown on a total of 1,497,829 hectares (MoAFS, 2012). Thus, maize/legume systems are predominant in Malawi and offer opportunity for this compliment. The trend of direction in Integrated Soil Fertility Management (ISFM) in sub-Saharan Africa is to scale out and intensify grain legume–cereal rotations and mixed cropping systems for soil fertility improvement and income generation and to improve the protein content in the diet through the high content of essential amino acids [2-11]. In 2005/06 season the Government of Malawi introduced the Farm Input Subsidy Program (FISP) which includes grain legume seeds for ISFM in addition to improved household food security and income [12].

1.2 Scope of witchweed problem in Malawi

Most grain legume crops, however, are prone to attack by parasitic weed species, including *Alectra vogelii* Benth and *Striga gesnorioides*, both of the family Orobanchaceae, which mainly attack groundnuts, cowpeas, soybeans and bambara nuts (*Vigna subterranea*) [13-16]. Serious infestation causes stunted crop growth, wilting and yield loss. In Tanzania yield losses of up to 50% have been reported [17] and in Kenya a total crop loss was reported [18]. Yield losses of 80 – 100% were reported on cowpeas in Botswana [19]. In soybeans, yield loss due to *A. vogelii* was estimated at 70-90 % [20]. For farmers to effectively adopt any effective control measure, extension workers require knowledge on the existence of the weed, its biology and its management. An integrated control approach utilizing several measures is considered as the best approach for control of witchweeds, as no single method may provide complete control [21-22]. In light infestations, uprooting before flowering and timely destruction through drying in a pit and burning is recommended. Hygiene during harvest to avoid weed seeds in grain reduces further spread of the weed. Other control measures include use of catch crops and trap crops. Catch crops are susceptible species which are ploughed in or harvested after parasite attachment but before emergence and seed production. On the other hand trap crops are plants that are not hosts to *Alectra* but do stimulate germination of the *Alectra* seed. These measures result in depletion of the seeds in the soil. Important trap crops for *A. vogelii* include sorghum (*Sorghum bicolor* L.), maize (*Zea mays* L.), pearl millet (*Pennisetum purpureum* L.), cotton (*Gossypium hirsutum* L.), sunflower (*Helianthus annuus* L.), dolichos beans (*Lablab purpureus* [L.] Sweet), guar beans and hyacinth bean [23]. Little has been done in establishment of *Alectra* resistant varieties in legumes in Malawi. However, in cowpeas a resistant variety, Mkanakaufiti (IT 99-494-6) has been released in Malawi and is recommended for growing in hot-spot areas [24]. Crop rotation using *Alectra* immune legumes such as pigeon peas, dolichos beans guar beans and hyacinth beans and non-susceptible crops can reduce *Alectra* seed bank in soil. In Malawi the parasitic weed has been reported in Lilongwe and Kasungu plains and parts of the southern region [25-26]. It has also been seen in Mzimba in northern Malawi by the authors of this report in the course of their studies. Interestingly, it was not reported by Banda and Morris (1986) in their book on 'common weeds of Malawi.

1.3 Justification and objectives

While a lot of research and outreach has been undertaken on *Striga asiatica*, a parasitic weed of cereals [28, 21, 13, 22; 29-30], relatively much less has been done on *A. vogelii*. For example, there was training on biology and control of parasitic weeds for all divisional crops officers in 1997 [31]. A cowpea project at Bunda College supported by McKnight Foundation grant no. 06-741 distributed posters on *A. vogelii* biology and control to various

extension offices in all ADDs in the country in 2009. A leaflet was further distributed carrying information on a newly released *A. vogelii* resistant cowpea variety, Mkanakaufiti (IT 99-494-6 ex IITA) to all District Agriculture Development Officers (DADOs) in March 2011. As efforts to popularize legumes are intensified, the incidence of *A. vogelii* appears to be on increase. In the course of conducting research activities supported by the McKnight Foundation project numbers 06-741 and 10-236, we have noted that many extension officers were unaware of *A. vogelii* and its control, even though the parasite was widely manifested in farmers' fields within their working areas. This study was therefore conducted to determine the awareness of the parasitic weed *A. vogelii* amongst extension officers in three districts in Malawi.

2. METHODOLOGY

2.1 Background to the survey area and respondents

Using a simple questionnaire, a survey was conducted from March 2012 to June 2012 in selected districts of Lilongwe, Mchinji, Kasungu and Balaka, in central and southern Malawi. The survey was conducted amongst extension personnel and farmers. The extension staff belonged to different categories in the organizational structure of extension service in Malawi. The structure starts from national level (Ministry of Agriculture)→Agricultural Development Division (ADD) → District Agriculture Development Office (DADO) → Extension planning Area (EPA) → Section. The frontline staff are in charge of a section within the EPA. The EPA is headed by agricultural extension development coordinator AEDC while the section is headed by an agricultural extension development officer, AEDO. There are subject matter specialists at the national, ADD, DADO levels (MoAFS, 2007). A total of 118 staff responded. There were 86 responding officers hailing from 23 EPA's in Lilongwe DADO, in Lilongwe ADD, 7 from Rivirivi EPA in Balaka district, Machinga ADD, and 25 from 9 EPAs in Kasungu and Mchinji districts (combined) in Kasungu ADD. Of these EPAs five were involved in activities connected to *A. vogelii* coordinated by the authors. In all cases the questionnaire was administered just before commencement of a meeting or field event.

2.2 The research tool

The main tool was a structured questionnaire which was designed to capture the following:- respondents' designation, whether they knew *A. vogelii* and other witchweeds, if they had received any information about control of witch weeds, how they received the information and the type of information they received. The questionnaire was however not designed to indicate the type of information received through a specific method of reception. The sample size was 118 across categories of staff. These categories included 66.9 % AEDOs, 19.5% AEDCs, 9.3% DADOs and other staff at district level and 4.2% ADD level staff.

2.3 Functional background to extension system to Malawi

The Agricultural Extension Development Officer (AEDO) is in charge of a section and interacts with farmers in their daily duties. They report to the Agricultural Extension Development Coordinator, AEDC, in their respective EPAs. At the EPA's there are fortnightly meetings during which all staff are appraised on recent developments in the EPA, including updates on new technology or simple skills and knowledge transfer sessions. Most of the staff and farmer training takes place at EPA level. Training is usually coordinated by subject matter specialists at the District, ADD or National levels. Most research and outreach projects operate at EPA levels and may disseminate information in direct or indirect ways to both staff and farmers. For this report, all staff other than AEDO have been

categorised as supervisors and the AEDO as frontline staff. Further to this government extension system, there is pluralistic agricultural extension system in Malawi [32-33].

2.4 Data analysis


The data were analysed by tabulation as percentages of frequencies using Statistics Package for Social Scientists (SPSS) 16th edition.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Awareness of various parasitic weed species.

The percentages of extension personnel aware of various witch weeds are presented in Figure 1. Of the 118 respondents, much more extension personnel (91 %) were aware of *S. asiatica* witchweed than *A. vogelii* (36 %). The other two species of *S. forbesii* and *gesnorioides* were seldom known (<2.0%).



3.1.2 Sources of information.

Table 1 shows the means of accessing information on *S. asiatica* or *A. vogelii* by the officers, broken down by category of staff (supervisors, frontline or all). For both species and all staff popular sources of information were brochures and meetings, while radio and college were important sources knowledge for supervisory staff with respect to *S. asiatica*. In general, more supervisory staff had access to knowledge than frontline staff. Even then, more information was on *S. asiatica* than *A. vogelii*. However, of the 39 frontline personnel, 27 % were aware of *S. asiatica* from college days, compared to 10% for supervisors. Notably, of the 79 supervisory staff 80 % were aware of *S. asiatica* through meetings, compared to 44 %. In terms of source of information for *A. vogelii* for frontline staff, the most important source of information was brochures followed by meetings and college. However knowledge were low (5-15 %). In comparison, the predominant source of information on *A. vogelii* for supervisory staff was brochure, followed by meetings.

167 Table 1. Percent of supervisory, frontline and all staff positive response to receiving
 168 information on control of *A. vogelii* and *S. asiatica* from various sources.
 169

Information source	<i>A. vogelii</i>			<i>S. asiatica</i>		
	All (n=118)	Supervisory staff (n=39)	Front line personnel (n=79)	All (n=118)	Supervisory staff (n=39)	Front line personnel (n=79)
Brochure	23.7	41.0	15.2	32.2	43.6	27.8
Meetings	16.9	24.1	13.9	55.1	79.9	44.3
News paper	0.8	2.6	0	5.1	7.7	3.8
Radio	1.7	0	2.5	16.1	23.1	12.7
College	3.4	0	5.1	21.2	10.3	26.6
Field days	1.7	0	2.5	5.9	5.1	6.3
Projects	1.7	2.6	1.3	0	3.6	0
Internet	0.8	2.6	0	0.8	2.6	0
Books	0.8	2.6	0	0.8	2.6	0

170 3.1.3 Control measures known

171 The control measures for *A. vogelii* most known by all staff were crop rotation (25 %),
 172 followed by resistant variety, manures and sanitation (Table 2). However, the levels were
 173 quite low (7-25 %). Early planting, use of trap crops, were seldom known (< 1 %). General
 174 awareness of control measures for *A. asiatica* was much higher, with crop rotation as most
 175 popular (76 %), followed by manure use (59 %), sanitation (30%) and resistant variety (13
 176 %). The others were seldom known (<2.5%). Among the supervisory staff, the commonly
 177 known methods for control of *A. vogelii* was crop rotation (36%), followed by manure use (23
 178 %), resistant variety and sanitation (8%). For frontline staff, crop rotations remained the most
 179 known method, followed by sanitation and manure, but the levels were quite low (5-19%).
 180

181 3.2 DISCUSSION

182 3.2.1 General awareness of *A. Vogelii* and *S.asiatica* and their control measures

183 The results of this study have shown that extension personnel at all levels are more aware of
 184 *S. asiatica* as a parasitic weed, including associated control measures compared to *A.*
 185 *vogelii*. This concurs with current literature which shows much research and outreach on *S.*
 186 *asiatica* compared to *A. vogelii*. The awareness of *A. vogelii* (36 %) is considerably much
 187 lower, considering the greater efforts being taken to promote legumes in Malawi [12, 2].
 188 Regarding type of information, the results indicate that officers were less knowledgeable on
 189 *A. vogelii*, compared to *S. asiatica*, an observation in line with the status quo in Malawi.
 190 There have been less circulars, seminars and conference presentation on this legume
 191 witchweed.
 192
 193
 194

Table 2. Percent of supervisory, frontline and all staff with positive response to receiving information on control of *A. vogelii* and *S. asiatica* from various sources.

Control method	<i>A. vogelii</i>			<i>S. asiatica</i>		
	All (n=118)	Supervisory staff (n=39)	Front line personnel (n=79)	All (n=118)	Supervisory staff (n=39)	Front line personnel (n=79)
Resistant variety	12.7	12.8	12.6	11.9	7.7	13.9
Sanitation	6.8	7.7	6.3	29.7	38.5	25.3
Crop rotation	24.6	35.9	19	76.3	79.5	74.7
Manure application	7.2	23.1	5.1	58.5	79.5	48.1
Early planting	0.8	0	2.6	2.3	5.1	1.3
Trap crops	0	0	0	0.8	0	1.3
Striga powder	0.8	0	1.3	1.7	0	2.5
Intercropping with <i>Tephrosia vogelii</i>	0	0	0	1.7	5.1	0

Many conferences and review papers which tackled *S. asiatica* did not tackle *A. vogelii* (e.g. [34-36]. We encourage that promotion of legumes should be simultaneous to awareness of important pests and diseases. We recommend that a target of 100 % knowledge is set for methods such as sanitation, manure application and rotations as these are preventive, least-cost and may also result in overall farm productivity [37, 29, 13]. However, it must be acknowledged that research on control of *A. vogelii* is low, and locally verified control measures are few and that that much of the knowledge on control is packaged from regional and international literature. It is recommended that research in Malawi should be intensified to verify, adapt and package most of these rotation or manure options. In Kenya, recent research has shown that use of cattle manure at 5 and 10 t ha⁻¹ reduced *A. vogelii* incidence and increased yield and yield components of some cowpea varieties, and not others [38].

About 12 % of all staff indicated awareness of resistant variety as control measure for *S. asiatica*, and yet there is no resistant variety available. It is likely that information that the combined use of hybrids, fertilizer and manures can result in improved yields under *S. asiatica* infestation [37, 21, 35] may be misunderstood to mean there are resistant varieties. Only about 13 % of all staff were aware of resistant variety as control measure for *A. vogelii*. The only legume variety with resistance to *A. vogelii* was released in 2011). Brochures on this were sent to all district agriculture offices, and it would have been expected that at least through meetings and brochures, higher levels of awareness would be recorded.

3.2.2 Sources of information.

The results on sources of information concur with a previous study with *Striga* spp on cereals by [28] who reported that extension bulletins were the most popular source at 35-66%, while other means such as posters (5-18 %), NGO's (5-13 %) or farmers and friends

(12 %) were low. Meetings, field days, colleges, radios were not mentioned. The finding in of this study that meetings were a predominant source of information is important is that staff meetings are routinely conducted and provide an opportunity for cost-effective technology transfer. However, the percent of positive responses to meetings is very low, particularly for *A. vogelii* information, we suggest a target of 100% as ideal. The low rating of field days suggests that there are hardly attempts to mention parasitic weeds despite their widespread nature, particularly in Mchinji and Lilongwe districts, where this study was conducted. We recommend that stakeholders should incorporate themes on control of these parasitic weeds in their legume promotional packages as these are getting widespread and cause considerable yield loss.

3.2 3 Access to information by frontline versus supervisory staff

The supervisory staff were more knowledgeable on control measures for both *A. vogelii* and *S. asiatica*. This is expected as supervisory staff are meant to be sources of information. However, staff are not directly passing information to frontline staff, noting that only 13.9 and 44.3 % of frontline staff obtained information on *A. vogelii* or *S. asiatica* from meetings. Interestingly, a high percent of supervisory staff obtained information from meetings, suggesting technology transfer amongst supervisors does take place, but is not passed on to frontline staff. The current extension policy in Malawi provides for pluralistic extension services, allowing for more players at frontline levels [32]. With this, it is surprising that field days and projects have not been dominant sources of information. A recent report indicated that there were up to 9 and 11 agricultural extension service providers in two villages in Mchinji district [33]. Such results suggest that while extension providers exist, there is less priority on these devastating parasitic weeds.

4. CONCLUSION

Results of this study have shown that very little awareness exist on the parasitic weed *A. vogelii* compared to *S. asiatica*. Meetings are predominant means of information source, and that information flow between extension personnel in supervisory positions and frontline staff is minimum. Knowledge on control methods is poor, particularly for frontline staff. With the standing pluralistic extension policy, we remind stakeholders to incorporate dissemination of parasitic weeds in their routine meetings, field days and media dissemination efforts. We recommend that a target of 100 % knowledge is set for methods such as sanitation, manure application and rotations as these are preventive, least cost and may also result in overall farm productivity.

ACKNOWLEDGEMENTS

We are grateful to the McKnight Foundation of USA for funds provided to Grant 06-741 and 09-236, which supported this work. We are also grateful to staff of Lilongwe ADD, Kasungu ADD and Balaka District for participating in this study. We acknowledge Dr Beston Maonga for proof reading the first manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

Author VHK designed the study, managed the literature searches, discussed the results and wrote manuscript. Author YLBT collected the data, performed the statistical analysis, wrote

the protocol and the first draft of the manuscript. 'Author EK collected data. All authors read and approved the final manuscript.

REFERENCES

1. Ministry of Agriculture and Food Security, MoAFS. Agricultural Production Estimates, 2007/08 season. Planning Department. MoAFS. Lilongwe. Ministry of Agriculture and Food Security. 2012.
2. Ngwira A R, Kabambe V H, Kambauwa G, Mhango WG, Mwale C D, Chimphero L, Chimbizi A, Mapfumo P. Scaling out best fit legume technologies for soil fertility enhancement among smallholder farmers in Malawi. *African Journal of Agricultural Research*, 2012; 7 (6): 918-928. DOI:10.5897/AJAR11.760.
3. Gwata ET. Potential impact of edible tropical legumes on crop productivity in the smallholder sector in Sub-Saharan Africa. *Journal of Food, Agriculture and Environment*, 2010; 8:939-944.
4. Nyemba RC, Dakora FD. Evaluating nitrogen fixation by food grain legumes in farmers' fields in the three agro-ecological zones of Zambia using the ^{15}N natural abundance. *Biol. Fertil. Soils*. 2010; 46: 461-470.
5. Misiko M, Tittonell P, Ramisch JJ, Richards P, Giller KE. Integrating new soybean varieties for soil fertility management in smallholder systems through participatory research: Lessons from western Kenya. *Agricultural Systems*, 2008; 97: 1-2.
6. Snapp, SS, Kanyama-PhiriGY, Kamanga B, Gilbert R, Wellard K. Farmer and researcher partnerships in Malawi: Developing technologies for the near-term and far-term. *Exp. Agric.* 2002; 38: 411-431.
7. Mwato IL, Mkandawire ABC, Mughogho SK. Combined inputs of crop residues and fertilizer for smallholder maize production in Southern Malawi. *Afr. Crop Sci. J*, 1999; 7:365-373.
8. Kumwenda JDT, Waddington SR, Snapp SS, Jones RB, Blackie MJ. Soil Fertility Management in Southern Africa. In: Byerlee D, Eicher CK (eds). *Africa's Emerging Maize Revolution*. Lynne Reiner Publishers, Colorado, 1997.
9. Kumwenda JDT, Saka AR, Snapp SS, Ganunga RP, Benson TD. Effects of organic residues and inorganic fertilizer nitrogen on maize yield. In: Benson TD, Kumwenda JDT (eds). *Maize Commodity Team Annual Report for 1996/97 season*. Chitedze Research Station Lilongwe, Malawi, 1988; Pp 77-91.
10. Baptist, NG. Essential amino acids of some common tropical legumes and cereals. *British Journal of Nutrition*, 1954; 8: 218-222.
11. Mahe S, Gausseres N, Tome D. Legume protein for human requirements. *Grain Leg.*, 1994; 7: 15-17.
12. Ministry of Agriculture, Irrigation and Food Security, MoAFS. (Annual Agricultural Statistical Bulletin. Planning Division. Government of Malawi. Lilongwe, Malawi, 2007.
13. Kabambe VH, Katunga L, Kapewa T, Ngwira AR. Screening legumes for integrated management of witchweeds (*Alectra vogelii* and *Striga asiatica*) in Malawi. *African Journal of Agricultural Research*, 2008; 3 (10): 706-713.
14. Riches CR. The identification of resistance to *Alectra vogelii* Benth (Scrophulariaceae) in cowpea. In: Weber, Chr. H and Forstreuter, W. (eds). *Parasitic Flowering Plants. Proceedings on the 4th International Symposium on Parasitic Seed Plants*, Marburg, 1987; pp 701 – 708.
15. Lagoke STO, Parkinson V, Agunbiade RM. Parasitic weeds and control in Africa. In: Kim SK (ed). *Combating Striga in Africa. Proceedings International Workshop organised by IITA, ICRISAT and IDRC*, IITA, Ibadan, Nigeria, 1988; pp. 3– 17.
16. Omoigui1 LO, Kamara AY, Ishiyaku MF, Boukar O. Comparative responses of cowpea breeding lines to *Striga* and *Alectra* in the dry savanna of northeast Nigeria. *African Journal of Agricultural Research*, 2012; Vol. 7(5), pp. 747-754.

17. Mbwaga AM, Kaswende J, Shayo. A reference manual on *Striga* distribution and control in Tanzania. Kilosa, Tanzania: Ilonga Agricultural Research Institute, 2000.
18. Bagnall-Oakley H, Gibberd V, Nyongesa TE. The incidence and control of *A. vogelii*, in Embu district, Kenya. In: Ransom JK, Musselman LJ, Worsham AD, Parker C (Eds.), Proceedings of the 5th International Symposium on Parasitic Weeds, CYMMIT, Nairobi, Kenya, 1991.
19. Riches CR. The biology and control of *Alectra vogelii* Benth. (Scrophulariaceae) in Botswana. PhD thesis, Reading, UK: University of Reading, 1989.
20. Kureh I, Katung PD, Orakwue FC. Reaction of soybean varieties to preconditioning and concentration of seed inoculum of *Alectra vogelii* (Benth). Science Forum. J. Pure Appl. Sci., 1999; 2(1): 116-124.
21. Kabambe VH, Mloza-Banda HR, Nyandule Phiri GY. Controlling witchweeds in cereals in Malawi. An extension bulletin for field staff. Department of Agricultural Research and Technical Services, Extension Bulletin no. 1/2002. Ministry of Agriculture and Irrigation, Lilongwe, Malawi, 2002.
22. Kabambe VH, Nambuzi SC, Kauwa AE. Integrated management of witchweed (*Striga asiatica* {L} Kunze) by means of maize-legume rotations and intercropping systems in Malawi. Bunda Journal of Agriculture, Environmental Science and Technology, 2008; 3 (2): 35 – 42.
23. Parker C and Riches CR. Parasitic weeds of the world. Biology and Control. CAB International. Oxon, UK. 1993; Pp332.
24. Mviha PJZ, Mtukuso A P, Banda MHP. A catalogue of agricultural technologies used by farmers in Malawi. Department of Agricultural Research Services, Lilongwe, Malawi, 2011; 40 pp.
25. Riches CR, Shaxson LJ. Parasitic weed problems in southern Malawi and use of farmer knowledge in the design of control measures. In: Munthali DC, Kumwenda JDT, Kisyombe F (eds.). Conference on Agricultural Research for Development organised by the Soil Pests Project (University of Malawi) and the Malawi Maize Research Team, Ministry of Agriculture. Mangochi, Malawi, 7-11 June 1993. Munthali, D.C., J.D.T. Kumwenda and F. Kisyombe (eds.). 1993; pp. 165-174.
26. Mainjeni CE. The host range of *Alectra vogelii* Benth. from Malawi and resistance in common bean and cowpea. Msc Thesis, University of Bath, UK. 1999; p. 83.
27. Banda EAK, Morris B. Common weeds of Malawi. University of Malawi; 1986
28. Mloza_Banda HR, Kabambe VH, Mphepo M, Chivinge O. Knowledge and perceptions of extension officers on *Striga* spp in Lilongwe Machinga Agricultural Development Divisions in Malawi. UNISWA Res. J. Agric. Sci. & Tech., 2000; 10:91-97.
29. Kabambe VH, Drennan DSH. Control of *Striga asiatica* in maize by means of crop rotation and intercropping in Malawi. In: Mloza-Banda HR, Salanje GF (eds.). Proc. of the 19th Biennial Weed Science Society Conference for Eastern Africa. Lilongwe, WSSEA, 2003; pp 105-111.
30. Sikwese M, Mloza-Banda HR, Kabambe VH, Phombeya HK. The potential of multipurpose tree species in the integrated management of *Striga asiatica* in Maize fields in Malawi. African Crop Science Conference Proceedings, 2003; 6:133-137.
31. Supra-Regional Project. Biology and control of *Striga* and *Alectra* in Malawi. Manual for the training of extension staff. GTZ GmbH and University of Hohenheim 1997; pp 380.
32. GOM (Government of Malawi). Agricultural Extension in the New Millennium: Towards Pluralistic and Demand Driven Services. Ministry of Agriculture, Lilongwe, Malawi, 2000.
33. Chowa C, Garforth C, Cardey S. Farmer Experience of Pluralistic Agricultural Extension, Malawi. The Journal of Agricultural Education and Extension, 2012. DOI:10.1080/1389224X.2012.735620
34. Kabambe VH. Integrated management of witchweed (*Striga asiatica*) in Malawi. Malawi Journal of Agricultural Sciences, 2002; 1: 47-51.

- 382 35. Mloza-Banda HR, VH Kabambe. Integrated management for *Striga* control in Malawi.
383 African Crop Science Journal, 1996; 1: 263-273.
- 384 36. Mloza-Banda HR, Salanje GF (eds). Proceedings of the 19th Biennial Weed Science
385 Society Conference for Eastern Africa. Lilongwe, WSSEA, 2003.
- 386 37. Kabambe VH. The development of cultural methods for control of *Striga* in maize in
387 Malawi. In: Rnasom JK, Musselman LJ, Worsham AD, Parcker C (Proc. 5th
388 International Symposium of Parasitic Weeds. Nairobi: CYMMYT, 1991 Pp 46-60.
- 389 38. Karanja J, Nguluu S, Gatheru M Farm yard manure reduces the virulence of *Alectra*
390 *vogelii* (Benth) on cowpea (*Vigna unguiculata*). African Journal of Plant Science, 2012;
391 6(3): 130-136