

HEAVY METAL CONTENT IN ACACIA SALIGNA AND ACACIA POLYACANTHA ON SLIME DAMS: IMPLICATIONS FOR PHYTOREMEDIATION

Masvodza D.R.^{1*}, Dzomba P.², Mhandu F.¹, and Masamha B.³

¹Bindura University of Science Education, Biological Sciences Dept, P. Bag 1020, Bindura

²Bindura University of Science Education, Chemistry Dept, P. Bag 1020, Bindura

³Bindura University of Science Education, Environmental Science Dept, P. Bag 1020, Bindura, Zimbabwe

ABSTRACT

Aims: To determine heavy metal content in *A. Saligna* and *A. Polyacantha* so as to ascertain their possible use in phytoremediation

Place and Duration of Study: Bindura University of Science Education, Chemistry and Biological Sciences Dept, P. Bag 1020, Bindura between December 2011 and January 2013.

Methodology: A total of four sampling points each in the form of 5x5m quadrants were established after every 100m in transects, 700m long separately established on both sites. Soil samples at 5-10cm and 10-15cm levels as well as roots, leaves and bark from five sampled plants were collected at each sampling point. The metals content was analyzed using Atomic Absorption Spectrometry and Inductively Coupled Plasma. The bioaccumulation factor and the shoot/root quotient were computed in Microsoft excel. Analysis of Variance was carried out using SPSS and Genstat Version 16.

Results: The present study shows that *A. Saligna* and *A. Polyacantha* accumulate heavy metals with BAF values results indicating significant differences between the slimes and control sites. Both species had BAF values for nickel, copper and iron greater than one except for zinc, lead and arsenic. The shoot/root quotients showed that nickel, copper and iron are translocated to the shoots in the species as compared to zinc, lead and arsenic.

Conclusion: *A. Saligna* and *A. Polyacantha* showed evidence of accumulation of nickel, copper and iron and therefore may be used for phytoremediation and restoration purposes at mine slime dams.

Keywords: [heavy metals, Phytoremediation, bioaccumulation factor, translocation factor, *Acacia polyacantha*, *Acacia Salina*]

1. INTRODUCTION

Contamination of soil, water and food plants with toxic heavy metals due to mining activities in mining towns is still a major environmental and human health problem (1). While methods such as excavation and burial of contaminated soil at designated waste sites have been suggested such methods are not popular due to huge costs. They cost over a million United States dollars per acre (2). There is still a need for researches in effective and affordable methods of counteracting this challenge (3; 4). Phytoremediation offers attractive options. It takes advantage of the fact that living plants can act as solar driven pumps that can extract and concentrate particular elements

from the environment (5). Harvested plant tissue that would have accumulated heavy metal contaminants may be easily and safely processed by drying, ashing or composting. Metals can then be reclaimed from the ashes. This generates recycling avenues and reduces the generation of hazardous waste (6). Major sources of heavy metal pollution in the environment are mostly anthropogenic, including mining activities, effluent discharges and waste disposal (7). In trace concentrations, many metals are essential to life and have several vital functions in biological processes but in excess the same metals can be toxic. It has been observed that even long after mining activities have ceased heavy metals continue to persist in the environment (8). They can enter the food chain when taken up by plants during farming and eventually affect human health (9). Heavy metals poison animals and humans by disrupting cellular enzymes, which use nutritional minerals such as magnesium, zinc and selenium for their function. Toxic metals replace these nutrients and bind their receptor sites, causing diffuse symptoms by affecting nerves, hormones, digestion and immune function (10).

A survey of most mine slime dumps in Bindura town shows that *Acacia polyacantha* and *Acacia saligna* grows very well in these areas. The plants depicted no stress or any stunted growth. Thus the proliferation of *Acacia polyacantha* and *Acacia saligna* on slime dams makes them an important object of research. While *A. polyacantha* is an indigenous species in Zimbabwe *A. saligna* is exotic and native to Australia and falls within a family of acacias commonly known as Australian wattles. The species spread to many parts of the world and is now considered the most widely planted non-timber species with around 600,000 ha established worldwide (11). Both species have been mentioned in other studies as ideal for phytoremediation (12; 13). *A. saligna* was reported to have a high biomass and fast growth rate (11). *A. polyacantha* out-performed indigenous species like *Bauhinia thonningii* in biomass production and growth rates (13, 14). Coates (11) indicated in his studies that the fast growth rate of *A. saligna* makes it ideal for reforestation of mine dumps (5). There is still need for more work on the capacities of both species as accumulators as well as the physiology involved (15). Therefore this research sought to assess heavy metal content of *Acacia polyacantha* and *Acacia saligna* with an aim of finding out if they can probably be used in heavy metal decontamination of the slime mine dumps.

2. MATERIALS AND METHODS

2.1 Study Area




Fig 1. A map of Bindura town where the study area is located.

Bindura town is about 88 km North-East of Harare. The area is generally rich in gold, nickel, copper and cobalt.

2.2 SAMPLING PROTOCOL

Sampling was carried out at the slime dams alongside a control site 50km outside, with a presumption that it was well separated from the mine. A line transect of 700 m long was established on the sites and a total of 4 sampling sites were systematically established after every 100 m in line transects. Four 5x5 m quadrants were established on the sites one at each established sampling point in the line transects.

2.3 Plant sample collection

Leaf, bark and root samples were collected from plants of *Acacia polyacantha* and *Acacia saligna* species at the slime dams and the control site. Five individual plants were sampled in each quadrant using simple random sampling. The plants of heights between 40 and 50cm were selected as this was an estimate for similar ages. The samples were separately collected from each plant and were wrapped in aluminum foil and labelled. All the samples were kept in a cooler box with ice during transportation and brought to the laboratory according to (16). Heavy metal contents were determined in the plant parts.

2.4 Soil collection

Three soil samples were also collected at each sampling point where plant samples were previously taken. The samples were then mixed to constitute composite samples per sampling point. In each plot, soil samples were collected at two depths, 5-10 and 10-15 cm using a soil auger (2). In all cases, soil samples were placed in clean plastic bags, sealed and transported to the laboratory. Soil samples were air-dried, ground into fine powder using pestle and mortar and passed through 2mm mesh sieve (2). Soil samples were analyzed for pH and heavy metal content.

2.5 SAMPLE ANALYSIS

2.5.1 Quality assessment and control

This was achieved by analyzing results in triplicates and use of certified reference standards to ascertain reproducibility and accuracy as shown in Table 1. Distilled water was used throughout the study. Glassware was properly cleaned and the reagents used were of analytical grade. Reagent blank determinations were used to correct the instrument readings.

Table 1. Plant certified reference material concentration (ppm) mean±SE.

Metal	Certified value	Measured value	Recovery
Copper	120±0.4	120±0.5	100
Zinc	260±0.5	260±0.5	100
Nickel	1.2±0.01	1.1±0.05	92
Lead	73±0.5	73±0.5	100
Arsenic	1.6±0.5	1.5±0.01	94

2.5.2 Soil pH

The soil pH was measured in a suspension at a soil-to-water ratio of 1:2 using a pH meter.

2.5.3 Soil heavy metal analysis

Two grams of the soil samples were weighed into nitric acid-washed glass beakers. Soil samples were digested by the addition of 20 cm³ of *aqua regia* (mixture of HCl and HNO₃, ratio 3:1) and 10 cm³ of 30 % H₂O₂ (2). The H₂O₂ was added in small portions to avoid possible overflow leading to loss of material from the beaker. The beakers were covered with a watch glass, and heated at 90°C for two hours. The beaker wall and watch glass were washed with distilled water and the

143 samples filtered out to separate the insoluble solid from the supernatant liquid and the filtrate was
144 collected. The volumes were adjusted to 100 cm³ with distilled water. All the samples and blanks
145 were stored in plastic containers.

147 **2.5.4 Plant heavy metal content analysis**

148
149 Root and shoot samples were thoroughly washed with distilled water to remove all adhering soil
150 particles. The plant samples were weighed to determine the fresh weight and dried in an oven at
151 80°C for 72 hours to determine their dry weight (2). The dry samples were crushed in a mortar and
152 the resulting powder digested by weighing 0.5 g of oven-dried ground and sieved soil (<1 mm) into
153 a nitric acid-washed porcelain crucible and placed in a muffle furnace for four hours at 500°C.

154
155 The crucibles were removed from the furnace and cooled. Ten mL of 6M HCl was added, covered
156 and heated on a steam bath for 15 minutes. Another 1 mL of HNO₃ was added and evaporated to
157 dryness by continuous heating for one hour to dehydrate silica and completely digest organic
158 compounds. Finally, 5 mL of 6 M HCl and 10 mL of water were added and the mixture heated on a
159 steam bath to complete dissolution. The mixture was cooled and filtered through a Whatman no.1
160 filter paper into a 50 mL volumetric flask and made up to the mark with distilled water.

161
162 Determination of nickel, copper, zinc, iron, arsenic, and lead in soil and plant samples was made
163 in triplicates directly on each of the final solution using Atomic Absorption Spectroscopy (AAS).
164 Determination of arsenic was also made in triplicates directly on the final solution using Inductively
165 coupled plasma (ICP).

167 **2.5.5 The bioaccumulation factor (BAF)**

168
169 The index of the plants' ability to accumulate metals from soils was calculated according to (11).

$$170 \quad \text{BAF} = \frac{C_{\text{plant}}}{C_{\text{soil}}}$$

172 where C_{plant} and C_{soil} represent the heavy metal concentration in plant parts and soils
173 respectively.

175 **2.5.6 The shoot/root quotient (SRQ)**

176
177 This may also be referred to as translocation factor (Ti) and it gives the leaf/root metal
178 concentration and it depicts the ability of the plant to translocate the metal species from roots to
179 leaves at different concentrations. It was computed as follows:

$$181 \quad \text{SRQ} = \frac{S}{R}$$

182 where S and R represent the level of heavy metals in the plant parts (leaves and bark) and R the
183 level of heavy metals in the roots respectively (13).

185 **3. RESULTS AND DISCUSSION**

187 **3.1 Soil pH**

188
189 The pH values for the soils on the mine dump samples ranged from 2.0 to 4.7 and can be
190 classified as acidic. The pH values at the control site ranged between 4.7 and 6.8 and can be
191 classified as acidic to neutral. Fig 2 shows the mean pH for the two sites.




Fig 2. A graph for the mean soil pH at the sites

3.2 Soil heavy metal contents

The sampling points within sites had non-significant variation ($P > 0.05$) for all the heavy metals, an indication of homogeneity among the sampling points. This means that the sampling strategy was effective and there was no bias in the strategy as all points within sites were uniform as shown in Table 2.

Table 2. Mean heavy metal concentration (ppm) for the soils at the control and slime dams.

		Ni	As	Cu	Fe	Pb	Zn
SITE	SLIMES	0.97ns	0.307ns	0.013***	0.18*	0.234***	2.45***
	CONTROL	0.51ns	0.057ns	0.201***	0.96*	0.043***	0.184***
	S.E	0.192	0.0949	0.0171	0.205	0.0169	0.0904
LEVEL	5-10cm	1.21**	0.0197*	0.146**	0.14*	0.121ns	1.454**
	10-15cm	0.28**	0.0616*	0.069**	0.99*	0.155ns	1.155**
	S.E	0.192	0.00949	0.0171ns	0.205ns	0.0169	0.0904
SAMPLING PT	1	0.76ns	0.00949ns	0.103ns	0.55ns	0.155ns	1.17ns
	2	0.72ns	0.0398ns	0.0103ns	0.55ns	0.123ns	1.43ns
	3	0.74ns	0.0405ns	0.109ns	0.56ns	0.137ns	1.24ns
	4	0.75ns	0.0424ns	0.115ns	0.6ns	0.138ns	1.377ns
	S.E	0.271	0.0401	0.02041	0.29	0.0239	0.1279

S.E.= standard error . ns= no significant difference ($p > 0.05$). * $p < 0.05$ ** $P < 0.01$ *** $p < 0.001$ ppm = parts per million 5-10cm – The top soil level at a depth of five to ten centimeters, 10-15cm bottom soil level at a depth of ten to fifteen centimeters, sampling pt: any one of four sampling points established at the two sites in the form of 5x5 quadrabts

Significant differences were observed between the sites for copper, zinc and lead ($P < 0.001$) and iron ($P < 0.05$) and non-significant differences for nickel and arsenic in the soil samples $P > 0.05$. Table 2. shows the means and standard errors. The analysis shows an indication that there was heavy metal accumulation at the slimes for four of the six metals considered. The non-significant differences for Nickel and Arsenic might be an indication that the control site was not ideal for these metals as it was only 10km from the mine dumps as substantive quantities of the metals were at both sites. Trojan Nickel Mine being a Nickel Mine would result in an abundance of the metal in the Bindura locality; hence a more far removed control site would have been better.

The soil depths, 5-10 and 10-15 cm are significantly different ($P < 0.01$) for nickel, copper, iron, zinc and arsenic and non-significantly different for lead. This shows a sign of movement of the metals between the levels as well as availability of the heavy metals in the soil layers. Fig 3 shows the heavy metal concentrations in the two levels.




Fig 3. Graph of heavy metal distribution in the two soil levels at the slime dams.
 5-10cm – The top soil level at a depth of five to ten centimeters, 10-15cm bottom soil level at a depth of ten to fifteen centimeters

It was observed that nickel is the most abundant heavy metal in the slimes at the 5-10 cm level, a confirmation that deposits of the metal were made at the slimes and on the other hand, copper, zinc and iron are most abundant at the control sites at different levels as shown in fig 4.




Fig 4. Graph of heavy metal distribution in the two soil levels at the control site
 5-10cm – The top soil level at a depth of five to ten centimeters, 10-15cm bottom soil level at a depth of ten to fifteen centimeters

The control site had a much less tree species density than the slimes which underwent reforestation for land reclamation by the mine. This may have resulted in the lower levels of some of the heavy metals observed at the slime dams. Iron is abundant at the control site in both levels.

The low pH at the slimes may have contributed to accumulation of heavy metals at the slimes (4). Soil acidity dramatically affects the cation exchange capacity (CEC) of soil by limiting the available exchange sites. H^+ bind to soil particles tighter than other cations, thus, any metal bound to a soil particle will get knocked off in the presence of excess H^+ (3). At high pH (>7), cations are less bioavailable because they have less competition from H^+ for available binding sites. Many cations bind to free hydroxyl groups (OH^-) and form insoluble hydrous metal oxides, which are unavailable for uptake, such as $CdCO_3$. This could explain why there was an accumulation of some of the heavy metals at the slimes.

3.3 Plant heavy metal contents

Generally the plants at the dams contain more metals than the control a reflection of the nature of the different sites. The roots at the slimes contain the most heavy metals especially nickel most likely as they are the first point of contact compared to the bark and leaves, whereas the leaves at the control contain the most iron a reflection of abundance of this metal at the site. Fig 5 summarizes mean heavy metal contents for different plant parts for the two species.




Fig 5. Heavy metal contents in the plant parts.

Bark – heavy metal contents in the bark, leaves – heavy metal contents in the leaves, root – heavy metal contents in the roots.

The contents become meaningful when expressed relative to soil contents, as the bioaccumulation factor.

3.4 Biological accumulation factor (BAF)

When BAF values were computed, the mean values shown in Table 3 were observed. Values greater than 1 indicate a net accumulation by the plant whereas values below 1 show net accumulation in the soil. Generally therefore the species are hyper accumulators for nickel, iron and copper as the BAFs are greater than 1. However they are comparable as the ANOVA shows they are the same.

The results indicated significant differences between the slimes and control sites for BAFs, which suggests the species are hyper accumulators for nickel, copper, iron, lead and zinc. However because the soil contents were not significantly different for nickel and arsenic, a better control site will be ideal in future studies with no abundance of this metal, so that precise conclusions are made for this metal.

The plant parts are not statistically different with respect to BAF values in other words no particular plant part leaves, bark or roots specifically accumulate the metals more than the other.

The species are also not significantly different with respect to BAFs but they are both on the high side as BAFs are greater than one except for zinc, lead and arsenic.

272

273 Table 3. Mean bioaccumulation factor for the species and plant parts at the sites

		Ni	As	Cu	Fe	Pb	Zn
SITE	SLIMES	3.32*	0.38ns	17.86***	2.57**	0.76**	1.66***
	CONTROL	0.17*	0.55ns	0.50***	0.77**	0.31**	0.11***
	S.E	0.762	0.078	2.72	1.195	0.168	0.111
SPECIES	A.POLYACANTHA	1.89ns	0.55ns	9.47ns	1.09ns	0.46ns	0.80ns
	A.SALIGNA	1.59ns	0.38ns	8.89ns	2.25ns	0.61ns	0.97ns
	S.E	0.933	0.078	2.72	1.195	0.168	0.111
PLANT PART	LEAVES	2.64ns	0.33ns	10.29ns	0.88ns	0.32ns	0.90ns
	ROOT	1.50ns	0.23ns	8.80ns	1.11ns	0.39ns	0.81ns
	BARK	1.09ns	0.84ns	8.45ns	3.02ns	0.89ns	0.95ns
	S.E	0.933	0.095	3.332	1.464	0.206	0.136

S.E=standard error . ns=no significant difference $P>0.05$ * $p<0.05$ ** $P<0.01$. *** $p<0.001$

From similar work on cabbage and broccoli grown on amended soils the overall bioaccumulation factor (BAF) of seven heavy metals in cabbage leaves and broccoli heads revealed that cabbage and broccoli were poor accumulators of Cr, Ni, Cu, Cd, and Pb (BAF <1), while BAF values were >1 for Zn and Mo (1). According to work by Maharia *et al.*, (20) the bioaccumulation factors (BAF) significantly lower BAF<1 values of Cu and Cr were found in the medicinal plants *Ocimum sanctum*, *Cassia fistula*, *Withania somnifera* and *Azadirachta Indica* Only *Withania somnifera* showed very high metal bioaccumulation BAF >1 (14).

281

282 3.5 Translocation factor

283

The results show that the metals are translocated from roots to shoots more in *A. Saligna* than *A. Polyacantha* as this species shows higher translocation factor across the metals, except for zinc whose upward mobility is comparable for the two species. The metals nickel, copper and iron are more upwardly mobile in the two species.

287




Fig 6. Translocation factors for the two species at the two sites.

Saligna/slimes – is the shoot/root quotient for *A. Saligna* at the slimes *Saligna/control* – is the shoot/root quotient for *A. Saligna* at the control site, *polyacantha/slimes* – is the shoot/root quotient for *A. Polyacantha* at the slimes *polyacantha/control* – is the shoot/root quotient for *A. Polyacantha* at the control site.

Fig 6 shows a graph of the translocation factors for the metals and the species. This means that destruction of the upper plant will be effective at removing these metals from the environment. The fast growth rate reported in literature for *A saligna* may contribute to the high translocation factor observed for the species. Both the ability of a species to accumulate high quantities of elements per biomass unit and the possibility of high biomass production over a given time and area are important.

Works by Ghafoori et al (19) confirm that the translocation factor increases with increased concentration of heavy metals (10). The ability of a species to tolerate high metal concentration makes it ideal as an accumulator. The values obtained in their study though on a different species, *Dyera costulata*, on three metals, lead, nickel and zinc also studied in this work showed maximum values of 0.62 for lead 4.00 for nickel and 3.1 for zinc, which are comparable to values obtained for a *A. Saligna* and *A. polyacantha*.

4. CONCLUSION

There was evidence of heavy metal accumulation by *A. Saligna* and *A. Polyacantha* according to data gathered in this study. These two species, *A. Saligna* and *A. Polyacantha* are two of the available options for phytoremediation at the dumps and the mine should continue to propagate them to remedy heavy metal toxicity in the environment. Harvesting and incinerating the plant will potentially facilitate remediation of the slime dams.

ACKNOWLEDGEMENTS

We appreciate the assistance made by Trojan Nickel mine as all the soil and heavy metal analysis was carried at their laboratory.

AUTHORS' CONTRIBUTIONS

Mr. Dzomba, Mr Mhandu and Ms Masvodza worked on formulating the project proposal. Mr. Mhandu and Mr. Dzomba carried out the sampling and laboratory analysis. Mr. Dzomba and Ms Masvodza wrote the manuscript. Mr Masamha assisted in statistical data analysis.

REFERENCES

1. P. Dzomba, S. Nyoni and N. Mudavanhu 2012. Heavy metal contamination risk through consumption of traditional food plants growing around Bindura town, Zimbabwe. *Journal of Toxicology and Environmental Health Sciences* Vol. 4(5): 92-95
2. Raskin I, Smith RD and Salt DE.1997. Phytoremediation of metals: Using plants to remove pollutants from the environment. *Current Opinion in Biotechnol* 8(2):221
3. Antonious GF, Kochhar TS and Coolong T 2012 Yield, quality, and concentration of seven heavy metals in cabbage and broccoli grown in sewage sludge and chicken manure amended soil. *Journal of Environmental Science and Health Part A Toxic/Hazardous Substances and Environmental Engineering* 47(13):1955-65
4. Gupta S, Nayek S, Saha RN and Satpati S. 2008. Assessment of heavy metal accumulation in macrophyte, agricultural soil and crop plants adjacent to discharge zone of sponge iron factory. *Environmental Geology* 55:731-73
5. Griffin A R, Midgley SJ, Bush D, Cunningham PJ, Rinaudo AT, 2011. Global uses of Australian acacias – recent trends and future prospects. *Diversity and Distributions* 17(5):837 - 847.
6. Ghosh M and Singh SP. 2005. A review on phytoremediation of heavy metals and utilization of its by products. *Applied and ecology and environmental research* 31:1-18.
7. Akan JC, Abdulraham FI, Sodipo OA and Lange AG. 2010. Physicochemical parameters of soil and vegetable samples from Gongulon Agricultural site, Maiduguri state Nigeria. *Journal Am. Science* 6:12.
8. Mupangwa JF, The growth performance of exotic and indigenous tree species in rehabilitating active gold mine tailings Dump at Shamva mine in Zimbabwe. *Journal of Applied Sciences and Environmental Management* (9):2
9. Bhargavi NVL and Sudha PN. 2011. Effect of Salinity and pH on the Accumulation of Heavy Metals in Sunflower (*Helianthus Annus*) Plant. Available at: <http://www.eco-web.com/edi/110706.html>
10. Chaney RL, Mielke HW and Sterrett SB. 1994. Speciation, mobility and bioavailability of soil lead. *environmental geochemistry and health* 9:105-129.
11. Coates W. 2005. Tree species selection for a mine tailings bioremediation project in Peru, short communication in *biomass and bioenergy* 28 (4): 418–423

- 373 12. Compton H. 1999. Using Phytoremediation to clean up sites.
374 www.epa.gov/superfund/accomp/news/phyto.htm
375
- 376 13. Prica M, Dalmacija, B, Roncevic S, Krcmar, D and Becelic M. 2007. A comparison of
377 sediment quality results with acid volatile sulfide (AVS) and simultaneously extracted
378 metals (SEM) ratio in Vojvodina (Serbia) sediments. Science of the Total Environment
379 389:235-44.
380
- 381 14. Peplow D. 1999. Environmental impacts of mining in eastern Washington. Center for
382 Water and Watershed Studies Fact Sheet. Seattle.
383
- 384 15. Raskin I and Ensley BD. 2000. Phytoremediation of toxic metals. eds. John Wiley and
385 Sons. New York.
386
- 387 16. Erakhrumen A and Agbontalor A. 2007. Phytoremediation an environmentally sound
388 technology for pollution prevention, control and remediation in developing countries.
389 Educational Research and Review 2(7): 15-156.
- 390 17. Erdei L, Mezsi LG, Mcs I, Vass I, Fglein F and Bulik L. 2005. Phytoremediation as a
391 program for decontamination of heavy-metal polluted environments. Proceedings of the
392 8th Hungarian Congress on Plant Physiology and the 6th Hungarian Conference on
393 Photosynthesis.
394
- 395 18. Friberg L, Nordberg GF and Vouk V. 1986. Handbook on the toxicology of metals.
396 Elsevier. Amsterdam.
397
- 398 19. Ghafoori M, Majid NM, Islam MM and Luhath S. Bioaccumulation of heavy metals by *Dyera*
399 *costulata* cultivated in sewage sludge contaminated soil. African Journal of Biotechnology
400 10(52) 10674-10682
401
- 402 20. Maharia RS, Dutta RK, Acharya R and Reddy AVR. 2010. Heavy metal bioaccumulation
403 in selected medicinal plants collected from Khetri copper mines and comparison with
404 those collected from fertile soil in Haridwar, India. Journal of Environmental Science and
405 Health Part B Pesticides Food Contaminants and Agricultural Wastes 45(2):174-81
406
- 407 21. Raskin I, Smith RD and Salt DE. 1997. Phytoremediation of metals: Using plants to
408 remove pollutants from the environment. Current Opinion in Biotechnol 8(2):221
409
- 410 22. Sastre J, Sahuquillo A, Vidal M, and Rauret G. 2002. Determination of Cd, Cu, Pb and Zn
411 in environmental samples: microwave assisted total digestion versus aqua regia and nitric
412 acid extraction. Anal. Chim. Acta, 462: 59–72.
413
- 414 23. Thompson GD, Dirk U Bellstedt, Margaret Byrne, Melissa A Millar, David M Richardson,
415 2012 Cultivation shapes genetic novelty in a globally important invader. Molecular
416 Ecology 21(13):3187-99.
417
- 418 24. Trueby P. 2003. Impact of heavy metals on forest trees from mining areas. Proceedings of
419 the International Conference on Mining and the Environment III. Sudbury. Ontario.
420 Canada.

- 421 25. Kindu M , Tadesse Y Glatzel G , Amha Y. 2012. Performance of eight tree species in the
422 highland Vertisols of central Ethiopia: growth, foliage nutrient concentration and effect on
423 soil chemical properties New Forests 32(3):285-298
- 424 26. Post JA, Kleinjan CA, Hoffmann JH and Impson FAC, 2009. Biological control of Acacia
425 cyclops in South Africa: The fundamental and realized host range of Dasineura dielsi
426 (Diptera: Cecidomyiidae). Biological Control 53(1):68-75.
- 427 27. Tadros JM. 2003. The physiology and biochemistry of the multipurpose forest trees
428 species: *Leucaena leucocephala*, *Prosopis juliflora*, *Acacia saligna*, and *Eucalyptus*
429 *citriodora* as influenced by water stress. Purdue University publication. Available at:
430 <http://docs.lib.purdue.edu/dissertations>
431