

Research paper

Evaluation of pre-emergence herbicides for weed control in cowpea (*Vigna unguiculata* (L.) Walp.) in a forest - savanna transition zone

Omovbude, Sunday and Udensi, Ekea Udensi

Department of Crop and Soil Science, University of Port Harcourt, East-West Road,
Choba P.M.B 5323 Port Harcourt, Nigeria

Corresponding authors: emails ueudensi@yahoo.co.uk or u.udensi@cgiar.org.

Abstract

Field experiment were conducted to evaluate the effect of pre-emergence herbicides on weed control in cowpea during the cropping seasons of 2005 and 2006 at the Teaching and Research Farm of Ambrose Alli University in Ekpoma (Lat. 6° 45'N, Long. 6° 8'E, Nigeria, in a forest-savanna transition zone of Nigeria. Six weed control treatments evaluated against unweeded control were; Codal gold^(R) (250g prometryne + 162.5 g metolachlor per litre) at a rate of 1.65 kg a.i ha⁻¹, Galex^(R) (Metolachlor 250g + metobromuron 250g per litre) at the rate of 3.0 kg a.i ha⁻¹; Pendilin^(R) (Pendimethalin 500g per litre) at a rate of 2.0 kg a.i ha⁻¹, one hoe – weeding at 3 weeks after planting (WAP) and two hoe weeding at 3 and 7 WAP. The treatments were laid out as randomized complete block design with four replications. Results showed that all the weed control treatments significantly ($p < 0.05$) reduced weed dry weight when compared to the unweeded control. Plots treated with herbicides and hoe weeded twice gave an acceptable ($\geq 70\%$) weed control in both years. Cowpea growth and yield were best under Codal gold at 1.65 kg a.i ha⁻¹ while the unweeded control had the poorest yield in both years. The highest profit was obtained under Codal gold at 1.65 kg a.i ha⁻¹ and the lowest in plot hoe weeded twice. The highest cost benefit ratio (CBR) of 24.61 in 2005 and 28.21 in 2006 were recorded with Codal gold^R treated plot while the lowest of 1.25 and 1.22 were obtained in 2005 and 2006 respectively under plot hoe weeded twice. Codal gold^(R) at 1.65 kg a.i ha⁻¹ was better than the other herbicides tried, in reducing weed infestation in cowpea as evidenced on the performance and yield of cowpea. With the range of herbicides tried in this study, Codal gold^(R) at 1.65 kg a.i ha⁻¹ appears to be the best herbicide for weed control in cowpea under the present study

Key words: Cowpea, Pre-emergence herbicides, forest – savanna transition zone, weed control.

1.0 INTRODUCTION

Cowpea (*Vigna unguiculata* (L) Walp.) is an annual legume that is widely cultivated in Nigeria mainly for its edible seeds. A more recent and reliable statistics, by Food and Agricultural Organization (FAO) reported that about 7.56 million tones of cowpea were produced annually on about 12.76 million hectares of land (IITA, 2007). Sub – Saharan Africa was reported to account for about 70% total World Production (IITA, 2007). Nigeria is still said to be the world largest cowpea producer where about 2 million tones are produced per annum. This is followed by Niger (650,000 tons) and Mali with 110, 000 tones (IITA, 2001).

Average world yield of cowpea grain is quite low at less than 0.3 ton/ha. Within Africa, average cowpea yields range from 0.05 to 0.55 ton/ha (Cisse *et al*, 1995). Alghali (1992) reported 200-300 kg ha⁻¹ in Nigeria and more than 2000kg ha⁻¹ under research environment (Rusoke and Rubaibage, 1994). Among other factors limiting cowpea production, inadequate weed control had been identified as a major contributory factor for yield gap. Uncontrolled weed growth and/or inadequate weed control in the crop have been reported to account for 40-80% reduction, in grain yield (Olofintoye and Adesiyun, 1989, Tijani – Eniola, 2001). Crop losses by weeds could be aggravated by delay in weeding or inability to weed through the entire crop growth period. However, studies of the thresh hold levels of weeds have shown that complete weed eliminations is not essential for high yields (Sangakkara, 1999), probably because the crop can also competed strongly with weeds, after the critical period of weed interference.

The critical period of weed competition for cowpea is the first 3 – 4 weeks of crop growth (Akinyemiju and Echendu, 1987). Weed competition has the capability of lowering vegetative growth, flowering, fruiting and seed production, which will determine yield. Whenever possible, several weed control methods which include, cultural, biological or chemical could be used separately or in combination to give the desired result (Gibson *et al* 2007).

Weed reduced crop yield by interfering with crop growth. Hand weeding limits agricultural productivity because there is a limit to the amount of land area that can be weeded even when the labor is free (Akobundu, 1987). Hand weeding required over 50% of the farmers' time leaving him and his family with little or no time for

other activities (Akobundu, 1987). Use of herbicides may therefore provide a timely and adequate alternative to hand weeding as this will not only remove the drudgery associated with it but also lower the cost of weeding and provide protection for crop against early weed competition when pre-emergence herbicides are used (Akobundu, 1987).

Presently, existing herbicides for pre-emergence weed control in cowpea are no longer **consistence** in their spectrum of **weeds effectively control**. This may be as a result of some weeds becoming **tolerant** to these herbicides. Codal gold^(R) which is one of the recently introduced new **herbicide** in the early twenty (20th) **centuries** in Nigeria had not **been** experimented on by farmers. Hence, the objective of this study was to assess the effectiveness of old herbicides (Galex^(R) and Pendilin^(R)) and Codal gold^(R) (new herbicide) for weed control and also their economic returns on cowpea in the forest savanna- transition zone.

2.0 MATERIALS AND METHOD

2.1 Experimental site

The experiment was conducted **in** between August and December, 2005 and repeated in the same season of 2006 at the Teaching and Research Farm of Ambrose Alli University (Latitude 6° 45¹, 6° 8¹ E and an altitude of 313 metres above sea level) in the forest – savanna transitions zone of Nigeria. The experimental site was under short fallow for about two (2) years, previously cropped to maize. Pre-cropping vegetation in the experimental site was dominated by Siam weed (*Chromoleana odorata* (L) R.M. King & Robinson), goat weed (*Ageratum conyzoides* Linn) ,broom weed (*Sida acuta* Burm F) ,guinea grass (*Panicum maximum* Jacq) and Cyperus spp. Soil **sample** were randomly collected from 10 spots (0-15cm depth) over the entire field using an auger before the commencement of the experiment. The soil samples were bulked and mixed thoroughly for analysis. The soil analysis results are presented in Table 1.

Table 1: Physic- chemical properties of the top soil (0-15cm) before cropping the experimental site in 2005 and 2006

Soil Properties	Values	
	2005	2006
Sand (g/kg)	950.00	951.00
Silt (g/kg)	45.00	46.00
Clay (g/kg)	5.00	3.00
p ^H (H ₂ O, 1.1)	6.20	6.50
Organic carbon (g/kg)	14.00	12.00
Total N (g/kg)	0.90	0.70
Available P (mg/kg)	15.40	16.78
Exchangeable cations (cmol/kg)		
Ca	3.90	3.70
Mg	2.10	1.90
K	0.29	0.26
Na	0.30	0.24
TE A (cmol/kg)	0.60	0.40
ECEC (cmol/kg)	7.19	6.50

TE A = Total Exchangeable Acidity

ECEC = Effective Cation Exchange Capacity

The soil in both years was neutral sandy, moderate organic carbon, low nitrogen and acidic. Phosphorus (P) and Potassium (K) values were adequate. Number of rainy days and total rain fall during the trials are presented in Table 2.

Table 2 Monthly and Total Rainfall during the trial, 2005 – 2006

Months	2005		2006	
	Rainy Days	Rainfall Days (mm)	Rainy Days	Rainfall Days (mm)
January	1	60.2	2	5.3
February	3	107.0	2	9.2
March	3	45.2	6	96.9
April	6	108.0	5	112.0
May	10	180.9	13	234.1
June	17	376.8	15	292.7
July	21	439.6	19	449.9
August	7	38.3	15	348.1
September	20	351.6	28	554.2
October	12	320.2	19	41.2
November	2	145.7	1	2.2
December	2	3.2	0	0.0
Total	104	2176.7	125	2145.8
Mean	8.69	187.4	10.42	178.8

Source: Edo State Agricultural Development Project (EADP), Irrua, Edo State, 2005 and 2006

The total rainfall was 2176.7mm distributed over 104 days in 2005 and 2145.8mm distributed in 2006 over 125 days in 2006 (Table 2).

2.2 Field Procedures

The land was manually slashed, stumped before leveling the soil surface with spades. Plot size was 3m×4m with an alley way of 1m among plots and between replicates. There was thus, a total of 28 plots occupying an experimental area of 23m× 19m (437m²) corresponding to approximately 0.04ha. There were six (6) treatments involved in the experiment and these were no weeding(control), weeding once at 3 weeks after planting (WAP), weeding twice at 3 and 7 WAP, Galex^(R) at 3.0kg a.i./ha, Codal gold^(R) at 1.65 kg a.i /ha and pendilin^(R) at 2.0kgai/ha. The chemical names of the herbicides are shown in Table 3.

Table 3: Chemical names of Herbicides used in the Study

Trade name	Chemical name
Pendilin [®]	[N-(1-ethylpropyl) – 3,4 – dimethyl 1-2, 6 dinitro benzenenamine]
Galex [®]	Metobromuron,(2-chloro – N-(2-ethyl – 6- methyl phenyl) – N-(2- methoxy -1-methyl ethyl acetamide + Metolachlor (N ¹ – 4 (4- bromophenyl) – N- methoxy – N- methyl urea]
Codal gold [®]	Prometryne, [N, N ¹ – bis (1-methyl ethyl) – 6- (methylthio) – 1, 3, 5-triazine – 2, 4-diamine + Metolachlor (2 chloro – N- (2-ethyl – 6- methyl phenyl) – N-) (2- methoxyl – 1- methyl ethyl) acetamide]

The cowpea cultivar Ife brown (IT84S-2246-4) used for the experiment was obtained from International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria. Cowpea was planted on 30th of August in 2005 and 31st of August 2006 at a spacing of 60cm×25cm. Two seeds were planted per hole and the seedling thinned to one per stand at 2WAP, giving a desired population of 66667 plants/ha. No fertilizer was applied to the crop stand. The herbicides were applied at the specific rate of each herbicides using CP3 knapsack sprayer calibrated to deliver approximately 250L/ha at 210kpa using a red poliject nozzle (chikoye *et al.*, 2005). The first weeding was done by hoeing at 3WAP and the second weeding at 7 WAP in both years.

2.3 Data collection and Analysis

Weed density was recorded at final harvest. Weed samples were collected from 1m x 1m quadrat per plot, bulked and oven dried at 80°C for 48 hours. Weed smothering efficiency (WSE) of the different weed control methods was calculated according to Subramanian *et al.* (1991).

$$WSE\% = \frac{WDWC - WDWT}{WDWC} \times \frac{100}{1}$$

Where:

WSE = Weed smothering Efficiency

WDWC = Weed dry weight of unweeded control

WDWT = Weed dry weight in treated plots

The crop parameters determined were:

Number of leaves/ Plant: The number of leaves obtained by direct physical counts by randomly selecting six (6) plants from the net plots (3m²) size and tagged. The average of the 6 plants was taken as per plant.

Plant height: This was done by randomly selecting 6 plants from the net plot size of 3m² and measured their vine length from the soil level to the tip of the apical bud with a measuring tape. The mean of the plant height was recorded per plant.

Leaf area: This was done by randomly selecting 6 plants and tagged. The leaf area was determined by graph tracing method. The total leaf area per plant was calculated as leaf area multiplied by the number of leaves using Remison (1997) procedures.

Days to 50% flowering: This was estimated by counting the number of days from sowing to the time when half of the plants start to produce flowers.

Number of seeds/plant: This was estimated as:

$$\frac{\text{number of seeds per net plot}}{\text{number of plants per net plot}}$$

Number of pods/plant: This was estimated as:

$$\frac{\text{number of harvested pods per net plot}}{\text{number of plants per net plot}}$$

1 **2.4 Cowpea grain yield:** Harvesting was assessed from net plot of 3m². The
 2 pods were handpicked and four pickings were done at weekly intervals starting from
 3 56 days after planting (DAP). The pods were sun dried for one week and then
 4 shelled. The grain yield in each net plot was weighed with weighing balance and the
 5 weight recorded. The yield per net plot was extrapolated to kilograms hectare⁻¹
 6 (kg/ha).

7 **2.5 Economic Assessment:** Economic evaluation of the weed control treatment
 8 was done using partial farm budgeting (Okoruwa *et al.*, 2005) at prevailing labor and
 9 market costs of materials to estimate revenue. Sale revenue was obtained by
 10 multiplying the final grain yield (kg ha⁻¹) by the market price (₦ kg⁻¹). This is
 11 represented below in the following formula:

12 (i) Revenue = Yc x Pc (where, Yc = cowpea yield in kilogram/ ha; Pc = Price of
 13 cowpea grain.

14 The profit was calculated by subtracting the costs of production from the sale
 15 revenue represented as follows:

16 (ii) Profit (net revenue) = Revenue – Total cost of Production

17 (iii) Simple proportion of total cost of weed control (cost of production) and net
 18 revenue (profit) were used to determine the cost/ benefits ratio (CBR) of
 19 each of the weed control treatment as follows:

20 Cost Benefit Ratio (CBR) = Profit (net revenue)/ Total cost of production.

21

1

2 3.0 RESULTS

3 3.1 Weed growth and Control

4 Weed density and weed dry weight were significantly influenced by weed
5 control treatments at final harvest in 2005 and 2006 cropping seasons (Table 4).

6 **Table 4: Effect of the herbicide treatments on weed density, biomass, and**

7 **control at harvest in 2005 and 2006 cropping season**

Treatment	Rate (kg a.i./ha)	Weed density (no/m ²)		Weed dry weight (g/m ²)		Weed smothering efficiency (WSE) (%)	
		2005	2006	2005	2006	2005	2006
No weeding	-	197.00	222.25	118.85	202.00	-	-
Codal gold ^(R)	1.65	61.25	67.00	15.58	19.50	87.08	90.35
Galex ^(R)	3.0	90.00	116.00	19.93	29.75	82.72	85.27
Pendilin ^(R)	2.0	90.25	117.00	20.78	28.00	83.06	86.07
Weeded x 1 (3WAP)	-	125.00	151.25	85.00	84.25	29.19	58.67
Weeded x 2 (3 + 7WAP)	-	90.25	115.75	23.50	31.00	80.18	84.56
LSD(P = 0.05)		4.667	9.548	7.139	2.510	1.134	3.286

8

9 All the herbicides reduce weed density considerably over the weedy checks. Codal
10 gold^(R) at 1.65 kg a.i /ha was most effective in suppressing weeds compared to the
11 rest of the treatments. The differences observed on weed density for plots treated
12 with Galex^(R) at 3.0 kg a.i /ha, Pendilin^(R) at 2.0 kg a.i /ha and plots hoe – weeded
13 twice were not significant, but they differ significantly from plots hoe weeded once.
14 Weed dry weight was significantly reduced by herbicide application in both years.

15 In 2005 the lowest weed dry weight was recorded in Codal gold^(R) treated
16 plots at 1.65 kg a.i /ha but it was at similar with Galex^(R) at 3.0 kg a.i /ha and
17 Pendilin^(R) plots at 2.0 kg a.i /ha but differs significantly from plots hoe-weeded twice.
18 Though plot hoe weeded twice had high weed dry weight, it was comparable with
19 that of Galex^(R) at 3.0 kg a.i /ha and Pendilin^(R) treated plots at 2.0 kg a.i /ha. In 2006,
20 plots hoe- weeded twice differ significantly (P < 0.05) in dry weight from Pendilin^(R)
21 treated plots at 2.0 kg a.i /ha, but it had a comparable value with that of Galex^(R)
22 treated plot at 3.0 kg a.i /ha. Weed smothering efficiency in both years was distinctly
23 higher in Codal gold^(R) plots at 1.65 kg a.i /ha than Pendilin^(R) treated plots at 2.0 kg

a.i /ha, Galex^(R) treated plot at 3.0 kg a.i /ha, plot hoe weeded twice (3 and 7 WAP) and plot hoe weeded once (3 WAP) in that order.

3.2 Vegetative Growth Characteristics

Leaves produced in unweeded plots was significantly lower than the treatment plots. Plots treated with Codal gold^(R) at 1.65 kg a.i /ha had the highest number of leaves. There were no significant differences ($P>0.05$) among Galex^(R) at 3.0 kg a.i /ha, Pendilin^(R) at 2.0 kg a.i /ha and plots hoe weeded twice in leaf production but they differ significantly from plots hoe – weeded once (Table5).

Plant height was significantly influenced by different herbicides treatments (Table5). Among the treatments Codal gold^(R) at 1.65 kg a.i /ha produced the tallest plants while the shortest plant was observed in weedy check. There were no significant different among Galex^(R) at 3.0 kg a.i /ha, Pendilin^(R) at 2.0 kg a.i /ha and plot- hoe weeded twice since uniform plant height were observed in them, but they differ from plot hoe weeded once, which had medium plant height.

Leaf area showed significance difference ($P<0.05$) among the treatments. Codal gold^(R) plot at 1.65 kg a.i /ha recorded the highest leaf area which differs significantly from other treatments. There were no significance difference in cowpea leaf area in Galex^(R) at 3.0 kg a.i /ha, Pendilin^(R) 2.0 kg a.i /ha and plots hoe weeded twice but they differ from plots hoe- weeded once.

Table 5: Effect of weed control treatments on cowpea height and leaf area

Treatment	Rate (kg a.i./ha)	Plant height (cm)		Leaves /plant (no)		Leaf Area /plant (cm ²)	
		2005	2006	2005	2006	2005	2006
No weeding	-	64.75	72.78	16.50	17.25	511.2	614.0
Codal Gold ^(R)	1.65	107.48	115.25	31.75	35.25	1591.6	1735.0
Galex ^(R)	3.0	97.10	102.50	25.50	27.25	1178.9	1271.3
Pendilin ^(R)	2.0	97.03	103.00	25.75	27.75	1191.7	1272.3
Weeded x 1 (3WAP)	-	89.88	91.75	20.00	22.00	817.0	910.8
Weeded x 2 (3+7WAP)	-	96.03	102.00	25.50	27.00	1173.0	1268.3
LSD(P = 0.05)		2.503	2.021	1.150	1.499	55.22	65.35

3.3 Effects of weed control treatment on yield and yield components of cowpea.

In 2005, there were no significant difference ($P>0.05$) among the herbicides treated plot and plot that were hoe weeded twice in terms of number of days to 50% flowering but they differ significantly from plot hoe weeded once and unweeded plots. (Table 6) The unweeded plot took the highest time to reach 50% flowering which was comparable to plots hoe weeded once. In 2006, unweeded plots differ significantly from other treatments by the time it took to reach 50% flowering. Codal gold^(R) at 1.65 kg a.i /ha had the shortest number of days to 50% flowering but it was not significantly difference from Galex^(R) at 3.0 kg a.i /ha, Pendilin^(R) at 2.0 kg a.i /ha and plot hoe weeded twice which differ significantly from plot hoe weeded once. (Table 6)

In 2006, Codal gold^(R) treated plot at 1.65 kg a.i /ha had significantly the highest number of pod than other treatments. The second highest number of pods was produced in plots treated with Pendilin^(R) at 2.0 kg a.i /ha but it was not different from Galex^(R) treated plot at 3.0 kg a.i /ha and plot hoe weeded twice. Plot hoe weeded once differ significantly from the rest of the treatment including the weedy check which had the lowest number of pods (Table 6)

In 2005, there was no significance difference ($P>0.05$) among the treatment although Codal gold^(R) at 1.65 kg a.i /ha had a slight number of seeds/pod. In 2006, the number of seeds/ pod differs significantly. Weedy plots recorded the lowest number of seeds/pod but it was not different from plots hoe weeded once. Codal gold^(R) treated plots at 1.65 kg a.i /ha recorded highest number of seeds/pod but the value was comparable with that of Pendilin plots at 2.0 kg a.i /ha. No significance difference was observed among Galex^(R) at 3.0 kg a.i /ha, Pendilin^(R) at 2.0 kg a.i /ha and plots that were hoe weeded twice. The grain yield of cowpea was significantly ($P<0.05$) affected by the weed control treatments in the two years of study (Table 6). Cowpea grain yield followed the trend Codal gold^(R) at 1.65 kg a.i /ha > Pendilin^(R) at 2.0 kg a.i /ha > Galex^(R) at 3.0 kg a.i /ha > Two hoe weedings > one hoe weeding > weedy plots.

Table 6: Effect of weed control treatment on yield and yield components of cowpea

Treatment	Rate (kg a.i /ha)	Days to 50 flowering		Pods/plant (no)		Seed \pod (no)		Grain yield (kg/ha)	
		2005	2006	2005	2006	2005	2006	2005	2006
No weeding	-	52	52	9	11	10	9	814.00	905.25
Codal Gold ^(R)	1.65	46	46	13	14	11	13	1652.50	1841.25
Galex ^(R)	3.0	46	46	11	13	10	12	1180.00	1236.50
Pendilin ^(R)	2.0	46	46	11	13	10	12	1185.83	1240.75
Weeded x 1 (3WAP)	-	50	50	10	12	10	10	999.93	1085.50
Weeded x 2 (3+7WAP)	-	46	46	11	13	10	12	1171.67	1230.50
LSD(P = 0.05)		3	21	2	0.6	NS	1	81.528	36.085

3.4 Economic Evaluation

The value of the output per hectares (total revenue) from the use of pre-emergence herbicides varied from ₦119, 991.6 to ₦198, 300 in 2005. (Table 7) The highest revenue (₦198, 300) was recorded under Codal gold^(R) plot at 1.65 kg a.i /ha followed by Pendilin^(R) treated plot at 2.0 kg a.i /ha while the lowest was recorded under plot hoe weeded once (₦119, 991.6) compared to the rest of the treatment. Similar trend was observed in 2006 with Codal gold^(R) at 1.65 kg a.i /ha recording the highest revenue (₦293, 362.5) followed by Pendilin^(R) treated plot at 2.0 kg a.i /ha while the lowest was recorded under plot hoe weeded once (₦141, 115) compared to the rest of the treatment. (Table 8) The above results were due to differences in yield/ha recorded by the different treatments with Codal gold^(R) at 1.65 kg a.i /ha resulting in the highest yields.

Table 7: Economics evaluation of the use of pre emergence herbicides in cowpea production in 2005 cropping season

Treatment	Rate (kg a.i./ha)	Rate (L/ha)	Herbicides cost N/ha	Time of treatment application (man – hr/ha)	Cost of treatment application (N/ha)	Total cost N/ha	Grain yield (kg/ha)	Revenue (N/ha)	Net Revenue (profit) N/ha	CBR
Codal gold ^(R)	1.65	4	7000	5	850	7,850	1,652.50	198,300.00	190,450.00	24.26
Gale ^(R)	3.0	5	6000	5	850	6,850	1,180.00	141,600.00	134,750.00	19.67
Pendilin ^(R)	2.0	5	6500	5	850	7,350	1,185.83	142,299.60	134,949.60	18.36
Weeded x1	3 WAP	-	0.00	120	31,200	31,200	999.93	119,991.60	88,791.60	2.85
Weeded x 2	3 + 7 WAP	-	0.00	240	62,400	62,400	1,171.67	140,600.40	78,200.40	1.25
Unweeded	-	-	0.00							

CBR = Cost of benefit Ratio = Net Revenue / Total Cost

1kg of cowpea = ₦120.00

1litre of Codal gold[®] = ₦1, 750.00

1litre of Gale[®] = ₦1, 200.00

1litre of Pendilin = ₦1, 300.00

Cost of application of herbicides = 150/hr. knapsack (hired) = ₦100 / treatment

Cost of hand weeding = ₦260/hr (₦600- ₦700 = (₦1300 average) between 7.00am – 12noon = 5 hours approximately

Table 8: Economics evaluation of the use of pre-emergence herbicides in cowpea production in 2006 cropping season

Treatment	Rate (kg a.i./ha)	Rate (L/ha)	Herbicides cost N/ha	Time of treatment application (man – hr/ha)	Cost of treatment application (N/ha)	Total cost N/ha	Grain yield (kg/ha)	Revenue (N/ha)	Net Revenue (profit) N/ha	CBR
Codal gold ^(R)	1.65	4	7200	5	1,020	8,220	1,841.25	239,362.50	231,142.50	28.12
Galex ^(R)	3.00	5	6250	5	1,020	7,270	1,236.50	160,745.00	153,475.00	21.11
Pendilin ^(R)	2.00	5	6750	5	1,020	7,770	1,240.75	161,297.50	153,527.50	19.76
Weeded x1	3 WAP	-	0.00	120	36,000	36,000	1,085.50	141,115.00	105,115.00	2.92
Weeded x 2	3 + 7 WAP	-	0.00	240	72,000	72,000	1,230.50	159,965.00	87,965.00	1.22
Unweeded	-	-	0.00							

CBR = Cost of benefit Ratio = Net Revenue/Total Cost

1kg of cowpea = ₦130.00

1litre of Codal gold^(R) = ₦1, 800.00

1litre of Galex^(R) = ₦1, 250.00

1litre of Pendilin^(R) = ₦1, 350.00

Cost of application of herbicides = 180/hr. knapsack (hired) = N120 / treatment

Cost of hand weeding = ~~N300/hr~~ (N700- N800) = (N1500 average) between 7.00am – 12noon = 5 hours approximately

Plots hoe weeded twice at 3& 7WAP recorded the highest cost of production (N62, 400) followed by plots hoe weeded once (N31, 200) compared to other treatments while the lowest was recorded under Galex^(R), treated plots at 3.0 kg a.i /ha (N6, 850) in 2005. In 2006, the same trend was observed with the highest cost of production incurred by weeding at 3&7WAP compared to all other treatments while Galex[®] treated plots at 3.0 kg a.i /ha resulted in the lowest cost of production.

In 2005, the highest net revenue (profit) (N190, 450) was obtained under Codal gold^(R) plots at 1.65 kg a.i /ha and the lowest under plots hoe-weeded twice (N78, 200.4). Similar trend was observed in 2006 with Codal gold^(R) at 1.65 kg a.i /ha having the highest profit of (N231, 142.5) and the lowest in plot hoe weeded twice (N87,965). The highest cost benefit ratio (CBR) (24.26) was recorded in Codal gold^(R) plot at 1.65 kg a.i /ha and the lowest in plot weeded twice (1.25). In 2006, the same trend was observed, Codal gold plot at 1.65 kg a.i /ha had the highest CBR (28.12) and the lowest in plot hoe weeded twice (1.22).

4.0 DISCUSSION

The high weed density recorded in the weedy plots in both years which invariably result to high weed dry weight could be attributed to low ground coverage by cowpea vines. Also, the low weed density observed in the herbicides treated plots could be attributed to effective weed control of the herbicides and their ability to control weed beyond the critical period of cowpea growth. The adequate weed coverage by cowpea vine led to smothering effect of the weeds judging from low weed population and low weed dry weight which invariably led to increase in weed smothering efficiency (WSE).

Hoe – weeding once at 3WAP was obviously not adequate for effective weed control in cowpea since it was not comparable to twice hoe weedings. The low weed density and low dry weight in plot hoe weeded twice results to high WSE. The cowpea vines were able to smothered weeds in plots hoe weeded twice compared to plot hoe weeded once which had moderate increase in weed dry weight. The moderate increase in weed dry weight could be attributed to frequent re-occurrence and persistent characteristics of weeds.

Weeding and herbicide application significantly encouraged vigorous growth of cowpea. Weed competition delayed the number of days to 50% flowering and

considerably reduced the yield and yield components of cowpea in the unweeded. Remison (1978) reported that low yield obtained from weedy plots may be due to competition from natural flora. The higher grain yields obtained from the herbicide – treated plots compared to hand-weeding could be due to minimal weed competition with cowpea and herbicides treated plots. Weedy plots gave the lowest grain yield as result of intense weed competition. The percentage yield reduction in the current study was 30.53% in 2005 and 26.43% in 2006. This study has confirmed earlier reports (Akobundu, 1979; Sangakkara, 1999) of a possible yield loss due to weed infestation in cowpea.

The yield difference per hectare recorded by the different treatments account for the variation observed in value of output/ha (total revenue) in both years. Codal gold^(R) at 1.65 kg a.i /ha was the most profitable method of weed control in both years. It had the highest grain yield coupled with low weed control cost. Although grain yield was high in plots hoe weeded twice, the cost of weeding was also high consequent upon scarcity of labor at time of weeding, thereby reducing the profit. This confirms the report of Adigun and Lagoke (2003) that hoe weeding is expensive. It also shows the advantage of herbicide application over hoe weeding in reduction of cost of production in cowpea. This agrees with the findings of Nazear *et al.* (2004) that a very lucrative cost – effective was recorded due to application of bactril-M on wheat and the gain in yield from hand weeded plot was nullified with additional cost of weeding. The results obtained from economic evaluation is also in conformity with that of Chikoye *et al.* (2005) who noted that chemical control is a better alternative to manual weeding because it is cheaper, faster and gives better weed control.

5.0 CONCLUSION

The applied herbicides enhanced cowpea productivity, through enhanced growth and yield of cowpea by reducing weed infestation. Consequent on the condition of the present study and spectrum of herbicide use or tried, Codal gold^(R) at 1.65 kg a.i /ha seem to be the best or most effective herbicide for reducing weed infestation in cowpea.

REFERENCES

- Adigun, I.A and Lagoke. S.T. O (2003). Weed control in transplant rain fed and irrigated tomatoes in the Nigerian savanna. *Nigerian Journal of Weed Science* 16:23-29
- Akinyemiju, O.A and Echendu, T.N.C (1987). Influence of tillage method and pre-emergency herbicide on weeds control in cowpea (*Vigna unguiculata* (L.) Walp). *Crop Protection* 6 289 – 294
- Akobundu, I.O (1979). Weed control strategies for multiple cropping systems of the humid and sub-humid tropics. In Akobundu, I.O. (Ed), weeds and their control in the humid and sub-humid tropics. IITA Proceedings series 3:80 -100.
- Akobundu, I.O (1987). *Weed Science in the Tropics. Principles and Practices*. 2nd Edition, John Wiley and Sons Inc Chichester 522p.
- Alghali, A.M (1992). Insecticide application and schedules to reduce grain yield losses caused by insect pests of cowpea in Nigeria. *Insect Science Applied* 13:725-730.
- Chikoye, D. Udensi, U.E and Lum, A.E (2005) Evaluation of new formulation of atrazine and metolachlor mixture for weed control in maize *Crop Protection* 24:1016-1020.
- Cisse N. Ndiaye, M. Thiaw, S. Hall. A.E (1995). Registration of Mouride cowpea *Crop Science* 35:1215-1216.
- Gibson, D.J, Miller, K.; Delong, M.; Commelly, J.; Kiwan L.; Wood A.J.; and Breyan G. (2007). The weed community effects on yield and quality of soybean (*Glycine Max* (L) Merk). *Journal of the Science of Food and Agriculture* 88 (3) 371 – 381.
- International Institute of Tropical Agriculture (IITA) (2001) Crop and farming systems. <http://www.IITA.org/cowpea.int.ml>.
- International Institute of Tropical Agriculture (IITA) (2007). Cowpea. IITA annual report. [http://www.IITA.org/cons/details/cowpea project details.aspx? Zonied 63 & artid=269](http://www.IITA.org/cons/details/cowpea%20project%20details.aspx?Zonied63&artid=269).
- Nazeer, H.S.; Gal, H.; Sad, U.R.; Nazir, A. and Eazles, S. (2004). Weed management in wheat on farmer field DASP command area. *Weed Science Society of Parkistan* 10 (1-2):25-35.
- Okoruwa, V.O.; Obadaki, F.O and Ibrahim ,G. (2005). Profitability of beef cattle farming fattening in the cosmopolitan city of Ibadan, Oyo State. *Moore Journal of Agricultural Research* 6 (1): 45-51

- Olofintoyen, J.A and Adesiyun, A (1989). Weed control in cowpea (*Vigna unguiculata* (L) Walp) with sethoxydim and galex *Nigerian Journal of weed Science* 2:29-34
- Remison, S.U (1978). The performance of cowpea (*Vigna unguiculata* (L) Walp) as Influenced by Weed Competition *Journal of Agricultural Science* Cambridge 90:523-530
- Remison S.U (1997). *Basic principles of crop physiology*. Sadoh Press, Benin City, Nigeria, 320p.
- Rusoke, D.D, Rubaihayo. P.(1994). The Influenced of some crop Protection Management Practices on Yield Stability of Cowpea. *Africa crop science journal* 2:143-146.
- Sangakkara, U.R (1999). Effect of weeds on yield and seed quality of two tropical grain legumes. *Tropical Science* 39;227-232.
- Subramaniam, S. Alli, A.M and Kumar R.J. (1991) All about weed control. Kalyani Publisher, New Delhi – 110002, India, 315p.
- Tijani – Eniola, H. (2001). Influenced of intra –row spacing and weeding regime on the performance of cowpea (*Vigna unguiculata* (L) Walp). *Nigerian Journal of weed science* 14:11-15