

Research Paper

Sugarcane Genotype Performance in Three Environments (Based on Crop Cycle) at Mardan, Pakistan

Comment [K1]: This is not G X E study but, it concerns different cropping systems. See: Sanchez (1976), p. 478. See Ottman (1996) for the definition of G x E study.

ABSTRACT

Sugarcane breeders often face significant genotype x environment interactions in their trials grown under multiple environments. Hence, genotypes need to be tested for their stability across different environments keeping in view the significant interactions. An experiment comprising 28 sugarcane genotypes (including 2 checks) was planted in two plant and one ratoon crops during 2010 to 2013 in a randomized complete block design with 3 replications. Data were recorded on cane, yield, and quality characters. Analyses of variance showed significant mean squares for crops, genotypes, and their interactions. The linear contrasts of two plant crops were found non-significant for tillering, vigor rate, stalk diameter, Brix, Pol, recovery, and cane yield. However, the contrast for plant crops versus ratoon was non-significant for stalk diameter only. Shukla's stability variances and yield stability indices (Ys_i) showed no stable character for all the that no single genotype except for cane yield recorded for was stable for all characters. However, genotypes MS-2003-CP-209, MS-2003-CP-275, CoJ-76, MS-2003-CR2-131, and MS-2003-CR5-245 as stable genotypes were stable for cane yield. The results of the study indicate the importance of genotype x environment interaction and stability in the ongoing varietal development program.

Key words: Sugarcane, Stability, Mardan, G x E interaction, Mardan, Pakistan

INTRODUCTION

Sugarcane genotypes in advanced evaluation stages need to be tested for their stability across different environments. Due to the varying interaction of genotypes with the environment the performance ranking of sugarcane genotypes is rarely the same across the environments they are tested in. Kang and Martin (1987) reported that underlying genotype x environment interaction reduced the correlation between phenotypic and genotypic values. Non significant genotype x location, genotype x year, and genotype x location x year interactions would indicate that little or no attention needs to be paid to locations and years in estimating genotypic effects and performance of the genotypes could be assessed based on one environment. However, if interaction of these sources of variation is significant, its causes, nature, and implications should be assessed. They further stated that a genotype would be regarded as stable if its associated interaction component is less than or equal to

38 | environmental variance (experimental error) and would be considered unstable otherwise
39 | [\(Ref???\)](#).

40 | According to Singh (1983), sugarcane is sensitive to environmental changes, with
41 | special regards to commercial sugar per unit area. Jackson and Hogarth (1992) reported that
42 | sugarcane clones interacted more with locations than crop-year in Australia. They suggested
43 | that for early stages of selection where many clones are under evaluation, testing only plant
44 | crops might be satisfactory. Comparatively larger genotypic mean squares showed that
45 | cultivars differed in their genetic potential for tons of cane per hectare (TCH) and Pol% of
46 | cane. Bissessur *et al* (2000) studied 154 sugarcane genotypes belonging to four families at
47 | two sites under both plant and ratoon conditions. They found significant differences between
48 | families, genotypes, and sites in stalk height, stalk diameter, recoverable sucrose % cane,
49 | and TCH. Significant $G \times E$ interactions were also found for stalk and yield characters. They
50 | concluded that selection for each specific environment should be carried out separately.

51 | Interaction of genotype, environment, and time of harvest and their implications for
52 | growers was studied by Gilbert *et al* (2006). The effect of these factors was assessed on
53 | kilograms of sugar per ton of cane (KST), TCH, and tons of sugar per hectare (TSH) for
54 | recently released cultivars in south Florida, USA. Cultivar, environment, time of harvest and
55 | their interactions significantly affected KST, TCH, and TSH. They concluded that significant
56 | [interaction between cultivar and environment justifying interactions supported](#) continued
57 | multi-location evaluation of sugarcane germplasm both during genotype selection and
58 | [subsequent released following cultivars release](#). In the same way, Zhou *et al* (2012) stated
59 | that genotype by environment interactions ($G \times E$) complicated the selection process and
60 | hindered the true effects of the genotypes. They found significant differential responses of
61 | genotypes across locations for irrigated and coastal long-cycle programs, indicating that it
62 | was important to identify and characterize sites in South Africa. Genotype by crop-year
63 | interaction was found more significant and larger in rainfed than in irrigated cropping systems.
64 | These results led them to conclude that ratooning ability was important in rainfed regions.
65 | They recommended that separating the coastal hinterland and coastal average potential were
66 | good for reducing $G \times E$ [interaction](#).

67 | The present study was planned to study the effect of different environments in the
68 | form of two plant and one ratoon crops on sugarcane genotypes across two years at Mardan,
69 | Pakistan, and attempt to identify genotypes that are stable across these environments.

70 |

71

72 MATERIALS AND METHODS

73 Twenty-six sugarcane genotypes and two checks were grown at Sugar Crops Research
74 Institute (SCRI), Mardan, during 2010 to 2013. The experimental design was a randomized
75 complete block with 3 replications. The first plant crop was planted in September 2010, while
76 2nd plant crop was planted in September 2011. A ratoon crop was maintained in 2011 as well.
77 Plot size for a genotype per replication was 6.7 x 10 meters (67 m²). There were 7 rows per
78 plot, 0.90 meters apart with a row length of 10 meters. The number of buds in the central row
79 was kept at 150. All agronomic and cultural practices were kept uniform across all plots. Data
80 were recorded on the following characters:

- 81 1. First Tillering: counting the number of tillers per 10 m central row, in the first week of
82 April.
- 83 2. Second Tillering: counting the number of tillers per 10 m central row, one month after the
84 first tillering.
- 85 3. Vigor Rate 1 (VR1): the difference of the two tillering counts divided by number of days
86 of the month as per the formula below (Tahir *et al.*, 2013).

$$87 \quad VR1 = \frac{(2ndTillering - 1stTillering)}{30}$$

- 88 4. First Growth: length of the standing plant from the ground to the top (rosette of leaves)
89 in centimeters, recorded during the first week of July.
- 90 5. Second Growth: length of the standing plant from the ground to the top in centimeters,
91 one month after the first growth.
- 92 6. Vigor Rate 2 (VR2): the difference of the two growth data divided by number of days of
93 the month as per the following formula (Tahir *et al.*, 2013).

$$94 \quad VR2 = \frac{(2ndGrowth - 1stGrowth)}{30}$$

- 95 7. Cane length: the length of the cane in centimeters to the point where tops were easily
96 removable.
- 97 8. Number of internodes per cane: counting the number of internodes per cane stalk.
- 98 9. Internode length (cm): length of the internode in the center of the plant in centimeters.
- 99 10. Stalk diameter (cm): Diameter of the stalk in centimeters with a digital Vernier caliper.
- 100 11. Millable canes: counting the number of millable canes (i.e. excluding the tillers which
101 have not developed into mature canes) in the central row of the plot.
- 102 12. Cane yield (tons per hectare): weighing the canes per plot without trash and converting
103 to tons of canes per hectare as follows:

$$104 \quad CaneYield = \left(\frac{XX \times 1000}{6.7 \times 1000} \right); \text{ Where } x = \text{cane yield in kgs per plot.}$$

13. Brix percentage: Five canes per sample were crushed using a cane crusher for estimation of Brix. Both Brix and temperature reading were taken with a hydrometer. Then, corrected Brix % was calculated using a Schmitz table (Spencer and Mead, 1948) for a particular temperature.

14. POL %:
Cane juice left from Brix reading was added with 1.5 g lead acetate and filtered. The filtered juice was then placed in a tube in a polarimeter. The reading taken was Pol% which was corrected for a particular Brix using Schmitz table (Spencer and Mead, 1948) to obtain corrected Pol%.

15. Purity %: was calculated by the following formula:

$$\text{Purity \%} = \frac{\text{POL \%}}{\text{Corrected Brix}} \times 100$$

Where Corrected Brix was the Brix adjusted with the ambient temperature.

16. Recovery %: Calculated by the following formula:

$$\text{Recovery \%} = [\text{POL \%} - 0.5 (\text{C. Brix} - \text{Pol \%})] \times 0.7$$

Data across 3 crops viz. 2 plant crops and 1 ratoon were analyzed to quantify genotype x environment interaction effects for all traits using [PROC Generalized Linear Model procedure](#) of SAS 9.1 (SAS Institute, 2003). A fixed effect model was used for genotypes and crops. When a G x E interaction was significant, means were calculated across three crops and 'stability variances' were calculated. According to Shukla (1972) a genotype is stable if its stability variance (σ_i^2) is equal to the environmental variance (σ_e^2) which means that $\sigma_i^2 = 0$. A relatively large value of σ_i^2 indicates greater instability of genotype i. As the stability variance is the difference between two sums of squares, it can be negative. These negative estimates of σ_i^2 may be taken as equal to zero (Shukla 1972). "Stability.par" procedure of the package "Agricolae" of R version 2.14 statistical software (R development core team, 2008) developed by Kang (1993) was used to calculate stability variances and YS_i values (which is determined by sum adjusted yield rank (Y) and stability rating (S) for each genotype). Genotypes having values for a character greater than mean YS_i and non-significant stability variance were judged as stable.

RESULTS AND DISCUSSION

Mean squares for crops were highly significant ($P < 0.01$) for all the characters except vigor rate 1 which was significant at $P < 0.05$, and stalk diameter was not significant (Table 1). Highly significant mean squares ($P < 0.01$) for the contrast of plant crop 2011 versus plant crop 2012 were recorded for growth1, growth2, vigor rate 1, cane length, number of internodes, purity, and number of millable canes, while significant ($P < 0.05$) for tillers2, and internode length, and non-significant for tiller1, vigor rate1, stalk diameter, brix, Pol, recovery, and cane yield. Mean squares of the contrast of plant crops versus ratoon were significant for all the characters except stalk diameter. Looking at the overall contribution of the contrasts mean

142 squares to the crops it was noted that plant crops versus ratoon contributed a substantial
143 amount to the crops mean squares.

144 Genotype mean squares were highly significant ($P < 0.01$) for all characters. In the
145 same way genotype x crop mean squares were highly significant ($P < 0.01$) for the characters
146 under study. Gilbert *et al* (2006) also found significant cultivar, environment, and time of
147 harvest interactions for sugar, cane yield, and sugar yield characters. Ramburan *et al* (2012)
148 reported that $G \times E$ interaction accounted for more variation than the main effect of genotype
149 in sugarcane. In our study the order of importance of sources of variation was crop,
150 genotypes and genotype x crop.

151 According to Kang and Martin (1987), theoretically, if the interactions are found non-
152 significant, performance of genotypes can be predicted from only one environment. However,
153 practicing breeders/geneticists are aware that this situation rarely occurs in typical
154 performance trials that examine large numbers of genotypes across many environments.
155 Shukla (1972) presented formulae for removing heterogeneity from the $G \times E$ interaction
156 variance, and partitioning the remainder of the $G \times E$ interaction variance and assigning it to
157 each genotype (parameter estimate). In our study, since the interaction component was
158 significant, this method was used for calculating stability analysis.

159 **Table 1. Analyses of mean squares, for 16 characters of 28 sugarcane genotypes analyzed in 2 plant-cane and one first-ratoon crop cycle.**

Source of variation	DF	Till1(number)	Till2(number)	VR1(ratio)	Gr1(cm)	Gr2(cm)	VR2(ratio)	Clength(cm)	Ninter(number)	IntLen(cm)	Cdia(cm)	Brix(%)	Pol(%)	Purity(%)	Recovery(%)	Mcanes(number)	Cyield(tonnes/ha)
Crops	2	531199.40**	1214753.69*	155.21**	123398.16*	104537.64*	4.54*	77399.40**	131.65**	188.08**	0.005	63.09**	122.53*	632.15**	78.83**	31473.91**	21695.05**
Reps within crops	6	30.67	1323.19	1.09	59.95	166.70	0.18	291.88	0.33	3.69	0.010	1.45	4.14	64.15	3.60	114.46	661.83
PC 2011 vs. PC 2012	1	2273.36	10465.93*	3.33	73246.20**	16656.31**	3.03**	9169.66**	113.31**	14.08*	0.003	2.41	0.38	144.10**	1.40	23347.50**	174.05
Plant crops vs. ratoon	1	1060125.45*	2419041.45*	307.10**	173550.11*	192418.98*	6.05**	145629.14*	150.00**	362.07**	0.006	123.77*	244.68*	1120.21*	156.26**	39600.31**	43216.05**
Genotype ^a	27	7770.20**	14322.62**	3.30**	1580.37**	4963.01**	1.43**	1484.59**	11.11**	15.13**	0.021**	2.43**	4.51**	45.70**	3.31**	1078.40**	296.74**
Genotype x Crop	54	3425.92**	5175.41**	3.56**	649.53**	1372.15**	1.02**	582.31**	5.24**	6.83**	0.013**	2.15**	2.89**	32.37**	2.05**	462.49*	276.84**
Error	162	880.46	2007.62	1.08	116.73	280.22	0.34	283.75	1.52	2.80	0.004	0.72	1.12	12.23	0.81	289.63	104.77
Total	251																
CV		18.90	15.82	24.66	10.47	9.06	20.43	9.87	8.48	12.31	6.62	4.38	6.58	4.22	8.90	21.54	15.31
Mean		156.98	283.20	4.21	103.18	184.78	2.87	170.72	14.53	13.58	0.93	19.37	16.70	82.82	10.09	78.98	66.83

160
161 **, * = Significant at P=0.01 and P=0.05, respectively.

162 Till= Tillers. Gr= Growth. VR= Vigor Rate. Clength= Cane Length. Ninter= Number of internodes. IntLen= Internode Length. Cdia= Cand diameter. Mcanes= Number of Millable Canes. Cyield= Cane Yield.

163 ^aMean squares for VR1, Cdia, Brix, Pol, Purity, Recovery, and Yield were tested against pooled error.

SHUKLA'S STABILITY VARIANCES (Σ_i^2) AND YIELD STABILITY (YS_i 'S)

Tillers and Growth

For tiller1, genotypes BF-162-166, MS-2003-CP-301, S-87-US-1767, MS-2003-CR2-131, and MS-2003-CR8-407, showed lower stability variances with mean values of 185.00, 167.83, 157.00, 147.33, and 192.50, respectively, while their yield stability values ranged from 12 to 30 which were above mean YS_i (11.68) (Table 2). For tiller2, genotypes MS-2003-CP-275, MS-2003-CP-378, CoJ-76, MS-2003-CR5-245, and MS-2003-CR8-407 showed mean values in the range of 312.5 to 342.06, lower and non-significant stability variances and higher values than mean YS_i (12.29). Vigor rate1 exhibited lower values of stability variances for genotypes MS-2003-CP-279, MS-2003-CP-300, MS-2003-CR1-50, MS-2003-CR2-129, and MS-2003-CR7-243, and means in the range of 4.13 to 5.13. YS_i values for vigor rate1 ranged from 13 to 29 which were above the mean YS_i .

The range of means for growth1 according to lower stability variances and higher values of YS_i values than mean was noted as 100.90 to 117.50 for genotypes MS-2003-CP-154, S-87-US-1767, CoJ-76, MS-2003-CR8-407, and MS-2003-CR9-451. For growth2 genotypes S-9883-CSSG-1139, MS-2003-CR1-50, MS-2003-CR2-131, MS-2003-CR5-245, and, MS-2003-CR9-451, showed non-significant stability variances and YS_i values in the range of 18 to 25 which were higher than means YS_i value (11.14). Genotypes MS-2003-CP-275, S-9883-CSSG-1139, MS-2003-CR1-50, MS-2003-CR2-129, and MS-2003-CR5-245, displayed lower and non-significant stability variances for vigor rate 1. Their YS_i values ranged from 20 to 25 and were higher than mean YS_i value (13.54).

Cane Characters

In stalklength, genotypes S-87-US-2787, CPF-225, S-9883-CSSG-1139, MS-2003-CR1-79, and MS-2003-CR7-243 exhibited lower stability with a range of 18 to 30 for YS_i values which were greater than mean YS_i value (13.39). Their means fell between 171.65 and 191.03. For number of internodes, genotypes MS-2003-CP-209, CPF-225, S-9883-CSSG-1139, MS-2003-CR5-245, and Mardan 93, had non-significant stability variances with a range of YS_i values between 12 and 26 and means ranging from 14.28 to 15.89. For internode length, genotypes MS-2003-CP-275, MS-2003-CP-279, CoJ-76, S-9883-CSSG-1139, and MS-2003-CR2-129 showed lower and non-significant variances with higher YS_i values and means ranging from 13.41 to 15.33.

Stalkdiameters of genotypes S-9883-CSSG-1139, MS-2003-CR2-129, MS-2003-CR2-132, MS-2003-CR7-243, and CP 77/400 gave non-significant stability variances with a mean range of 0.93 to 1.02 and YS_i values between 12 and 28.

Quality Characters

Comment [K2]: Symbols are not meanfull in a title!

Comment [K3]: Please, include discussion in this subsection.

Genotypes MS-2003-CP-300, MS-2003-CP-378, CPF-225, MS-2003-CR8-407, and CP 77/400 manifested lower and non-significant values for stability parameter and high values for yield stability statistic which ranged from 14 to 27. These genotypes showed mean values in the range of 19.42 to 20.03. For Pol, genotypes MS-2003-CP-301, CPF-225, MS-2003-CR1-51, MS-2003-CR5-245, and MS-2003-CR8-407 displayed lower values for stability variances and were non-significant. Their means ranged from 16.13 to 17.07 and YS_i values from 17 to 28.

For purity, genotypes BF-162-166, CPF-225, S-9883-CSSG-1139, MS-2003-CR1-50, and MS-2003-CR5-245 showed non-significant values for stability statistic and higher values for YS_i than mean YS_i . Means for these genotypes ranged from 82.52 to 85.01. For recovery, genotypes MS-2003-CP-301, CPF-225, MS-2003-CR1-51, MS-2003-CR7-243, and MS-2003-CR8-407 showed non-significant values with higher YS_i values ranging from 14 to 27.

Millable canes and cane yield

Genotypes MS-2003-CP-209, CoJ-76, MS-2003-CR2-131, MS-2003-CR5-245, and MS-2003-CR8-407 manifested lower and non-significant stability variances for millable canes. Their means ranged from 81.72 to 99.31 with YS_i in the range of 19 to 30. For cane yield, MS-2003-CP-209, MS-2003-CP-275, CoJ-76, MS-2003-CR2-131, and MS-2003-CR5-245, showed values for stability parameter which were non-significant. Mean cane yield for these genotypes ranged from 67.22 to 75.44 with YS_i values greater than mean YS_i value (12.00).

Shukla's stability variances and Y_{si} (Table 2) showed that no single genotype was stable across all the characters. Similar results have been reported by Tahir *et al.* (2013) wherein a pronounced effect of environments was found and no single genotype stood out stable for all the characters across all environments. However, genotype MS-2003-CR8-407 showed stability for tiller1, tiller2, growth1, brix, Pol, recovery, and millable stalks. MS-2003-CR5-245 was relatively stable as it showed good stability for 6 characters including growth2, vigor rate2, number of internodes, Pol, millable canes and cane yield. Genotypes MS-2003-CP-209, MS-2003-CP-275, CoJ-76, MS-2003-CR2-131, and MS-2003-CR5-245 showed stability for cane yield.

Stability across different environments is an important attribute to consider in varietal breeding program. Those genotypes will be of interest, which are stable for most of yield characters. These statistics should be a regular part of the breeding program at Sugar Crops Research Institute, to help find out most stable genotypes.

????????

CONCLUSIONS

Analysis of variance for the characters under study showed significant differences among sugarcane genotypes for various sources of variation, including genotype x crop

Comment [K4]: Please, can you explain the results? By this way, it look likes a technical report!

Comment [K5]: I miss a serious discussion in this paper.

235 interactions. While no genotypes were stable for all characters studied, two genotypes were
236 stable for most characters. MS-2003-CR8-407 was stable for sugar-related traits, and MS-
237 2003-CR5-245 was stable for both a sugar-related trait (Pol) and cane yield. Therefore,
238 although genotype x environment is a significant factor to consider for sugarcane production
239 and cultivar development in Pakistan, it is interesting to note that at least for some traits, we
240 found two of 28 genotypes that were stable across two plant-cane environments and one first-
241 ratoon crop. Since these results show that for the overwhelming percentage of genotypes
242 tested for interaction with crop cycle, we should continue using stability analysis as an
243 essential tool for identifying genotypes that have good performance in both the plant-cane
244 and first-ratoon crops. This also suggests that we should conduct further studies to test the
245 importance of continuing selection to the second-ratoon crop and beyond. Based on the
246 extremely low percentage of genotypes that were stable across all three environments, we
247 strongly discourage against making selections based on plant-cane results only.

248 **Table 2.Means, Shukla's stability variance (σ_i^2), and yield stability statistic (YS_i) of 28 genotypes for 16 traits analyzed in 2 plant-cane and 1 first-ratoon crop**
249 **cycle.**

Genotype		Till1(number)			Till2(number)			VR1(ratio)			Gr1(cm)		
S.No.		Means	σ_i^2	YS_i	Mean	σ_i^2	YS_i	Mean	σ_i^2	YS_i	Mean	σ_i^2	YS_i
1.	MS-2003-CP-154	133.83	2046.88	5	219.50	8177.92 *	-5	2.86	2.73	0	100.90	102.83	12 +
2.	BF-162-166	185.00	1320.69	23 +	331.17	3103.78	27 +	4.87	4.92 *	22 +	101.70	581.45 **	5
3.	MS-2003-CP-209	177.50	3889.90 *	17 +	294.67	5597.30	18 +	3.91	0.25	5	81.70	710.80 **	-9
4.	MS-2003-CP-275	192.67	2839.84 *	23 +	312.50	192.29	22 +	4.00	1.49	8	108.53	511.86 *	17 +
5.	MS-2003-CP-279	174.83	5310.77 **	12 +	307.67	9611.96 **	12	4.43	1.42	20 +	98.10	134.52	8
6.	MS-2003-CP-300	187.17	3271.21 *	20 +	331.83	10256.17 **	20 +	4.82	2.35	23 +	95.13	393.16 *	3
7.	MS-2003-CP-301	167.83	[†] 67.60	18 +	297.17	7716.77 *	15 +	4.31	7.15 **	10	94.23	926.60 **	-4
8.	MS-2003-CP-377	147.17	1021.35	11	269.67	649.46	10	4.08	1.38	11	79.97	655.61 **	-10
9.	MS-2003-CP-378	189.33	4013.82 *	21 +	310.83	2199.72	21 +	4.05	0.13	9	93.83	271.44	3
10.	S-87-US-1767	147.33	-31.45	12 +	231.83	62.08	2	2.81	0.40	-1	117.50	336.58	28 +
11.	S-87-US-2787	113.33	393.83	1	231.00	9389.11 *	-3	3.92	11.18 **	-1	100.73	101.71	11
12.	CoJ-76	195.67	12263.74 **	21 +	328.00	2837.63	25 +	4.41	7.49 **	11	112.53	391.55 *	22 +
13.	CPF-225	194.17	4058.51 *	24 +	351.50	6207.32 *	26 +	5.24	18.89 **	22 +	94.87	86.76	6
14.	S-9883-CSSG-1139	144.50	758.33	10	267.00	918.93	9	4.08	0.04	11	98.97	306.12	10
15.	MS-2003-CR1-50	97.83	4663.26 **	-10	221.83	7273.24 *	-4	4.13	0.21	13 +	108.33	101.56	20 +
16.	MS-2003-CR1-51	129.00	1071.00	3	237.00	6465.61 *	1	3.60	3.22	3	82.40	311.03	0

Genotype		Till1(number)			Till2(number)			VR1(ratio)			Gr1(cm)		
S.No.		Means	σ_i^2	YS _i	Mean	σ_i^2	YS _i	Mean	σ_i^2	YS _i	Mean	σ_i^2	YS _i
17.	MS-2003-CR1-79	136.50	5182.61 **	0	282.00	11099.63 **	4	4.85	2.67	25 +	107.00	553.65 **	11
18.	MS-2003-CR2-129	134.50	2157.20	7	284.00	4436.48	15 +	4.98	0.72	27 +	123.30	1135.25 **	22 +
19.	MS-2003-CR2-131	157.00	-34.65	15 +	290.67	6013.87	17 +	4.45	5.57 **	13 +	110.73	836.22 **	16 +
20.	MS-2003-CR2-132	131.50	1411.35	4	234.00	5234.22	4	3.42	2.19	2	109.43	155.18	22 +
21.	MS-2003-CR2-155	109.17	661.27	-1	233.17	1107.08	3	4.13	4.21 *	9	86.37	-12.17	2
22.	MS-2003-CR5-245	192.50	6285.40 **	18 +	318.33	2886.12	24 +	4.19	1.63	15 +	106.30	122.41	17 +
23.	MS-2003-CR6-295	159.67	2492.97	16 +	276.83	7581.72 *	7	3.91	1.75	5	98.73	442.14 *	5
24.	MS-2003-CR7-243	174.33	3038.09 *	15 +	328.17	6459.48 *	22 +	5.13	2.55	29 +	110.53	642.15 **	15 +
25.	MS-2003-CR8-407	196.60	1191.40	30 +	342.06	903.75	29 +	4.85	1.60	24 +	114.34	298.23	27 +
26.	MS-2003-CR9-451	163.73	2289.62	17 +	285.61	7269.48 *	12	4.06	1.44	10	103.66	227.26	16 +
27.	CP 77/400	139.33	13631.15 **	1	249.94	3573.57	7	3.69	9.04 **	-4	143.01	3981.60 **	23 +
28.	Mardan 93	123.33	10795.92 **	-6	261.72	7686.09 *	4	4.61	3.13	22 +	106.32	3881.30 **	10
Means		156.98		11.68	283.20		12.29	4.21		12.25	103.18		11.00

***, = Significant at P=0.1, 0.05 and 0.01, respectively, and also indicate that genotype was judged to be unstable.

*= Selected genotypes

Till= Tillers. Gr= Growth. VR= Vigor Rate

†negative values for stability variance may be taken as zero.

255

256 **Table 2.(Contd.) Genotypes means, Shukla's stability variance (σ_i^2) and yield stability statistic (YS).**

S.No.	Genotypes	Gr2(cm)			VR2(ratio)			Clength(cm)			Ninter(cm)		
		Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i
1.	MS-2003-CP-154	171.40	813.68	4	2.47	0.72	3	166.30	117.68	9	14.11	4.06	10
2.	BF-162-166	176.10	1817.79 **	-1	2.74	1.09 *	8	171.56	583.06	17 +	13.22	1.21	3
3.	MS-2003-CP-209	153.67	2460.35 **	-7	2.69	0.91	10	171.28	1192.56 *	12	14.28	0.89	12 +
4.	MS-2003-CP-275	188.73	124.53	20 +	2.97	0.00	20 +	176.89	847.98	20 +	13.05	-0.18	0
5.	MS-2003-CP-279	180.43	469.60	12 +	2.96	0.74	19 +	159.17	360.60	5	12.94	5.68 *	-5
6.	MS-2003-CP-300	168.00	16.70	3	2.65	0.32	7	166.00	90.65	8	14.33	1.01	13 +
7.	MS-2003-CP-301	176.17	1261.23 *	4	2.73	0.30	11	147.17	796.11	-1	13.05	3.00	0
8.	MS-2003-CP-377	141.40	294.95	-2	2.62	2.37 **	-2	152.20	-21.90	1	13.55	0.66	5
9.	MS-2003-CP-378	177.63	1341.17 **	1	2.77	0.01	13	154.26	444.39	2	16.61	2.22	30 +
10.	S-87-US-1767	193.63	904.61 *	17 +	2.65	0.27	8	170.61	192.29	13	13.17	10.96 **	-6
11.	S-87-US-2787	173.83	814.04	6	2.48	0.89	4	174.57	268.65	19 +	15.44	11.37 **	16 +
12.	CoJ-76	183.03	1003.53 *	9	2.44	0.25	2	166.30	53.41	9	14.00	1.57	8
13.	CPF-225	179.57	114.84	10	2.94	0.47	18 +	183.44	193.17	23 +	15.89	0.23	26 +
14.	S-9883-CSSG-1139	187.53	280.02	19 +	3.10	0.09	22 +	171.65	203.79	18 +	14.89	0.03	19 +
15.	MS-2003-CR1-50	195.77	260.05	22 +	2.98	-0.04	21 +	167.45	100.87	12	15.28	5.28 *	19 +


S.No.	Genotypes	Gr2(cm)			VR2(ratio)			Clength(cm)			Ninter(cm)		
		Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i
16.	MS-2003-CR1-51	145.23	602.39	-1	2.27	0.22	0	158.89	2873.35 **	-4	13.61	1.74	7
17.	MS-2003-CR1-79	184.87	1909.81 **	8	2.68	0.71	9	160.67	210.67	6	14.78	7.92 **	10
18.	MS-2003-CR2-129	215.70	1540.83 **	22 +	3.21	0.03	24 +	188.22	248.69	27 +	15.00	4.72 *	17 +
19.	MS-2003-CR2-131	198.57	175.72	25 +	3.15	0.47	23 +	187.72	428.62	26 +	16.67	11.82 **	23 +
20.	MS-2003-CR2-132	185.03	646.63	17 +	2.60	0.18	5	167.17	329.80	11	14.06	7.25 **	1
21.	MS-2003-CR2-155	145.27	1889.53 **	-8	2.10	1.72 **	-9	147.22	1088.99 *	-4	16.39	18.61 **	20 +
22.	MS-2003-CR5-245	198.47	370.35	24 +	3.24	0.20	25 +	189.33	2025.52 **	21 +	15.89	0.94	26 +
23.	MS-2003-CR6-295	180.00	2369.12 **	3	2.91	0.58	17 +	177.33	1072.61 *	18 +	15.22	6.54 *	18 +
24.	MS-2003-CR7-243	210.90	1353.00 **	19 +	3.25	0.96	26 +	191.03	280.90	30 +	14.61	12.35 **	9
25.	MS-2003-CR8-407	214.93	1322.16 *	25 +	3.45	0.62	28 +	189.22	471.49	28 +	14.21	7.97 **	3
26.	MS-2003-CR9-451	187.03	617.93	18 +	2.88	0.89	16 +	177.06	346.63	21 +	14.35	9.28 **	6
27.	CP 77/400	249.58	11051.87 **	23 +	3.67	13.14 **	21 +	162.42	180.91	7	13.22	9.14 **	-4
28.	Mardan 93	211.42	2594.15 **	20 +	3.68	0.43	30 +	185.13	1322.98 *	21 +	14.89	0.55	20 +
	Means	184.78		11.14	2.87		13.54	170.72		13.39	14.53		10.93

**, = Significant at P=0.1, 0.05 and 0.01, respectively, and also indicate that genotype was judged to be unstable.

*= Selected genotypes

Gr= Growth. VR= Vigor Rate. Clength= Cane Length. Ninter= Number of internodes.

†negative values for stability variance may be taken as zero



262

263 **Table 2 (contd). Genotypes means, Shukla's stability variance (σ_i^2) and yield stability statistic (YS_i).**

264

S.No.	Genotypes	IntLen(cm)			Cdia(cm)			Brix(%)			Pol(%)		
		Means	σ_i^2	YS_i	Means	σ_i^2	YS_i	Means	σ_i^2	YS_i	Means	σ_i^2	YS_i
1.	MS-2003-CP-154	12.98	0.88	8	1.03	0.02 **	23 +	20.57	4.58 **	22 +	17.90	6.40 **	23 +
2.	BF-162-166	15.25	1.78	27 +	0.89	0.00	5	19.83	1.68	25 +	16.41	3.30	22 +
3.	MS-2003-CP-209	13.19	1.92	11	0.88	0.03 **	-8	18.23	0.93	-1	15.26	0.20	4
4.	MS-2003-CP-275	15.33	0.06	28 +	0.90	0.01	6	19.54	1.70	19 +	15.40	0.31	5
5.	MS-2003-CP-279	13.61	0.20	17 +	0.93	0.01	16 +	19.16	1.17	8	15.65	-0.04	7
6.	MS-2003-CP-300	12.44	0.83	3	0.92	0.00	13 +	19.72	0.65	24 +	16.42	8.70 **	15 +
7.	MS-2003-CP-301	12.78	1.91	7	0.93	0.00	14 +	19.89	1.63	26 +	16.31	0.62	21 +
8.	MS-2003-CP-377	12.22	1.86	1	0.98	0.01	24 +	19.47	1.97	17 +	15.80	3.06	9
9.	MS-2003-CP-378	11.14	7.01	0	0.91	0.01	10	19.42	0.23	14 +	15.94	0.01	12
10.	S-87-US-1767	15.11	4.89	26 +	0.93	0.08 **	6	19.45	1.72	15 +	15.87	0.68	11
11.	S-87-US-2787	12.55	0.28	5	1.02	0.01 *	24 +	19.59	4.72 **	12	16.84	5.90 **	17 +
12.	CoJ-76	13.41	0.87	13 +	0.94	0.02 **	13 +	19.68	6.70 **	15 +	16.98	10.37 **	19 +
13.	CPF-225	13.83	3.21	19 +	0.90	0.00	6	20.03	0.10	27 +	17.07	-0.06	28 +
14.	S-9883-CSSG-1139	13.89	1.53	20 +	0.93	0.00	20 +	19.46	1.75	16 +	16.31	2.01	20 +

S.No.	Genotypes	IntLen(cm)			Cdia(cm)			Brix(%)			Pol(%)		
		Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i
15.	MS-2003-CR1-50	13.16	13.61 **	2	0.97	0.01	23 +	19.23	1.16	9	15.95	2.18	13 +
16.	MS-2003-CR1-51	13.08	38.78 **	1	0.90	0.05 **	0	19.59	2.97 *	16 +	16.13	0.11	17 +
17.	MS-2003-CR1-79	13.33	9.26 *	8	0.90	0.01	8	18.60	2.74 *	-3	15.48	0.04	6
18.	MS-2003-CR2-129	14.75	1.05	23 +	0.96	0.00	22 +	18.93	0.62	4	15.20	0.57	1
19.	MS-2003-CR2-131	12.64	1.10	6	0.91	0.01	10	18.97	5.19 **	-3	15.25	3.61 *	-1
20.	MS-2003-CR2-132	13.50	6.82	14 +	0.93	0.00	16 +	19.06	0.29	7	14.97	0.50	-1
21.	MS-2003-CR2-155	10.72	14.19 **	-10	0.88	0.01 *	-4	18.86	2.13	3	15.01	5.01 *	-4
22.	MS-2003-CR5-245	13.64	10.66 *	14 +	0.88	0.00	0	19.34	-0.03	11	16.23	0.13	19 +
23.	MS-2003-CR6-295	12.44	8.67 *	-1	1.01	0.00	27 +	18.99	2.92 *	2	16.14	7.95 **	10
24.	MS-2003-CR7-243	16.25	29.22 **	23 +	0.91	0.00	12 +	18.38	2.73 *	-4	15.72	0.95	8
25.	MS-2003-CR8-407	15.44	4.49	29 +	0.88	0.04 **	-8	19.53	1.11	18 +	15.95	0.11	13 +
26.	MS-2003-CR9-451	14.45	12.05 *	18 +	0.86	0.01	-1	19.27	3.61 **	2	15.81	7.40 **	2
27.	CP 77/400	14.93	9.34 *	21 +	1.02	0.00	28 +	19.61	0.59	22 +	16.77	2.49	24 +
28.	Mardan 93	14.18	4.88	21 +	1.00	0.01	26 +	20.16	4.77 **	21 +	17.15	8.19 **	21 +
	Means	13.58		12.64	0.93		11.82	19.38		12.29	16.07		12.18

***, = Significant at P=0.1, 0.05 and 0.01, respectively, and also indicate that genotype was judged to be unstable.

* = Selected genotypes

IntLen= Internode Length. Cdia= Cand diameter. Mcanes= Number of Millable Canes. Cyield= Cane Yield.

268 †negative values for stability variance may be taken as zero.

269 **Table 2 (contd). Genotypes means, Shukla’s stability variance (σ_i^2) and yield stability statistic (YS_i).**

S.No.	Genotypes	Purity(%)			Recovery(%)			Mcanes(number)			Cyield(tonnes/ha)		
		Means	σ_i^2	YS_i	Means	σ_i^2	YS_i	Means	σ_i^2	YS_i	Means	σ_i^2	YS_i
1.	MS-2003-CP-154	86.95	64.51 **	22 +	11.58	4.45 **	23 +	71.44	911.58 *	3	67.22	6.93	16 +
2.	BF-162-166	82.52	9.72	13 +	10.28	2.07	19 +	75.61	194.68	11	66.67	1.02	11
3.	MS-2003-CP-209	83.65	31.85	19 +	9.64	0.46	5	81.72	116.14	19 +	68.56	27.48	19 +
4.	MS-2003-CP-275	78.72	19.81	0	9.33	0.29	1	75.50	1247.48 *	6	67.78	-2.36	18 +
5.	MS-2003-CP-279	81.64	34.63	8	9.72	0.38	6	80.98	370.51	18 +	63.56	16.08	8
6.	MS-2003-CP-300	83.37	151.71 **	10	10.34	8.14 **	15 +	84.44	1223.28 *	17 +	67.56	1151.74 **	9
7.	MS-2003-CP-301	81.85	33.11	10	10.15	0.74	18 +	90.50	410.40	25 +	71.44	886.54 **	14 +
8.	MS-2003-CP-377	80.95	16.21	5	9.77	1.94	8	64.43	410.61	1	56.22	91.95	-1
9.	MS-2003-CP-378	82.01	16.81	11	9.96	0.26	12	76.11	115.22	12	57.89	473.57 *	-4
10.	S-87-US-1767	81.55	7.03	6	9.86	0.22	10	71.78	76.00	8	74.11	772.09 **	18 +
11.	S-87-US-2787	85.72	18.00	28 +	10.82	3.45 *	22 +	57.33	235.14	-1	59.34	360.02 *	0
12.	CoJ-76	86.12	24.94	29 +	10.93	6.21 **	20 +	87.42	52.31	23 +	75.44	25.65	29 +
13.	CPF-225	85.01	4.40	23 +	10.92	0.00	27 +	87.28	933.41 *	18 +	66.78	81.11	13 +
14.	S-9883-CSSG-1139	83.83	4.49	20 +	10.31	1.08	21 +	71.00	290.21	6	63.11	85.18	6
15.	MS-2003-CR1-50	82.69	8.65	14 +	10.02	1.46	13 +	70.92	4.81	5	64.45	334.68 *	5

S.No.	Genotypes	Purity(%)			Recovery(%)			Mcanes(number)			Cyield(tonnes/ha)		
		Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i	Means	σ_i^2	YS _i
16.	MS-2003-CR1-51	82.16	24.46	12	10.09	-0.05	15 +	60.37	761.88	0	69.78	43.72	21 +
17.	MS-2003-CR1-79	83.30	66.41 **	9	9.73	0.52	7	67.33	1436.00 **	-5	69.33	497.82 **	12
18.	MS-2003-CR2-129	80.26	14.86	3	9.34	0.51	2	73.11	219.88	9	74.00	93.98	25 +
19.	MS-2003-CR2-131	80.43	40.30 *	0	9.38	2.09	3	94.39	171.64	28 +	65.00	17.80	10
20.	MS-2003-CR2-132	78.49	35.68	-1	9.03	0.96	-1	82.78	534.15	20 +	58.67	251.98	2
21.	MS-2003-CR2-155	79.62	47.82 *	-3	9.17	3.58 *	-4	67.56	280.29	4	66.67	40.64	11
22.	MS-2003-CR5-245	83.94	6.88	21 +	10.29	0.21	20 +	94.17	132.56	27 +	71.78	-5.47	23 +
23.	MS-2003-CR6-295	85.13	49.45 *	20 +	10.31	5.48 **	13 +	80.78	200.53	17 +	63.33	636.69 **	-1
24.	MS-2003-CR7-243	85.46	11.32	26 +	10.06	0.23	14 +	96.83	529.97	29 +	75.00	733.18 **	20 +
25.	MS-2003-CR8-407	81.56	14.14	7	9.92	0.15	11	99.31	134.30	30 +	75.56	564.86 **	22 +
26.	MS-2003-CR9-451	81.79	45.90 *	5	9.85	4.97 **	1	89.21	1175.19 *	20 +	58.00	303.07	1
27.	CP 77/400	85.36	54.14 *	21 +	10.75	2.46	24 +	77.89	337.63	13	72.67	225.44	24 +
28.	Mardan 93	84.86	49.90 *	18 +	10.94	5.29 **	21 +	80.64	444.83	16 +	61.33	36.70	5
	Means	82.82		12.71	10.09		12.36	78.96		13.54	66.83		12.00

**, = Significant at P=0.1, 0.05 and 0.01, respectively, and also indicate that genotype was judged to be unstable.

*= Selected genotypes

Mcanes= Number of Millable Canes. Cyield= Cane Yield.

†negative values for stability variance may be taken as zero.

274 **AUTHORS' CONTRIBUTIONS**
 275 1. Mohammad Tahir: Designed, and laid out the experiment; compiled the study results, followed by
 276 statistical analyses; wrote the first draft.
 277 2. Dr Iftikhar Hussain Khalil: Supervised the research project/relevant literature search.
 278 3. DrPerH.Mccord: Reviewed the first draft.
 279 4. Barry Glaz: Critically reviewed and contributed to writing of the manuscript.

280 **REFERENCES CITED**
 281

- 282 1. Gilbert RA, JM Shine Jr, JD Miller, RW Rice, CR Rainbolt. The effect of genotype, environment,
 283 and time of harvest on sugarcane yields in Florida, USA. *Field Crops Research*. 2006; 95: 156-
 284 170
 285 2. Jackson, PA, DM Hogarth. Genotype × environment interactions in sugarcane. I. Patterns on
 286 response across sites and crop-years in North Queensland. *Aust. J. Agric. Res.* 1992; 43: 1447-
 287 1460.
 288 3. Kang MS. Simultaneous selection for yield and stability: Consequences for growers. 1993; *Agron.*
 289 *J.* 85:754-757.
 290 4. Kang MS, FA Martin. A review of important aspects of Genotype-Environment Interactions and
 291 practical suggestions for sugarcane breeders. 1987; *J. Am. Soc. Sugarcane Tech.* 7: 36.
 292 5. Tahir, M., R. Gul, H. Rahman, and M. Khalid. 2013; Genetic divergence in sugarcane genotypes.
 293 *Am. J. Exp. Agric.* 3(1): 102-109
 294 6. R Development Core Team. R: A language and environment for statistical computing. R
 295 Foundation for Statistical Computing, Vienna, Austria. 2008; ISBN 3-900051-07-0, URL
 296 <http://www.R-project.org>.
 297 7. Ramburan S, M Zhou, M Labuschagne. Integrating empirical and analytical approaches to
 298 investigate genotype x environment interactions in sugarcane. *Crop Sci.* 2012; 52: 2153-2165.
 299 8. SAS Institute. SAS system for Windows Release 9.1. 2003; SAS Inst., Cary, NC.
 300 9. Shukla GR. Some statistical aspects of partitioning genotype-environmental components of
 301 variability. *Heredity*. 1972; 29:237-245.
 302 10. Singh HN. Stability parameters for yield and quality characters in sugarcane. Paper presented to
 303 the 18th cong. ISSCT. 1983; 12pp. Abstracted in sugarcane *Int. J. Cane. Agri. No.* 1983; 1: 26.
 304 11. Bissessur, D, RAE.Tinley-Bassett, LYCChong, MHRJulien. Family x environment and genotype x
 305 environment interactions for sugarcane across two contrasting marginal environments in
 306 Mauritius. 2000; *Expl. Agric.* 36: 101-114.
 307 12. Tahir M, H Rahman, A Ali, S Anwar, M Khalid. Assessment of genotype x environment interaction
 308 and stability of promising sugarcane genotypes for different agronomic characters in Peshawar
 309 valley. 2013; *Am. J. Exp. Agric.* 3(1): 142-151.
 310 13. Zhou M, SV Joshi, T Maritz. Trends and implications of genotype by environment interactions in
 311 South African sugarcane breeding. *J. Crop Impr.* 2012; 26: 163-176.