

REVIEW OF AQUACULTURAL PRODUCTION AND MANAGEMENT IN NIGERIA

¹Ozigbo Emmanuel, ¹Anyadike Chinenye, ²Adegbite Oluwatobi and ³Kolawole Peter¹Department of Agricultural and Bioresources Engineering, University of Nigeria Nsukka.²Department of Biological Sciences, Covenant University, Ota, Nigeria. ³Post Harvest Unit, International Institute of Tropical Agriculture, Ibadan, Nigeria.

ABSTRACT

Fish is an important source of food, income, employment, and recreation for people around the world and it is a very important source of animal protein for both man and livestock in developed and developing countries. In Nigeria, the current demand for fish is about four times the level of local production. Humans consume approximately 80 percent of the catch as food. The remaining 20 percent goes into the manufacturing of products such as fish oil, fertilizers, and animal food. Fish farming named aquaculture is an integral part of agriculture which was found to have the capacity to increase the country's GDP and can solve the unemployment problem for our teeming youths if adequately managed. Therefore, this paper reviewed the status of Aquacultural production in Nigeria, problems and prospects of Aquacultural development in the country, different fish species that can adequately thrive in Nigerian pond and its preparation and finally showed some first-rate management practices that can boost aquacultural production in Nigeria.

KEYWORDS: Aquaculture, Development, Management, Production, Stocking

INTRODUCTION

The demand for fish in Nigeria mostly outstrips the local production. Nigeria is the largest fish consumer in Africa and among the largest fish consumers in the world with over 1.5 millions tons of fish consumed annually. Yet, Nigeria imports over 900,000 metric tons of fish while its domestic catch is estimated at 450,000 metric tons/year (Ozigbo *et al.*, 2013). The table 1 below showed the projected human population figures, the fish demand and supply in Nigeria from 2000 – 2015. The statistic available indicates that the growth in fish production is due to increased activities of Aquaculture and the need for aquaculture arose from the decrease in supply from ocean fisheries as a result of over-fishing, habitat destruction and pollutions (Adedeji and Okocha, 2011). Aquaculture is farming of aquatic organisms in fresh, brackish or salt water. A wide variety of aquatic organisms are produced through aquaculture. These include: fishes, crustaceans, molluscs, algae, and aquatic plants. Unlike capture fisheries, aquaculture requires deliberate human intervention in the organisms' productivity which results in yields that exceed those from the natural environment alone. Such interventions are stocking water with seed (fingerlings), fertilizing the water, feeding the organisms, and maintaining water quality (Green, 2008). Objectives of aquaculture are as follows: production of protein rich, nutritive, palatable and easily digestible human food benefiting the whole society through plentiful food supplies at low or reasonable cost; providing new species and strengthening stocks of existing fish in natural and man-made water-bodies through artificial recruitment and transplantation; production of sport-fish and support to recreational fishing; production of bait-fish for commercial and sport fishery; production of ornamental fish for aesthetic appeal, recycling of organic waste of human and livestock origin, land and aquatic resource utilization: this constitutes the macro-economic point of view benefiting the whole society. It involves (a)

Comment [SM1]: Fisheries and Aquaculture

Comment [SM2]: Organisms and plants

maximum resource allocation to aquaculture and its optimal utilization; (b) increasing standard of living by maximising profitability; and (c) creation of production surplus for export (earning foreign exchange especially important to most developing countries) (Jhingran, 2001). The objective of study is to enumerate the status of Aquacultural production in Nigeria, highlight the problems and prospects of Aquacultural development in the country and provide good production and management practices that can boost aquacultural production in Nigeria.

Table 1: Projected Human Population, Fish Demand and Supply in Nigeria (2000 – 2015)

Year	Projected Population (Million)	Projected Fish Demand (Tonnes)	Projected Domestic Fish Supply (Tonnes)	Deficit (Tonnes)
2000	114.4	1,430,000.00	467,098.00	962,902.00
2001	117.6	1,470,000.00	480,163.60	984,836.40
2002	121.0	1,412,500.00	507,928.20	1,004,572.00
2003	124.4	1,555,000.00	522,627.10	1,063,082.60
2004	128.0	1,600,000.00	536,917.60	1,063,072.40
2005	131.5	1,643,750.00	552,433.10	1,091,317.00
2006	135.3	1,691,250.00	567,948.60	1,23,301.40
2007	139.1	1,732,750.00	583,872.40	1,154,873.00
2008	143.0	1,782,300.00	600,612.80	1,186,887.20
2009	147.1	1,838,750.00	617,353.20	1,221,397.00
2010	151.2	1,810,000.00	634,500.20	1,255,440.00
2011	155.5	1,943,750.00	652,606.60	1,291,143.00
2012	160.0	2,000,000.00	689,958.00	1,328,508.00
2013	164.0	2,113,750.00	709,683.10	1,365,042.00
2014	169.1	2,175,000.00	730,248.00	1,404,067.10
2015	174.0	2,055,000.00	671,492.30	1,444,752.10

Source: (FDF, 2008).

STATUS OF AQUACULTURE IN NIGERIA

Aquaculture was developed more than 2,000 years ago in countries such as China, Italy, and Egypt. Not long after, aquacultural practices in Europe, China, and Japan commonly involved stocking wild-caught fingerlings — for example, carp fingerlings (juvenile fish) captured from rivers — in ponds or other bodies of water for further growth (Green, 2008). The World leading aquaculture producers today are Asia and America. The United States of America (U.S.A) aquaculture production rose more than 400 percent between 1980 and 2000 (Jhingran, 2001). The substantial increase in aquacultural production in countries like U.S.A is as result of aquacultural mechanization which has led to increased productivity, labour efficiency and improved product quality. Unlike Asia, Africa has little aquaculture tradition and has been affected by a number of external problems that have prevented proper management and development despite the investment. A number of countries in sub-Saharan Africa are characterized by low agricultural production, poor management of resources, economic stagnation, persistent political instability, lack of technical know-how, increasing environmental damage, and severe poverty.

Table 2: Aquaculture Production (volume and value) in 2005 in different region of the world

Comment [SM3]: delete

Country/Region	Production volume (million tonnes)	Production volume (%)	Production value US\$	Production Value (%)
China	32.4	67.3	35.99	51.22
Rest of the Asia Pacific	10.7	22.3	20.6	29.3
Western Europe	2	4.2	5.42	7.72
Latin America and the Caribbean	1.4	2.9	5.24	7.47
North America	0.6	1.3	1.3	1.86
North Africa	0.6	1.2	0.83	1.19
Central and Eastern Europe	0.3	0.6	0.67	0.91
Sub-Saharan Africa	0.1	0.2	0.25	0.36
World total	48.1	100	70.3	100

Source: (FAO, 2006)

Although aquaculture activities in Nigeria started about 50 years ago (Olagunju *et al.*, 2007), yet Nigeria has not been able to meet domestic production demand for the populace. According to Ekunwe and Emokaro (2009) statistics indicate that Nigeria is the largest African aquaculture producer, with production output of over 15,489 tonnes per annum, this is closely followed by Egypt with output of about 5,645 tonnes. Only five other countries: Zambia, Madagascar, Togo, Kenya and Sudan produce more than 1,000 tonnes each. This result shows that Africa in general is far behind in aquacultural production. In Nigeria, aquaculture development has been driven by social and economic objectives, such as nutrition improvement in rural areas, generation of supplementary income, diversification of income activities, and the creation of employment. This is especially true in rural communities, where opportunities for economic activities are limited. Only in recent years has aquaculture been viewed as an activity likely to meet national shortfalls in fish supplies, thereby reducing fish imports. The fisheries sector accounts for about 2% of national G.D.P, 40% of the animal protein intake and a substantial proportion of employment, especially in the rural areas; the sector is a principal source of livelihood for over three million people in Nigeria (Adedeji and Okocha, 2011). In terms of economic and trade, aquaculture makes a minor contribution to overall fish and protein supply and G.D.P. this can be attributed to emerging nature of the sector when compared with agriculture and fisheries which are important primary sectors. Agriculture contribution to G.D.P between 1999 and 2002 was 34% while Aquaculture contribution alone to G.D.P within the same period was 0.154%. In relation to meat production, capture fisheries and aquaculture contributed 475, 162 and 30,677 respectively to meat production. Culture fisheries contribution to fish supply was 6.06%. Per capita fish consumption between 2003 and 2004 was 7.3% per Kg per year as compared to the recommended rate of 12.0% per Kg per year (Adeogun *et al.*, 2008).

Nigeria has the natural resources (such as lands, rivers, streams, reservoirs and lakes; and human resource) and potentials to compete with the world leading aquacultural countries. In 1990, 82 million hectares out of Nigeria's total land area of about 91 million hectares were found to be arable, although only 42 percent of the cultivable area was farmed. Much of this land was farmed under the bush fallow system, whereby land is left idle for a period of time to allow natural regeneration of soil fertility. 18 million hectares were classified as

permanent pasture, but had the potential to support crops. Most of the 20 million hectares covered by forests and woodlands are believed to have agricultural potential (Helen, 1991). Also in terms of water resources, Nigeria has about 264 medium and large dams with a combined storage capacity of 33 billion cubic meters of water. Of these dams, 210 are owned by the Federal Government, 34 are owned by states, while 20 are owned by private organizations. The dams, which have about 350,000 hectares of irrigable land in their vicinities, are largely underutilized (Magdalene, 2013).

The Maximum Sustainable Yield of Nigerian sea was observed when Ikenweuwe (2013) stated that Nigeria has about 853km coastline and a maritime water of 210,900km² including the Exclusive Economic Zone (EEZ) (while Nwosu et al (2011) and FAO (2013) noted that the figures of coastline and marine water are about 900km, 217,313 km² and 960km, 309,120 km² respectively). The continental shelf is narrow, extending, for only about 15km in the western area and ranges from 60 – 80km in the eastern tip. This condition limits the trawlable grounds to 3200nm² of the 1147nm² continental shelf area. The inshore waters (0 – 50m) are characterized by a variety of small fish species varying from 25 to 50cm in total length. The most predominant is the *Pseudotolithus*. Estimated potential yield of the inshore waters is about 16620mt for finfish and between 3500 – 4020mt for shellfish resources, which are exploited by both the artisanal and industrial operators, the potential fisheries resources are estimated at about 9460mt, and consists of mostly tuna and tuna-like fishes. She further added that more than 90 percent of recorded fish production in Nigeria emanates from the artisanal fisheries sector (marine, brackish and inland or fresh water). The principal pelagic fishes in inshore artisanal catches are bonga (*Ethmalosa fimbriata*) and sardine (*Sardinella Maderensis*) while the most common semi-pelagic is the shad (*Ilisha Africana*). Maximum Sustainable Yield (MSY) for pelagic fishes at sea in Nigeria was put at 60,000 tonnes while Ajayi and Talabi (1984) summed up the yield of the fishery in coastal and brackish waters to be 70,000 – 90,000 tonnes. Although, an annual landing of 40,000 tonnes was projected for the fishery at sea during 1975 – 1980, far less catches were made.

Table 3: Nigeria fish supply by sector

Sector	2000	2001	2002	2003	2004	2005	2006	2007
Artisanal	418,069.00	433,537	450,965	446.203	434.830	513,537	53,332	504,227
Aquaculture	25,720	26,398	30,664	33,667	43,950	56.355	84,523	85.087
Industrial trawler	23,308.3	28,378	30,091	33,882	30,421	33,778	33,778	28,193
Imported Fish	557,884	681,152	663,180	663,152	611,152	646,484	646,484	737,666

Source (FDF, 2007)

In spite of these great potentials of natural resources and man-power availability to fish farming in Nigeria, the country is still currently unable to bridge the gap in the short fall between total domestic fish production and the total domestic demand. Meanwhile, millions of Nigerian youths are ready and equally available to work any time of the day. Kareen et al (2008) reported that the growth in general agricultural sector in Nigeria has fallen short of expectations. Value added per capita in agriculture has risen by less than 1% per year for the past 20

Comment [SM4]: delete

Comment [SM5]: delete

Comment [SM6]: These dams can be used for cage aquaculture and pen aquaculture.

Comment [SM7]: (MSY)

Comment [SM8]: *Pseudotolithus*

Comment [SM9]: Delete
It relates to capture fisheries. Paper is about aquaculture.

Comment [SM10]: Add "Nigerian coastline has huge potential for inland aquaculture and offshore cage culture."

123 years and food productions gains have not kept pace with population growth, resulting in rising food imports
124 and declining levels of national food self-sufficiency.

125

126 **Aquacultural Farm Systems and Techniques**

127 There are different systems and techniques used in aquaculture for fish production. We have three major
128 systems based on feeding methods; extensive, intensive and semi-intensive. In extensive system, the fish
129 feeding is based on local photosynthetic production (natural foods like phytoplanktons and zooplanktons). That
130 is, no supplementary feeding is required from the aquaculturalists. Intensive system is the one in which the
131 fish are fed with external food supply. Whereas, in semi-intensive, the fish are fed with supplementary feed in
132 support of the natural food supply.

133

134 There various techniques used are as follows: flow through systems, ponds, cages, tanks and recirculating
135 systems:

136 **A Flow-through System**, also known as a raceway, is an artificial channel used in aquaculture to culture
137 aquatic organisms. Raceway systems are among the earliest methods used for inland aquaculture. A raceway
138 usually consists of rectangular basins or canals constructed of concrete and equipped with an inlet and outlet.
139 A continuous water flow-through is maintained to provide the required level of water quality, which allows
140 animals to be cultured at higher densities within the raceway (Mirzoyan and Gross, 2010). Freshwater species
141 such as trout, catfish and tilapia are commonly cultured in raceways (Gupta and Acosta, 2004).

142 **Fish Cage aquaculture** refers to the rearing of aquatic species, within enclosures in natural waterways. The
143 fish cages are placed in lakes, bayous, ponds, rivers or oceans to contain and protect fish until they can be
144 harvested (Offshore, 2013). The method is also called "off-shore cultivation" when the cages are placed in the
145 sea. Fish are stocked in cages, artificially fed, and harvested when they reach market size. Open systems are
146 being implemented in a wide range of environments including freshwater rivers, brackish estuaries and coastal
147 marine regions. Floating mesh cages are anchored to the seafloor and vary in size depending on the scale of
148 operation and the species cultured.

149 **Fish Tanks:** fish farming can also be carried out in outdoor or indoor concrete or plastic tanks. Tanks can be
150 inform of small aquaria (glass or plastic) or large fibreglasses. Production tanks varies in size and shape,
151 however, round tanks between 5,000 to 10,000 liters are most commonly used (EAI, 1990). Tanks need to be
152 non corrosive therefore plastic or fiberglass is recommended. Smooth round tanks with a conical shaped
153 bottom are considered advantageous as this will assist with waste solids disposal during draining. Some
154 growers will use small cages within the tanks to aid in introducing new stock into the cage. Using this method,
155 growers can isolate and protect newly introduced fingerlings from cannibalism and disease. Alternatively,
156 nursery tanks can be incorporated into the system.

157 **A Fish Pond** is a controlled pond, artificial lake, or reservoir that is stocked with fish and is used in aquaculture
158 for fish farming, or is used for recreational fishing or for ornamental purposes. || A Pond is an intensive system,
159 though at times, it can be made semi-intensive system and can give high yield production provided that good

Comment [SM11]: delete

Comment [SM12]: delete

Comment [SM13]: delete

Comment [SM14]: delete

Comment [SM15]: Add "Mostly earthen ponds are used for culture of carps, tilapia, catfishes and seabass."

160 feeding and management procedures are adopted. According to FAO accessed online (2013), Fish pond
161 aquaculture production is currently very low, estimated at below, 10,000 mt/year from a total water surface
162 area of about 5,500 hectares. It has been estimated that if Nigeria is to be self sufficient in fish production
163 through aquaculture, a total of about 900,000 hectares of water surface area must be cultivated to produce a
164 minimum of about 900,000 mt of fish a year (i.e. estimating at a minimum production of 1 mt per hectare per
165 year).

166 **A Recirculating System** is essentially a closed system and involves fish tanks and filtration and water
167 treatment systems. The fish are housed within tanks and the water is exchanged continuously to guarantee
168 optimum growing conditions. Water is pumped into the tanks, through biological and mechanical filtration
169 systems and then returned into the tanks. Not all water is 100% exchanged however as it is difficult to ensure
170 that all waste products are converted or removed by the treatment process. Most culture systems recommend
171 at least 5% to 10% water exchange rate per day depending on stocking and feeding rates (EAI, 1990).
172 Recirculating systems occupy a very small area and allow the grower to stock fish at high densities and
173 produce high yields per unit area. Recirculating systems are very intensive and therefore require a high level in
174 management of stock, equipment and water quality. Recirculating systems provide a predictable and constant
175 environment for growing fish. Recirculation systems can be expensive to purchase and operate. For this
176 reason it is usually only economically viable to farm high value species in these systems.

Comment [SM16]: delete

Comment [SM17]: Recirculating Aquaculture System (RAS)

Comment [SM18]: RAS

Comment [SM19]: Theses systems

Comment [SM20]: They

Comment [SM21]: RAS

177 FISH SPECIES GROWN IN NIGERIA

178 Fish is very important in the diet of many Nigerians, high in nutritional value with complete array of amino
179 acids, vitamins and minerals (Akinrotimi *et al.*, 2007c). In addition, fish products are relatively cheaper (Table
180 5) compare to beef, pork and other animal protein sources in the country (Amao *et al.*, 2006). FAO (2007)
181 reported that fish contribute more than 60% of the world supply of protein, especially in the developing
182 countries. The major species cultured in Nigeria include tilapias, catfish and carp. However, the African catfish
183 species (*Clarias gariepinus* or *lazera*) are the most resistant and widely accepted and highly valued fish that
184 are cultivated in Nigeria. According to Adewumi & Olaleye (2011), it is major specie reared in Nigeria.

185 **Clarias spp. (*C. lazera* or *gariepinus*)** popularly known as Catfish

- 186 • Its body has no scales
- 187 • Has omnivorous feeding habit
- 188 • It preys heavily on other species and even on its own fry and fingerlings.
- 189 • Usually it is not stocked alone but along with tilapia which provides food for it.
- 190 • It has relatively slow growth rate (when compared to with common Carp and Heterotis)
- 191 • The flesh is very tasty and free from bones.

192 Other most common fish species that are grown in Nigeria earthen ponds include:

193 **Tilapia spp. (*T. guineensis* and *T. niloticus*)**

- 194 • Its whole body is heavily covered with scales
- 195 • It has omnivorous feeding habit

- Very prolific. Hence it reproduces at a very rapid rate.
- Its flesh is very bony but the bones can be separated and removed especially when the fish is big.
- The flesh has a delicious taste.

199 *Cyprinus carpio* (Common Carp)

- 200 • It has few large shiny scales
- 201 • It is sluggish and this makes it an easy prey for Clarias. Hence it is not advisable to stock it together
- 202 with Clarias in the same pond.
- 203 • Does not grow well without supplementary feeding.
- 204 • It grows at a fast rate when fed well and can attain 2kg in 8-10 months
- 205 • It has genetic potential to grow to a very large size (up to 10kg).

206 **Table 4: Production technique and fish cultured**

	%
Culture System	
Monoculture	80.4
Polyculture	5.4
Integrated	14.2
Production Technique	
Stagnant water	35.7
Flow Through System	48.2
Water Recirculatory	16.1
Type of fish raised	
Tilapia	5.4
Carp	14.6
Cat fish	73.2
Ornamental	5.4
Others	1.8

Comment [SM22]: of which species?

Comment [SM23]: of which species?

208 **Source: Field Survey 2003 – 2006**

209 **Table 5: Average Prices of some commonly consumed meat products in Southern Nigeria**

Meat product	Year		
	2000 N. K	2003 N. K	2010 N. K
Beef	300.00	500.00	800.00
Pork	280.00	380.00	700.00
Chicken	380.00	500.00	750.00
Goat meat	400.00	500.00	800.00
Cat fish	350.00	400.00	500.00
Tilapia	200.00	280.00	350.00

211 **Source (FDF, 2008)**

213 **Table 6: Production of Cultured Clarrid cat fish in best 20 countries in 2001**

S/N	Country	Quantity (t)
1	Thailand	146,000

2	Indonesia	77,332
3	Nigeria	61,916
4	Uganda	20,941
5	Malaysia	18,486
6	Netherlands	4,800
7	Philippines	2,376
8	Hungary	1,724
9	Syria	1,030
10	Cambodia	800
11	Poland	380
12	Brazil	362
13	Kenya	302
14	Mali	300
15	Belgium	250
16	Togo	200
17	Romania	118
18	Italy	115
19	Cameroon	110
20	South Africa	100

Source: FAO (2009)

PROBLEMS AND PROSPECTS OF AQUACULTURE IN NIGERIA

Nigeria as a nation is endowed with good natural resources which can enhance aquacultural development to a greater extent, however, the reverse is the case. Many researchers are wondering whether it is leadership problem or lack of skills required for the production or lack of awareness on the importance of aquaculture to national development. Of course, aquaculture benefits are numerous; it fulfils protein demand of the country, helps prevent food insecurity, creates jobs which settle the unemployment issue and it is prolific and profitable, generate income for individuals, serves as recreation centers for relaxation, can attract foreign exchange capable of increasing the country's GDP and so on. Most fishermen interviewed along the floodplain of the upper Niger expressed the view that their major problem is that of input supply rather than capital to acquire the input. This was the common opinion among flood pond fishermen whose capital inputs were simple and fairly cheap, being limited to gourds as floating objects instead of plank boats and with only clapnets and foul hooks as the principal fishing gears. The total capital input for a fishing unit among this category of fishermen is currently about \$70.00 or N700.00 (Naira). Their income, estimated at an average daily catch of 8 kg and 4 kg, for the good and bad fishing seasons respectively, and for a total of 160 fishing days in the year, amounts to about \$960.00 at an average selling price of \$1.00/kg of fish or N10.00 per kg of fish at 1989–90 prices (FAO, 2013).

The Problems: let us look at some researchers' view on the militants of aquacultural development in our country: Adewumi & Olaleye (2011) found out that a number of problems confront the production of catfish; being a major species in Nigeria. Prominent among these are: poor management skills, inadequate supply of good quality seed, lack of capital, high cost of feed, faulty data collection, lack of environmental impact consideration and marketing of products. If the associated problems of production, especially the twin issue of feed production and fingerling supply are tackled, Nigeria will soon become a world exporter of catfish. According to Oota cited in Oyinbo and Rekwot (2013), high cost of input, lack of credit facilities to fish farmers

Comment [SM24]: Delete
This paragraph is about capture fisheries only. Not relevant.

at low interest rate, lack of skilled manpower and lack of aquacultural extension service are attributes contributing to underdevelopment of aquaculture in the country. While George *et al* (2010) stated that the major problem hindering the promotion and development of the aquaculture industry in Nigeria has been the scarcity of fish fingerlings and that the major factors militating against the production of high quantity of fish seed are energy and water quality related problems arising from skills gap in the industry.

The Prospects: over the last 2 decades, Federal Government effort was on artisanal fisheries development in Nigeria because the fisheries were known to contribute over 95% of the local fish production. The National Scheme (National Accelerated Fish Production Programme) introduced by the Federal Government through the Federal Department of Fisheries provided fishing inputs such as engines, gillnets, ropes, lead and twines to fishermen with a 50% subsidy in order to assist them in attaining maximum capture efficiency (FAO, 2013). However, overfishing activities made by these fishermen have reduced the yielding capacity of the most fisheries. Aquaculture which is the current tool capable of increasing the country's total fish production if properly managed has not received a fair treatment by the government. International assistance for aquacultural development in Nigeria, government subsidy aquacultural inputs, shifting interest and resources of National Accelerated Fish Production Programme to aquaculture industry, provision of basic infrastructures, electricity and quality water supply, provision of extension services including education program to aquaculturalists and government support and involvement in production program at the federal, state and local level will significantly boost the country's aquacultural production.

FISH POND PREPARATION

Before a pond can be stocked with new fish, the excessive wastes, which accumulate in the pond during the previous farming, must be removed and the soil and water conditioned or upon a newly constructed pond, the following preparations have to be taken into consideration: cleaning, liming, eradication of predators, fertilization, and aeration.

Cleaning – There are two methods of cleaning; the drying method and wet method. The drying method occurs where the ponds can be completely dried like a concrete pond. Whereas, the wet method takes place in ponds like earthen ponds in which the water in the pond cannot be completely dried.

Liming – Once the pond is cleaned, it is then filled with water and left overnight before flushing out to remove debris and elevate the pH. This process should be repeated until the pH of the water remains above 7, and only then the lime is applied. The types of lime to be used depend on the water pH. It is recommended that agricultural lime (CaCO_3) or dolomite [$\text{CaMg}(\text{CO}_3)_2$] should be used in a pond with water pH near neutral and the hydrated lime [$\text{Ca}(\text{OH})_2$] should be used in a pond with water pH below 5 (KAU, 2013). The amount of lime to be used should be carefully calculated to avoid inducing an excessively high water pH, which may increase ammonia toxicity and result in the mortality of the fish stocked. When the pond is properly limed and filled with water, the average water pH should be between 7.5 - 8.5 with daily fluctuation of less than 0.5.

Eradication of Predators – After liming, the pond should be filled with water to the maximum depth through a screen with fine mesh to prevent the predators and competitors from entering the pond. Some chemicals

277 should be used to eradicate these animals in the pond before stocking. Fish can be killed by the application of
278 tea seed powder at the rate of 20-30 ppm. After the application of tea seed, the pond should be left for 3 days
279 before the fry, fingerlings and post larvae can be stocked. Hypochlorite, either calcium or sodium salt, is
280 currently used at 15-20% (60% active ingredient) to eliminate both vertebrates and invertebrates. The pond
281 must be cleaned prior to the application of hypochlorite since hypochlorite may react with the organic matters
282 and produces the toxic organochlorine compounds. Hypochlorite should be applied after the pond is filled to
283 the maximum height and left for 3 days to allow the hatching of planktonic organisms. After the hypochlorite
284 application, the pond should be aerated and the application of lime and fertilizer should be conducted on Day
285 3, then stock the pond with fish on Day 7. During the first month, water must not be added to the pond, unless
286 the water quality is poor, to prevent the introduction of competitors and predators.

287 **Fertilization** – The pond must be fertilized with either organic or inorganic fertilizer to stimulate the plankton
288 bloom in order to provide shade to the pond bottom and utilize the nitrogenous and phosphate wastes within
289 the pond. The shade will also prevent the growth of harmful benthic algae. The sun dried chicken manure is
290 the most common organic fertilizer to be used in amounts of 200-300 kg/ha. The manure must be soaked in
291 water for 24 hours before it is spread over the surface of the water. Inorganic fertilizers, such as urea (46% N)
292 and compound fertilizers like, ammonium phosphate (16:20:0) or those with N:P:K combination of (16:16:16)
293 can be used at 20-30 kg./ha (KAU, 2013). The fertilizer must be dissolved in water before it is spread over the
294 water surface to avoid precipitation of the fertilizer onto the pond bottom, which will enrich the soil and
295 accelerate the growth of benthic algae.

296 **Aeration** – 0.5-1.0 ha pond would require, four aerators installed at the corners of the pond, approximately 3-5
297 m from the bottom of the dike and positioned at an angle that will encourage the maximum water flow within
298 the pond. The type of aerator to be used depends on the depth of the water. One horsepower paddle wheel
299 aerators should be used in ponds of less than 1.2 m water depth and the 2 HP (horsepower) paddle wheel
300 aerators should be used in ponds deeper than 1.2 m. The aerators should be switched-on 24 hrs before the
301 post larvae are stocked to allow enough time to create the current and clean up the feeding area (KAU, 2013).

302 **FISH FEED AND FEEDING**

303 **Fish feed** is a plant (phytoplankton) or animal (zooplankton) material intended for consumption by fish, kept in
304 aquacultural systems. Fish feeds normally contain macronutrients, proteins, fats, fibre, trace elements and
305 vitamins necessary to keep captive fish in good health. Depending on the type of the system in which the fish
306 is being reared (intensive, semi-intensive and extensive system). The fish can be fed by natural and
307 supplementary feeds or combination of both. Riche and Garling (2003) reported that fish reared in intensive
308 tank systems requires all nutrients in a complete pelleted diet since natural food is limited and fish cannot
309 forage freely for natural foods. There are three (3) types of aqua feeds are produced worldwide: floating, slow-
310 sinking and sinking to meet the nutritional requirement of fish species under culture and the culture system.
311 The sizes are as follows: 0.5mm, 0.8mm, 1mm, 2mm, 3mm, 4.5mm, 6mm, and even 10mm (1cm) for larger
312 fish. The sinking (hard) pellets usually sink to the bottom of the aquacultural facilities while the floating does

313 not. The floating ability depends on the degree of fineness of the grounded food components and mixing power
314 of the machine.

315 **Feeding** is usually the largest operational cost of growing fish in aquaculture (Paolucci et al., 2012). The feed
316 can represent greater than 50% of the variable costs in growing fish. Labour and feeding equipment are
317 contributors to the total cost of feeding the aquacultural crop. Delivering the feed to fish at the right time, in the
318 correct form, and in the right amount is necessary for optimal growth. The choice of feeding equipment
319 depends on factors such as type and life stage of fish, type and size of feed, size of the operation, available
320 labour, and the type of culture system. Therefore, a careful analysis of these factors is essential for successful
321 and profitable fish culture.

322 There are three groups of feeders and feeding methods—hand, automatic, and demand—used to deliver
323 formulated diets. The type of feeding system depends on the life stage of fish being cultured, size of operation,
324 type of diet, available resources, and personal preferences. This is because larval and smaller fish require
325 frequent feeding, they are often fed with automated feeders that can be set to feed throughout the day at
326 regular intervals. Smaller aquacultural operations may have sufficient labour to feed all stocks by hand while in
327 larger operations, hand feeding becomes a problem unless larger, automated equipment is used. Whatever
328 method is used to facilitate feeding, the operator must periodically check for feeding effectiveness by sampling
329 and weighting fish, calculating feed-conversion ratios, and adjusting the feeding rates (Summerfelt and Wade,
330 1997).

331 **Hand Feeding** – Manual or hand feeding can be used in all types of aquacultural systems - cage, net, pond,
332 flow-through, and recirculating and is often the method of choice for small systems. Hand feeding can reduce
333 capital expenditures for equipment, provided sufficient labour is available to perform the feeding duties. There
334 are other advantages to hand feeding; the fish can be observed at each feeding for feeding behaviour to
335 determine if they are actively eating all of the food presented to them. Floating pellets make observations
336 easier. Fish can only be routinely fed to satiation by hand feeding. Observing the fish during feeding can also
337 be critical in the early determination of disease or parasite problems.

338 **Demand Feeders** – The cultured fish can be allowed to determine for themselves how much feed is made
339 available when demand feeders are used. Demand feeders can be used to make food continuously available
340 and allows for the fish to feed to satiation. Typically, fish fed with demand feeders consume amounts of feed
341 similar to what they would eat. The demand feeders must, however, be readily accessible to the fish, and
342 sufficient numbers of feeders must be strategically located around larger systems. Fish are trained to hit the
343 trigger of a demand feeder when hungry and then become self-feeding. Some demand systems use a touch-
344 sensitive trigger to activate some kind of mechanical delivery system. Other modifications of the demand
345 feeder use an in-water feed tray that activates feed release as feed is eaten from the tray. Usually, all of the
346 demand feeders are adjustable for food size and amount released per trigger activation. It takes classically
347 about 7 to 10 days for rainbow trout to train and feeding to stabilize on demand feeders (Summerfelt and
348 Wade, 1997). Demand feeders make feed available 24 hours per day and result in less size variability in the
349 harvested fish. Trout fed with demand feeders can have up to 10% better feed conversion than in hand-fed

systems. Other advantages from demand feeders are lower production costs from reduced labour requirements and increased feeding efficiency. Trout on demand feeders showed fewer disease problems and fewer problems with dissolved oxygen sags and ammonia spikes from heavy feedings.

Automatic Feeders – An automatic feeder is basically a mechanism that delivers a prescribed amount of feed to the fish at desired time intervals. The major advantage of automatic feeders is the reduction in labour equipment necessitated by hand feeding, especially in larger operations. Automated feeders have been adapted from other livestock operations to provide cost-effective means to meet the array of needs for aquaculture. One key feature of automatic feeders is some kind of programmable device that allows the operator to vary the frequency and amount of feeding. This may be a combination of electrical and mechanical components that broadcast feed into the culture system. Some systems use electrical timers to determine the frequency, which are available from most aquacultural equipment suppliers. Ideally, the settings for frequency will allow enough flexibility to meet the needs of the fish. For smaller fish, the frequency may need to be several times per hour, and for larger fish the frequency could be hourly or longer (Goddard, 1996). Automatic feeders use some kind of mechanical or pneumatic delivery system to present feed to the fish.

PRODUCTION MANAGEMENT

The purpose of fisheries management is to provide good fishing that will lead to increased yield. Pond owners must decide what they want from their pond and tailor their management to meet their goal(s). Fish business is generally prolific if proper handling and management precautions are adapted. The major management practices necessary for effective production are as follows; fish stocking density, feeding, water quality control, diseases control and record keeping. In the case of feeding, it has already been discussed in this article.

Comment [SM25]: delete

Fish Stocking Density – Stocking marks the beginning of production cycle. Stocking density of any aquacultural pond has to be first and foremost considered in management principles. This is because if a pond exceeds its carrying capacity, fish stress is bound to occur which can eventually lead to fish mortalities. The process of stocking referred to here, starts with the collection of fingerlings from the hatchery, transporting them to the farm and, finally, putting them into the pond. Poor stocking procedures, are among the major causes of low survival in grow-out ponds Nelly *et al* (2009). They result in stress, diseases and reduced growth and eventually lead to mortality and financial losses. Also, quality fingerlings are another important factor to note while stocking. Poor quality stock will give poor production performance regardless of other factors. The most important practical criteria for assessing the quality of fingerlings are source, physical appearance and how they swim. A Pond's carrying capacity is influenced by the following: the size of fish in the pond (because this influences the feeding rate); the species of fish being raised because fish like *clarias spp* become air breathers and do not need to rely on dissolved oxygen in the pond, therefore the carrying capacity is higher for *clarias spp* compared to Tilapia; the amount and type of feed or fertilizer added to the pond and the water volume and quality.

The equation below illustrates how the number of fish to be stocked in a pond can be estimated based on the pond's carrying capacity by Nelly *et al* (2009).

$$\text{Number to stock} = \frac{\text{Estimated maximum carrying capacity of a pond (kg)}}{\text{Desired weight at harvest (kg)}} + 10\%$$

387

388

389 EXAMPLE: The size of pond to be stocked is 1,000 m². The size of fingerlings is 15 g and the targeted size for
390 harvest is 800g. The intended management regime is catfish monoculture fed entirely on commercial
391 nutritionally-complete pellets in static water ponds.

392 How many fish should be stocked?

393 The critical standing crop for catfish monoculture fed commercial pellets is estimated at 1.8 kg/m² (18 tons/ha).

394 Add 10% to account for mortalities.

395

396
$$= \frac{(1,000 \text{ m}^2 \times 1.8 \text{ kg/m}^2)}{0.8 \text{ kg}} + 10\% \text{ of } \frac{(1,000 \text{ m}^2 \times 1.8 \text{ kg/m}^2)}{0.8 \text{ kg}}$$

397

398
$$= 2,250 + 225$$

399
$$= 2,475 \text{ fingerlings should be stocked}$$

400

401 **Water Quality** – Water quality is the second aspect to consider in aquacultural management. The failure or
402 success of production lies on this key factor. Poor water quality at high level does not spare the life of the fish
403 for a minute. Therefore, this has to be checked regularly as prescribed below. The basic water quality
404 parameters that are important in maintaining fish health include dissolved oxygen (>4.0 mg/l), salinity (15-32
405 ppt), temperature (26-30°C), pH (7.0-8.5), nitrogen compounds [ammonia (< 0.15 mg/l), nitrate and nitrite (<
406 4.5 mg/l)], BOD [5day (50/30mg/l)], hydrogen sulphide (< 0.1 mg/l) and pesticides (Andrew, 2007).

407 These should all be checked at different times: oxygen and temperature are measured at least twice daily to
408 determine the influence of photosynthesis on concentrations. Salinity and pH are measured daily, and nitrate,
409 ammonia and nitrite can be measured 2-3 times per week. Pesticides (those known to be used in watershed)
410 should be tested periodically and at different rainfall levels to determine the effect of runoff on concentrations.
411 Also check during periods of high rainfall increased water flow that influences water quality and may increase
412 pesticides, sediment, and organic loading. Measure other parameters several times over a month period to
413 know ranges. Know what pesticides are being used in the watershed and when, understand their toxicity, and
414 test water periodically for those compounds.

415 **Fish Diseases** – High productivity in aquacultural production only transpire when fish are healthy and free
416 from diseases. Fish disease management is aimed at preventing the onset of disease and measures to reduce
417 losses from disease when it occurs. Let us see some common fish diseases, their causes and management
418 control:

419 **Vibriosis** is a bacterial disease causing significant losses of fish in marine fish farms. Vibriosis accounts for an
420 estimated two-thirds of disease reported in grouper species. Vibriosis results in severe skin, muscle, fin, eye
421 and internal organ damage of fish. Stressors that trigger vibriosis outbreaks include high water temperatures,
422 high stocking densities, poor handling of fish, and an organically polluted culture environment. Antibacterial
423 medication, reducing stocking densities, careful handling of fish, improving the culture environment through the

use of clean pelleted feeds, stocking of fish in cooler season of the year and vaccination are important control measures (Roger *et al.*, 2011).

Streptococcosis is a disease of freshwater pond fish cultures, particularly *Streptococcus agalactiae* of tilapia (*Oreochromis spp.*). Clinical signs include cloudy eyes, exophthalmos and focal, ulcerative skin lesions along the caudal peduncle area. Control of streptococcosis relies on the administration of an antibiotic to which the isolate is sensitive together with control of any associated ectoparasites and appropriate risk reducing husbandry measures. Preventative measures are dependent on an effective vaccine against the pathogenic strains of Streptococcus present in the fish environment or to which the susceptible fish stocks are likely to encounter (Roger *et al.*, 2011). Others diseases include Benediniasis, viral hemorrhagic septicemia, Myxobolus cerebralis (whirling disease), *Ichthyophthirius multifiliis*, *Epinephelus lanceolatus*, Viral Encephalopathy and Retinopathy (VER) Disease.

Record Keeping

Keep thorough records from the onset of stocking to the harvesting period — they are valuable in health management. Keep accurate records of numbers and sizes of fingerlings, time, type and quantity of the feed fed, periods of water changed, time of aeration, date of treatment and vaccination, etc. These records will help you evaluate the status of your fish populations and provide you with proper guide to next farming.

CONCLUSION

Standards for development of aquaculture are not beyond the country's potentials. As can be deduced from the review on factors that hinder the development of aquaculture in the country, the problem is simply considered to be lack of management skills, support and the scarcity of the fingerlings. Nevertheless, proper aquacultural system design, good stocking and water quality management are essential to successful and quality fish production. Consequently, maintaining a good culture environment through use of proper management practices will reduce the risk of disease and increase production, fish quality, and marketability. Therefore, it is essential to implement and follow good aquaculture management practices to ensure quality and quantity yield. The agricultural sector is a key to unlocking the padlock of food insecurity, let alone its subsector, aquaculture. Statistics indicate that Nigeria spends N100 billion on fish importation annually. We cannot continue to rely on importation of every consumer goods, even the ones the country has both natural resources and man-power capacity to produce comfortably and successfully. Aquaculture business is viable in Nigeria from all indications but needs more efforts from the both public and government to increase its local production so as to bridge the gap between the fish demand and supply in the country.

REFERENCES

- Adedeji O.B. and Okocha R.C. Constraint to Aquaculture Development in Nigeria and Way Forward" Veterinary Public Health and Preventive Medicine University of Ibadan, Nigeria, 2011.
- Adeogun O.A, Ajana A.M., Ayinla O.A., Yarhere M.T. and M.O. Adeogun. Application of Logit Model in Adoption Decision: A Study of Hybrid Clarias in Lagos State, Nigeria. American-Eurasian J. Agric. & Environ. Sci., 4 (4): 468-472, ISSN 1818-6769 © IDOSI Publications, 2008

460 Adewumi, A.A & Olaleye, V.F. Catfish culture in Nigeria: Progress, Prospects and Problems. African Journal of
461 Agricultural Research, 2011; 6(6): 1281-1285

462 Ajayi TO, Talabi S.O. The potential and strategies for optimum utilization of the fisheries resources of Nigeria.
463 N.I.O.M.R. Technical Paper No. 18, 25p,1984.

464 Akinrotimi, O.A. Ansa, E.J. Owhonda, K.N. Edun, O.M. Onunkwo, D.N. Opara, J.Y. Anyanwu P.E. and
465 Amachree D. Variation in oxygen carrying capacity of Sarotherodon melanotheron blood in different
466 acclimation media. Journal of Animal and Veterinary Advances. 2007; 6(8): 932 – 937.

467 Akinrotimi, O.A.; Abu, O.M.G; Aranyo, A.A. Environmental Friendly Aquaculture Key to Sustainable Fish
468 Farming Development in Nigeria. Continental J. Fisheries and Aquatic Science 5 (2): 17 – 31. ISSN:
469 2141 – 4246 © Wilolud Journals, 2011

470 Amao, J.O., Oluwatayo, I.B. and Osuntope, F.K. Economics of Fish Demands in Lagos State, Nigeria. Journal
471 of Human Ecology. 2006; 19(1):25 – 30.

472 Andrew Lazur. Grow-out Pond and Water Quality Management. JIFSAN (Joint Institute for Safety and applied
473 Nutrition) Good Aquacultural Practices Program, University of Maryland, 2007.

474 EAI, Emperor Aquatics, Inc. (1990). Aquatic Life Support Systems (ALSS). Sanatoga Station Road, Pottstown,
475 Pennsylvania, USA. info@emperoraquatics.com

476 Ekunwe, P.A and C.O. Emokaro. Technical Efficiency of Catfish Farmers in Kaduna, Nigeria Journal of Applied
477 Sciences Research., 2009; 5(7): 802-805.

478 FAO, Food and Agriculture Organization of the United Nation Fish stat: Universal Software for Fishery
479 statistical time series V.23. Data and statistics Unit, FAO, Rome Italy, 2009.

480 FAO, Food and Agriculture Organization of the United Nations. The state of World Fisheries and Aquaculture
481 2006. FAO Fisheries Department, Rome Italy, 30pp, 2007.

482 FAO, Food and Agriculture Organization of the United Nations. The State of the World Fisheries and
483 Aquaculture FAO, Rome Italy, 2006.

484 FAO, Food and Agriculture Organization of the United Nations. Inland Fisheries Resources of Nigeria.
485 Corporate Document repository. Produced by Fisheries and Aquaculture Department.
486 <http://www.fao.org/documents/en/detail/64969>. Accessed online on 7th of Decemebr, 2013.

487 FDF, Federal Department of Fisheries. Fisheries Statistics of Nigeria, Fourth Edition 1995 – 2007; 49pp.

488 FDF, Federal Department of Fisheries. Fisheries Statistics of Nigeria Projected human population; fish demand
489 and supply in Nigeria from 2000 – 2015, 2008; 56pp.

490 George, F.O.A., Olaoye, O. J., Akande, O. P. & Oghobase, R. R. Determinants of Aquaculture Fish Seed
491 Production and Development in Ogun State, Nigeria. Journal of Sustainable Development in Africa,
492 2010; 12(8): 22 – 34.

493 Goddard, S. Feed Management in Intensive Aquaculture. Chapman and Hall, New York, 1996.

494 Green, Bartholomew W. Aquaculture. Microsoft® Encarta® 2009.

495 Gupta, M.V. & Acosta, B.O. Tilapia farming: A global review. WorldFish Center, Penang, Malaysia, 2004.
496 Accessed online on 29.9.2011.

497 Helen Chapin Metz, ed. Nigeria: A Country Study. Washington: GPO for the Library of Congress, 1991.

498 Ikenweawe N. Bolatito. Lecture Note on Fisheries Stock Assessment (2 units) FIS 318. Department of

499 Aquaculture and Fisheries Management, University of Agriculture Abeokuta, Nigeria. Accessed on 7th

500 December, 2013.

501 Jhingran, V.G., Introduction to aquaculture United Nations Development Programme, Food and Agriculture

502 Organization of the United Nations, Nigerian Institute for Oceanography and Marine Research, 2001.

503 Kareen R.O., Dipeolu., Aromolaran A.B and Williams S.B. Economic efficiency in Fish Farming: hope for agro-

504 allied industries in Niagara. Chinese Journal of Oceanology and Limnology. Vol. 26 No. 1. P. 104-115,

505 2008.

506 KAU, Kerala Agricultural University Pond Preparation Infotech Portal Centre for E –Learning Kerala Agricultural

507 University. Accessed online 10th December, 2013.

508 Magdalene Ukuedojor. Adequate drinking water still a tall dream in Nigeria. Nigerian newspaper: Weekly Trust.

509 Published on Saturday, 05 January 2013 06:00.

510 Mirzoyan, N., Tal, Y., Gross, A. Anaerobic digestion of sludge from intensive recirculating aquaculture systems:

511 Review In Aquaculture 306 (2010) 1–6.

512 Nelly A. Isyagi, Karen L. Veverica, Rashid Asimwe, and William H. Daniels. Manual for the Commercial Pond

513 Production of the African Catfish in Uganda Department of Fisheries and Allied Aquacultures; Auburn

514 University, Alabama, USA, 2009.

515 Nwosu FM, Ita EO and Enin UI. Fisheries management in Nigeria: A case study of the marine fisheries policy.

516 International Research Journal of Agricultural Science and Soil Science Vol. 1(3) pp.070-076, May

517 2011. Available online <http://www.interestjournals.org/IRJAS> Copyright © 2011 International Research

518 Journals

519 Off-shore fish farming term. Accessed online on 7th December, 2013.

520 Olagunju F.I, Adesiyani I.O, Ezekiel A.A. Economic viability of cat fish production in Oyo state, Nigeria. J. Hum.

521 Ecol. 21(2):121 – 124, 2007.

522 Olomola Ade S. Strategies for Managing the Opportunities and Challenges of the Current Agricultural

523 Commodity Booms in SSA in Seminar Papers on Managing Commodity Booms in Sub-Saharan Africa:

524 A Publication of the AERC Senior Policy Seminar IX. African Economic Research Consortium (AERC),

525 Nairobi, Kenya, 2007.

526 Oyinbo Oyakhilomen and Rekwot Grace Zibah. Fishery Production and Economic Growth in Nigeria: Pathway

527 for Sustainable Economic Development. Department of Agricultural Economics and Rural Sociology,

528 Ahmadu Bello University, Nigeria – Journal of Sustainable Development in Africa. Vol. 5. No.2, 2013.

529 Ozigbo Emmanuel, Anyadike Chinenye, Forolunsho Gbadebo, Okechuckwu Richardson and Kolawole Peter.

530 Development of an Automatic Fish Feeder” International Institute of Tropical Agriculture Postharvest

531 Unit, Ibadan – African Journal of Root and Tuber Crop. Vol.10 No.1 Page 27-32, 2013.

532 Paolucci, M., Fabbrocini, A., Volpe, M.G., Varricchio, E., Coccia, E. Development of biopolymers as binders for
533 feed for farmed aquatic organisms, in: Muchlisin, Z.A., (Ed.), Aquaculture, 2012. Vol. 1, InTech, (ISBN
534 979-953-307-107-6), Open access publisher.

535 Riche, M., D. Garling. Feeding Tilapia in intensive recirculatory systems. North central Regional Aquaculture
536 Centre and United State Department of Agriculture USDA, 2003: 1-4pp.

537 Roger Chong, Barry Bousfield and Richard Brown. Fish Disease Management. Veterinary Bulletin -
538 Agriculture, Fisheries and Conservation Department Newsletter, Hong Kong. Volume No. 1 Issue No. 8.
539 December, 2011.

540 Summerfelt, S. T. and E.M. Wade. Recent advances in water treatment processes to intensity fish production
541 in large recirculating systems. in M. Timmons (ed) successes and failure in commercial Aquaculture
542 (Conference Proceedings), 1997.

543

544

545