

Original Research Article

Identification of Hematological Markers Suitable for Improving Productivity of Helmeted Guinea Fowl *Numida meleagris*

ABSTRACT

The research was designed to estimate variations in hematological variables between sex and varieties of extensively reared helmeted guinea fowl with the objective of recommending parameters that are suitable as markers during selection. The study was carried out in the teaching and research farm of Department of Animal Science, Ahmadu Bello University, Zaria between May and July 2012. Five varieties of helmeted guinea fowl (45 males and 45 females) that were clinically healthy were sampled for white blood cells (WBC), packed cell volume (PCV), red blood cells (RBC), hemoglobin, platelets, Mean Corpuscular Hemoglobin Concentration (MCHC), Mean Corpuscular Hemoglobin (MCH) and Mean Corpuscular Volume (MCV). The results showed that there were no significant ($P > .05$) effect of sex on WBC, MCHC, RBC, hemoglobin, PCV and MCH. There were significant ($P = .05$) effects of varieties on all the hematological parameters determined in this study. The study revealed that sex has influence on MCV and platelets while RBC, WBC, PCV, hemoglobin, MCHC, MCH, MCV and platelet differed with varieties. Therefore, the study of the variations in hematological parameters as well as their association with economic traits could be used to develop marker assisted technology which can be incorporated into the traditional animal breeding methodology to fast track improvement to aid selection, improve efficiency and speed of selection in breeding programs.

Keywords: *Helmeted Guinea Fowl; Hematology; Selection; Markers.*

1. INTRODUCTION

In most African countries especially Nigeria, guinea fowl production is in its infancy [1,2] yet guinea fowl is a promising genetic resource for evolving a low input – grain saving poultry alternative for production in the developing world. The rearing of guinea fowl is thus a potential alternative poultry system. Guinea fowl is gaining importance in the Nigerian poultry industry [3] hence the need for extensive and intensive studies into every aspect that is likely to contribute to the future improvement of this species.

Determination of hematological parameters is essential in diagnosing various pathological and metabolic disorders [4]. It provides highly valuable information on the physiological state and form the cornerstone of medical diagnosis of diseases [5,6]. However, hematological parameters provide information that can be useful not only for diagnostic and management purposes, but could also be incorporated into breeding programs for genetic improvement [4].

There is a lack of exact data on the hematological profiles of helmeted guinea fowl reared in Nigeria. Values for the hematology of helmeted guinea fowl have been reported [7,8,9] in the southern parts of Nigeria as it affects sex and system of production [10]. The objective of the study is to identify hematological parameters during the breeding season that are suitable as markers for both sexes and varieties of the extensively-reared helmeted guinea fowl in Northern Nigeria.

2. MATERIAL AND METHODS

2.1 Study Location

The research was conducted on the teaching and research farm of Department of Animal Science, Ahmadu Bello University, Zaria, Kaduna State, Nigeria.

2.2 Sample Size and Sample Procedure

A total of 45 males and 45 females of five varieties of helmeted guinea fowl classified in an earlier study [11] comprising of Ash, Black, Pearl Ash, Pearl Black and White varieties were used for this study. The birds which had been extensively reared were sourced from seven different locations in Katsina and Kaduna states, Northern Nigeria [12]. Blood collection was as described in an earlier study [13]. About 5ml of whole blood was collected by superficial venipuncture of a wing vein using a 5ml syringe. The whole blood was collected into Ethylene Di – amine Tetra Acetic acid (EDTA) bottles and taken to the laboratory for analysis within 3 hours of collection.

The following variables were analyzed for by an automated system using the COULTER™ LH 750 Hematology analyzer by BECKMAN COULTER, Inc, Orange county, California, United States of America;

- i) Red blood cell (RBC)
- ii) White blood cell (WBC)
- iii) Packed Cell Volume (PCV)
- iv) Hemoglobin
- v) Mean Corpuscular Hemoglobin Concentration (MCHC)
- vi) Mean Corpuscular Hemoglobin (MCH)
- vii) Mean Corpuscular Volume (MCV)
- viii) Platelet

2.3 Statistical Analysis

The data was analysed using General Linear Model (GLM) procedure of Statistical Analysis System [14]. The differences between sex and between varieties of helmeted guinea fowl in haematological parameters were determined using the Tukey's studentized range (HSD) test.

The statistical model used was;

$$Y_{ijk} = \mu + V_i + S_j + e_{ijk}$$

Where,

Y_{ijk} = observation on the j^{th} sex of the i^{th} variety of helmeted guinea fowl

μ = overall mean.

V_i = the effect of i^{th} variety of helmeted guinea fowl ($i = 1-5$)

S_j = the effect of j^{th} sex of helmeted guinea fowl ($j = 1-2$)

e_{ijk} = residual error

3. RESULTS AND DISCUSSION

3.1 Sex differences in hematology of helmeted guinea fowl

The hematological parameters determined in the male and female guinea fowls are as indicated in Figure 1. The WBC and MCHC were slightly higher in females while RBC, hemoglobin, PCV and MCH in males though the differences were not significant ($P > .05$). However, the males had significantly ($P = .05$) higher MCV than the females while the females had significantly ($P = .05$) higher platelet value.

The values of PCV in this study were similar to values obtained for the Pearl helmeted guinea fowl [10,15,16]. The RBC values of both sexes were similar to those reported by¹⁵ and within the general reference range [17] of avian ($2.1 - 5.5 \times 10^{12}/l$). Hence, any of the values was representative of a healthy helmeted guinea fowl. Hemoglobin is a component of the red blood cells and in this study is also within reference range [17] for avian (12.20 – 20.00g/dL). This result is also consistent with the report of no sex effect on WBC, RBC, MCHC, PCV, MCV, MCH and hemoglobin in yellow – legged gulls [18]. Similarly, no significant ($P > .05$) influence of sex on RBC, hemoglobin and WBC was reported in male and female Ring – necked pheasants [19] and in turkey [20]. This result however differed from those obtained by some authors who reported significant effect of sex on blood hematology in helmeted guinea fowl [10] and in Sudanese indigenous chicken [21]. In their report, PCV, RBC and hemoglobin were higher ($P = .05$) in males than females and these differences were attributed to increased courting of the females by the males during their breeding season which was when they were bled, as hemoglobin and PCV were influenced by reproductive hormones; androgen and estrogen which together depress erythropoiesis [22] whereas androgen and thyroxine stimulates erythropoiesis [23]. The difference in result on sex influence may be attributed to differences in experimental setup and environment.

RBC, PCV and hemoglobin are responsible for enhancing oxygen delivery to the testes of naked neck cocks [21] and are also associated with improved semen quality [24]. More so, increased number of RBC and PCV in cocks correspond with the time of androgen production [25,26]. Thus, increase in PCV could be used to predict spermatozoa maturation in testes, sexual maturity and the start of semen production [27] while can be used as indicator of feed conversion efficiency [28].

WBC which protects the body against invading microorganisms and body cells with mutated DNA, and clean up debris [29] is similar in both sexes and are within reference range of similar birds reared extensively [19,21] indicating that genetically, both sexes had equal potentials for immunity to diseases.

MCHC and MCH values were within the range obtained for Nigerian Indigenous chicken [30] and Sudanese Indigenous chicken [21] respectively. MCHC and MCH were not affected by sex which indicates that there was similar cellular haemoglobin content in blood samples obtained. An increase in MCHC value does not occur since RBCs do not become supersaturated with hemoglobin. However, a reduced value of MCHC usually indicates a deficiency anemia resulting in an insufficiency of hemopoietic factors [17]. MCH has been reported as not such a useful index as the MCHC [17].

Sex significantly influenced MCV and platelets in this study. MCV values in both sexes were lower than those reported for the Sudanese indigenous chicken and helmeted guinea fowl [10,21] but close to the value ($176.73 \pm 33.08fL$) reported for helmeted guinea fowl [15]. MCV give an indication as to the type of anemia and may also suggest the etiology of the


135 disease process [17]. An increase in this value is most useful for indicating a regenerative
 136 anemia. In a non-regenerative anemia, the cells are either normal or reduced in size.
 137

138 Sex also influenced platelets values in this study in accordance with the reports of [23,31,32]
 139 on avian hematology. Hematological parameters that have bearing on productivity of
 140 livestock should be incorporated in the formulation of selection indices for optimum
 141 productivity [33]. Platelets are essential for the repair of blood vessels when damaged; they
 142 also provide growth factors for healing and repair [29]. Low platelet concentration suggests
 143 that the process of clot-formation (blood clotting) will be prolonged resulting in excessive loss
 144 of blood in the case of injury [34] and this study indicated that the sexes had differing abilities
 145 to prevent excessive bleeding from occurring.
 146
 147

148

149

150






Figure 1: Influence of sex on hematological parameters in helmeted guinea fowl

3.2 Differences in hematology of helmeted guinea fowl varieties

The values of the various parameters of the hematological parameters in five varieties of helmeted guinea fowl are indicated in Figure 2. There were significant ($P = .05$) differences between the five varieties of the helmeted guinea fowl in all the hematological parameters determined in this study.

The Black, Pearl Black, Pearl Ash and White varieties were similar in WBC, RBC, hemoglobin and PCV; and had significantly ($P = .05$) higher values for the WBC than Ash variety. The MCV values were similar ($P > .05$) for Ash, Black, Pearl black and White but the Pearl Ash had significantly ($P = .05$) higher values than the other four varieties. The MCH value of the Pearl Ash was significantly ($P = .05$) higher than the other four varieties. The values in the Black, Pearl Black and White were similar ($P > .05$) but higher ($P = .05$) than in the Ash variety. The platelets value was significantly ($P = .05$) higher in the Black than the other four varieties. The White variety also had significantly ($P = .05$) higher value than in the Ash, Pearl Ash and Pearl Black had similar values ($P > .05$).

The significance of the testing for the WBC levels is for detecting and measuring the ability to destroy diseases that may be present in the body. Significant differences were observed in this study due to variety of the helmeted guinea fowl. This indicated that although the helmeted guinea fowl used appeared clinically healthy and had been treated for ecto – and endo – parasites, given broad spectrum antibiotics, good quality feed and housing before commencement of blood collection, there may still be some undiagnosed infections which the bird may be fighting against due to the fact that they were reared on free – range from the juvenile stage to adulthood and may have developed immunity against such. The five varieties of helmeted guinea fowl had varying abilities to fight diseases as indicated by the varying levels of white blood cells. The very high white blood cells could also be a factor that enhances higher disease resistance of the helmeted guinea fowl compared to other poultry. It is also possible that the change in the management system accounted for the high white blood cells levels. WBC was similar to the range ($21.50 - 28.23 \times 10^3/\mu$) reported [35] for Nigerian local and exotic chickens and the birds may be immunologically challenged which may be largely informed by the associated inherent resistance to tropical diseases [36]. It is also noteworthy that the subsisting ecological condition support several vectors and parasites of economic significance.

The RBC is a function of other components such as PCV, hemoglobin, MCH, MCHC etc and in this study the variety of the helmeted guinea fowl influenced the values. The Black, Pearl Ash, Pearl Black and White varieties have superior values of red blood cells to the Ash variety. This indicated varietal differences since all the birds used were of similar ages, were

all sexually matured and the females were in – lay and under same environmental factors. The RBC values in this study were within the range of $2.20 \times 10^{12}/L$ [37] and $2.15 (T/L)$ [15] for helmeted guinea fowl and $2.10 - 2.58 (T/L)$ for feathered games. They were however higher than $1.13 \times 10^{12}/L$ in another report [10] for helmeted guinea fowl. The ideal range of red blood cells of helmeted guinea fowl during the reproductive phase based on this study was $2.06 - 2.20 \times 10^6/\mu l$ since there were no significant differences between varieties except in the Ash.

PCV is a component of red blood cells hence the pattern of significant differences was similar. The PCV values obtained in this study ($35.71 - 37.14\%$) can be regarded as normal as the birds were in – lay and the values were similar to previous reports for laying Pearl helmeted guinea fowl [29] and adult helmeted guinea fowl [10,15] where values of 34.52% , 36% and 37% respectively were reported. The values were however lower than the reference range for psittacines, $40 - 55\%$ [32] and $42.60 \pm 1.60\%$ for frizzled feathered Nigerian indigenous chicken [35] which have similar phenotypic features as helmeted guinea fowl. This is because higher PCV values are usually due to a higher weight gain as well as reduction in heat load [38]. Selection of helmeted guinea fowl with significantly high PCV would therefore lead to a boost in the growth and productive life [35]. The range of values ($35.71 - 37.14\%$) obtained in this study, except in the White variety could be used to establish the ideal range for PCV as a reference range for the helmeted guinea fowl during the breeding season.

The MCV values obtained in this study were within the reference range for small psittacines, $121 - 200fl$ as reported [17] and could be used as reference values for the helmeted guinea fowl during breeding. The values, obtained in this study were similar to $176.73(fl)$ [15] for helmeted guinea fowl but higher than the $130.82 - 141.19(fl)$ [35] for Nigerian local and exotic chickens. MCV values were also higher than 86.99 , 105.88 and 116.66 cubic micron for Fayoumi, Assil and local chicken, respectively [31]


Hemoglobin is also a component of RBC hence the pattern of significant differences was similar. Except for the White variety, the other four varieties had similar hemoglobin values in this study. This indicates that the birds were in similar health status since variations in hemoglobin can be due to living at high altitudes, excessive/under production by bone marrow of blood cells, anemia due to nutritional deficiency, destruction of blood cells internally, liver tumor (hepatomas), kidney tumor and lung diseases. The range ($13.61 - 14.19g/dL$) obtained could be used for as a reference for breeding *Numida meleagris* during the breeding season in Northern Nigeria. The reference range as obtained in this study was similar to $12.36 - 16.37g/dL$ for Nigerian local and exotic chickens [35]. This indicated that vital physiological relationship of hemoglobin with oxygen and the transportation of gases (oxygen and carbon dioxide) to and fro the tissues of the body has been maintained and was normal [35].

The MCHC values in this study indicated significant differences between varieties. MCHC denotes the volume and character of hemoglobin and is the most accurate of red cell indices. The values ($33.92 - 40.63g/dL$) in this study could serve as reference for breeding *Numida meleagris* in Northern Nigeria by recommending them as a range because Pearl Ash variety had a significantly higher value while the Ash variety had a significantly lower value. The values obtained were similar to previous reports for feathered games: helmeted guinea fowl *Numida meleagris* 35.44% ; common pheasant *Phasianus colchicus* 31.27% ; Chukar *Alectoris chukar* 30.40% ; Wild turkey *Meleagris gallopavo* 30.71% (Strakova *et al.*, 2010). Wide variations ($33.92 - 40.63g/dL$) in the results could be due to age of the cells since the values were high. It has been reported that if cells are old and not replaced often (sperocytosis) and *in vivo* and *in vitro* hemolysis occur high MCHC values result; and if the

values are obtained when there are too many young RBC (reticulocytes) or when there is iron deficiency, low values are obtained. The MCHC values (33.92 – 40.63g/dL) in this study were within the reference range of 28 – 38g/dL for psittacine [17].

The MCH values in this study differed significantly between varieties of the helmeted guinea fowl. The MCH value followed a similar pattern as red blood cells; since it is a component of the RBC. The test defines the type of anemia and cause of anemia if present in species. Wide variations in MCH values could be due to anemia as a result of vitamin B₁₂ or/ and folate deficiency, ineffective production in the bone marrow, recent loss and replacement by newer reticulocytic cells from bone marrow and *in vivo* or *in vitro* hemolysis. Similar values of 60.88 [37] and 62.07(pg) were reported which were lower than the 120.33(pg) in another study [10]. The values (64.19 – 71.56pg) obtained in the present study can be recommended as a range for the extensively raised helmeted guinea fowl.

The platelets in this study were significantly ($P = .05$) different across variety. This indicated that varieties had differing abilities to prevent excessive bleeding from occurring.






Figure 2: Influence of variety of helmeted guinea fowl on hematological parameters

^{abc} columns with different superscripts differ significantly ($P = .05$)

4. CONCLUSION

In conclusion, we found that MCV and platelets significantly differed with sex while RBC, WBC, PCV, hemoglobin, MCHC, MCH, MCV and platelets differed with varieties of extensively reared guinea fowl during breeding season in northern Nigeria. Selection of *Numida meleagris* males with high RBC, hemoglobin and PCV for breeding may improve fertility due to improved semen quality, and androgen production. PCV values may be useful markers for predicting spermatozoa maturation in testes, sexual maturity and the onset of semen production. The WBC could serve as a marker indicating immunity and inherent resistance to tropical poultry diseases and parasites while hemoglobin and MCV values similar to or above (but within the reference range of avian) those obtained in this study could be suitable markers for evaluating and improving the selected population of helmeted guinea fowl for feed conversion efficiency and adequate oxygen supply for maintenance of body functions. PCV can serve as a marker for selection of helmeted guinea fowl for improved growth and productive life of the bird through improved weight gain and heat – tolerance.

COMPETING INTERESTS

There are no competing interests

REFERENCES

1. Dondofema FA. Survey of agricultural production and marketing in Nenyunga Communal lands with special emphasis on livestock. B.Sc Hons Dissertation, University of Zimbabwe, Harare, Zimbabwe. Pp30; 2000.
2. Smith J. Guinea fowl, Diversification Data Base, Scottish Agricultural College; 2000. Available <http://www.sac.ac.uk/management/external/diversification/Tableofcontents>.
3. Onyeausi BI. Serum Progesterone Concentrations in Indigenous Nigerian Guinea fowls. International Journal of Poultry Science. 2007; 6 (8): 608 – 609
4. Ibrahim AA, Aliyu J, Hassan AM 2011. Effects of Age and Sex on some Haematological parameters of Turkey (*Melegris gallopavo*) reared in the Semi – arid environment of northern Nigeria. Proceedings of 35th Annual Conference of Genetic Society of Nigeria. Held at Ahmadu Bello University, Zaria, Kaduna State on 10th – 13th October, 2011 .Pp 51 – 54
5. Harr KE. Clinical chemistry of companion avian species: A review. Veterinary Clinical Pathology. 2002; 31:140–151.

- 305 6. Hauptmanová K., Malý M, Literák I. Changes of haematological parameters in
306 common pheasant throughout the year. *Vet Med Czech*. 2006; 51: 29-34
- 307 7. Oyewale JO, Ogwuegbu SO. Hematological studies on the guinea fowl (*Numida*
308 *meleagris* Pallas). *Bulletin on Animal Health and Production in Africa*. 1986; 34, 61-
309 65.
- 310 8. Orji BI, Okeke GC, Akanyiba AO. Hematological studies on the guinea fowl (*Numida*
311 *meleagris* galeata Pallas). I. Effect of age sex and the time of bleeding on the
312 haematological values of guinea fowls. *Nigerian Journal of Animal Production*. 1986;
313 13: 94 - 99.
- 314 9. Oyewale JO. Some aspects of the haematology of the guinea fowl (*Numida*
315 *meleagris* galeata, Pallas) Ph.D Thesis. University of Ibadan, Ibadan; 1988.
- 316 10. Okoro VMO, Ogundu UE, Ogbuewu IP, Obikaonu H, Emenyonu C. Effect of sex and
317 systems of production on the haematological and serum biochemical characters of
318 helmeted guinea fowls (*Numida meleagris* pallas). *International Journal of*
319 *Biosciences*. 2011; 1(3): 51 -56.
- 320 11. Adedibu II, Ayorinde KL, Musa AA, Orunmuyi M, Kabir M. 2013a. Multifactorial
321 analyses of morphological traits of helmeted guinea fowls *Numidia meleagris*. Book
322 of abstracts of the 37th Annual conference of Genetics Society of Nigeria held at
323 Federal University, Lafia, Nassarawa state on the 22nd – 24th October, 2013. Pp 93 –
324 98.
- 325 12. Adedibu II, Ayorinde KL, Hassan MR, Owolabi J. Socio economic contribution of
326 helmeted guinea fowl in Northern Nigeria. *International Journal of Applied Research*
327 *and Technology*. 2013b; 2(5): 42 – 50.
- 328 13. Ayorinde KL, Song Y, Durmmond P, Smith EJ. Molecular genetic analysis of diversity
329 in village chickens in Nigeria. *Nigerian Journal of Animal Production*. 2001; 28(2): 128
330 – 134.
- 331 14. SAS Institute Inc. Statistical Analysis System User's Guide. SAS/STAT Version 9.0
332 for windows. SAS Institute Inc., Inc. Cary, North Carolina, USA; 2002.
- 333 15. Strakova E, Suchy P, Kabelova R, Vitula F, Herzig I. Values of selected
334 haematological indicators in six species of feathered game. *Acta Veterinaria*. Brno.
335 2010;79: S3 – S8: doi:10.2754/avb201079S9S003
- 336 16. Sturkie PD. *Avian Physiology*. Comstock Publishing Associates, New York. 24 -35;
337 1965
- 338 17. Coles BH. *Avian medicine and surgery*. Blackwell Science, Oxford, UK, pp. 58; 2005
- 339 18. Antonio Garcia MT, Hermosa Y, Aguirre JI. Does breeding status influence
340 haematology and blood biochemistry of yellow-legged gulls? *Acta Biologica*
341 *Hungarica* 2010; 61(4): 391– 400 DOI: 10.1556/ABiol.61.2010.4.3
- 342 19. Schmidt EM, Paulillo AC, Santin E, Dittrich RL, Oliveira EG. Hematological and
343 serum chemistry values for the Ring – necked pheasant (*Phasianus colchicus*):
344 variation with sex and age. *International Journal of Poultry Science*. 2007; 6(2): 137
345 – 139
- 346 20. Priya M, Gomathy VS. Haematological and blood biochemicals in male and female
347 turkeys of different age groups. *Tamilnadu Journal of Veterinary and Animal*
348 *Sciences*. 2008; 4(2): 60 – 68
- 349 21. Elagib HAA, Ahmed ADA. Comparative study on haematology values of blood of
350 indigenous chickens in Sudan. *Asian Journal of Poultry Science*. 2011; 5: 41 – 45.
351 Doi:10.3923/ajpsaj.2011.41.45
- 352 22. Oke UK., Herbert U, Akinmutimi AH. Early lay characteristics and haematology of
353 Pearl Guinea fowls as influenced by dietary proteinand energy levels. *International*
354 *Journal of Poultry Science*. 2003; 2(2): 125 – 132.
- 355 23. Oke UK., Herbert U, Akinmutimi AH. Early lay characteristics and haematology of
356 Pearl Guinea fowls as influenced by dietary proteinand energy levels. *International*
357 *Journal of Poultry Science*. 2003; 2(2): 125 – 132.

- 358 24. Herbert R, Nanney J, Spano JS, Pedersoli WM, Krista LM.. Erythrocyte distribution
359 in ducks. American Journal of Veterinary Research. 1989; 50: 958-960.
- 360 25. Elkomy AEA, El – ghalid OAH, Abd – El – Hady AM. Effect of gonadotrophin
361 stimulant (Theriogon) treatment on Alexandria cockerels reproductive performance.
362 Egyptian Poultry Science. 2009; 29(2): 609 – 621
- 363 26. Atwal OS. Mcfarland LZ, Wilson WO. Hematology of coturnix from birth to maturity.
364 Poultry Science. 1964; 43: 1392 – 1401
- 365 27. Bierer BW, Eleazer TH, Roebuck DE. Sedimentation rate, packed cell volume, buffy
366 coat volume and rectal temperature of Chickens and Turkeys at various ages.
367 Journal of the American Veterinary Medicine Association. 1964; 144: 727 – 730.
- 368 28. Helene CC, Murray RB. Correlation of Organ weights, hematocrit and testosterone
369 with sexual maturity of the male turkey. Poultry Science. 1990; 70: 1252 – 1257
- 370 29. OpenStax College. *Anatomy and Physiology*. OpenStax College. 2013. Retrieved
371 from the Connexions Website:<http://cnx.org/content/col11496/1.6/>
- 372 30. Mitruka BM, Rawnsley HM. Chemical, biochemical and haematological reference
373 value in normal experimental animals, Masson Dubling U.S.A. Pp 88 – 90; 1977
- 374 31. Isidahomen, C. E. & Njidda, A. A. Haematology and Carcass Characteristics of
375 Naked Neck, Frizzled and Normal Feathered Local Chickens in Southern Nigeria.
376 Savannah J. Agric.2012; 7: 12–19
- 377 32. Islam MS, Lucky NS, Islam MR, Ahad A, Das BR, Rahman MM, et al. Haematological
378 parameters of Fayomi Assil and local chickens reared in Saylhet Region in
379 Bangladesh. International.Journal of Poultry Science. 2004; 3: 144 – 147.
- 380 33. Kececi T, Col R. Haematological and biochemical indices of the blood of Pheasant
381 (*Phasianus colchicus*) of different ages. Turkish Journal of Veterinary and Animal
382 Science. 2010; 35(1): 149 – 156
- 383 34. Mmereole FUC. Haematological and serological profiles of the local and exotic
384 chickens in Southern Nigeria. Ph.D Thesis submitted to the Department of Animal
385 Science, Delta State University,Asaba, Delta State; 2004
- 386 35. Isidahomen EC, Ozoje MO, Njidda AA. Haematological and Serum Biochemical
387 Indices of Local and Exotic Chickens in a subtropical Enviroment. European Journal
388 of Biological Sciences. 2011; 3(1): 16 – 21.
- 389 36. Emenalom OO, Esonu BO, Etuk EB, Anaba C. Effect of *Mucuna puriens* (velvet
390 beans) leaf meal on performance and blood composition of finisher broiler chickens.
391 Nigerian Journal of Animal Production. 2009; 36(1): 52 – 60
- 392 37. Olayemi FO. Haematolgical changes in guinea fowls (*Numida meleagris* galeta,
393 Pallas) following haemorrhage. International Journal of Poultry Science. 2009; 8(11):
394 1093 – 1095.
- 395 38. El Safty SA, Ali UM, Fathi MM. Immunological Parameters and laying performance of
396 naked necke and normally feathered genotypes of chicken under winter conditions in
397 Egypt. International Journal of Poultry Science. 2006; 5:780 – 785