

Original Research Article

Qualitative characteristics and Suture Measurements of Two Breeds of Snail Rearing in Nigeria

Abstract

Heliculture is gradually gaining more attention in Nigeria. There are still constraint to snail production en mass which include breed and strain identification feeding, management and health. In this study, the Qualitative characteristics and suture measurement of two breed of snails were observed. A total of 90 snails comprising of 50 Archachatina marginata and 40 Achatina achatina were purchased at different markets in towns around Ayetoro. Most (64%) of the A. achatina had brown shell colour, a few (36%) had brown with light yellow shell colour. 100% of A. achatina had brown with light brown-yellow shell colour. 22% of A. marginata had black foot, 64% had brown and 14% had brown-black foot colour. 100% of A. achatina had brown-black foot colour. There is a significant difference ($P<0.01$) in the brown-black foot colour of A. achatina and A. marginata. 94% of A. marginata had broad and wide apex, 6% had pointed apex, (0%) narrow and wider apex while 100% A. achatina had narrow and wider apex. A. achatina had zig zag thick brown lines on the shell which was significantly different ($P<0.01$) from the A. marginata. The distance between the 1st and 2nd suture, 2nd and 3rd suture of A. marginata respectively (1.69 ± 0.08 , 0.79 ± 0.04) cm is longer than the respective lengths (1.59 ± 0.03 , 0.78 ± 0.02) cm of A. achatina. This distance in suture of the 1st and 2nd, 3rd and 4th, 4th and 5th of A. marginata is not significantly different ($P>0.01$) from A. achatina, the 2nd and 3rd suture is significantly different ($P<0.01$). Snail live weight is highly positively correlated with the distance between the 1st and 2nd suture (0.889) and the 2nd and 3rd suture (0.718) in A. marginata and A. achatina respectively. The SLW is determined more by the distance between the 4th and 5th suture in A. marginata.

Keywords: Heliculture, Qualitative Characters A. marginata, A. achatina, suture, broad and wide apex, pointed apex

Introduction

The importance of protein can not be over emphasized in the nutrition of man [7]. Animal protein is of high biological value and posses all the essential amino acids in desirable quantities [3]. The major sources of meat protein for the Nigerian populace comes mainly from poultry, beef, mutton while the protein of animal source in Nigeria is inadequate because the demand exceeds supply [7]. Some sources of animal protein include macro and micro livestock species [1]. Micro livestock are excellent and cheaper sources of animal proteins [3] [6]. Snail have been well known and highly appreciated as a valuable source of nutrition in many countries [4]. In ancient Rome, Snails were fattened up in

cochlear gardens before they are eaten [9]. In Nigeria, Snails constitute an important component of the food of numerous rural dwellers especially in the rainforest belt as well as other area of the country where snail eaten is not regarded as a taboo [4]. A type of snail known as prewinkle is being relished more among the Efik tribe in Nigeria in states like Akwa Ibom and Cross Rivers. In West Africa, the common breeds of snails reared are African giant land snail *Archachatina marginata*, *Achatina achatina*, *Achatina fulica*, *Limicolaria* species [5] [2]. The most popular edible snails in West Africa are the giant Snail *Achatina achatina* and the big black *Archachatina marginata* [1]. [7] noted that the foot (skin) of *A. marginata* can be black or white in colour. *A. marginata* has bulbous shell with brown stripes and a wide apex [4]. *A. achatina* has a brown shell with conspicuous zig zag streaks and narrow pointed apex [4]. The shell of the matured snail form elongated cone which coils spirally around a central axis usually 4-5 whorls at maturity [7]. In this study the qualitative characters and suture measurement of *Archachatina marginata*, and *Achatina achatina* were observed.

Materials and Method

Study Location

The experiment was carried out at the Teaching and Research farm of Olabisi Onabanjo University, Ayetoro in Ogun State. Ayetoro is located in latitude 7° 15' N and longitude 3° 31'E in a deciduous/ derived savannah zone of Ogun State. Climate is sub-humid tropical with an annual rainfall of 1,909.3mm. Raining season is between early April and late October, Rainfall pattern is bimodal with two peaks in June and September. Maximum temperature varies between 29°C during the peak of the wet season and 34°C at the onset of the wet season. Mean annual relative humidity is 81% (Onakomaiya et al, 1992).

Measurement of Characteristics

Two breeds of Snails were used in this study. Fifty *Archachatina marginata* and forty *Achatina achatina*. Animals were purchased at different markets in towns around Ayetoro.

Animals were identified with numbers using paint to write on the shell. Snails were numbered 1-90. Suture was measured using the vernier caliper. A scout pro sensitive scale (0.600g) was used to take the weight. Data collected include shell colour (SC), foot colour (FC), shape of Apex Shell (SAS). Types of stripes on shell (TSS), Distance between first and second suture (DBS1 & 2), second and third suture (DBS2 & 3), third and forth suture (DBS3 & 4), forth and fifth suture (DBS 4 & 5) and snail have weight (SLW).

Statistical Analysis

Data collected were analysed using statistical package for social scientist 2006; which include chi square, descriptive statistics and T-test of means, correlation and regression analysis.

Results and Discussion

36% of *A. marginata* snail had brown with light yellow shell colour and 64% had brown shell colour while 100% *A. achatina* had brown with light brown yellow shell colour. The brown with light brown yellow colour of *A. achatina* is significantly different from the brown with light brown yellow colour of *A. marginata* at $P < 0.01$ (Table 1 & 2). The result agrees with Akinnusi (2004). In *A. achatina*, it was observed during this study that there is a red colouration on the aperture shell. 22% of *A. marginata* had black foot colour, 64% had brown and 14% had brown black while 100% of *A. achatina* had brown-black foot colour. The brown-black foot colour of *A. achatina* is significantly different from the brown-black foot of *A. marginata* at $P < 0.01$ (Table 1 & 2). The result agrees with the statement of Akinnusi (2004), that the foot colour of *A. marginata* snail is usually brown black which is different from *A. Achatina*. 94% of *A. marginata* had broad-wider apex, 6% had pointed apex, there is no (0%) narrow-wider apex while 100% of *A. achatina* had narrow and wider apex.

The narrow and wider apex of *A. achatina* is highly significantly different from the broad and wider apex of *A. marginata* ($P < 0.01$). This result agrees with Akinusi (2004). 80% of *A. marginata* snail had straight lines with brown lines on shell, 20% had spotted stripes with straight stripes while 100% of *A. achatina* had zig zag thick brown lines. The zig zag thick brown lines on shell of *A. achatina* is highly significantly different from the straight

lines with thick brown lines of *A. marginata*. This result agrees with Akinnusi (2004). In this study the result revealed that some of the *A. marginata* snail had spotted stripes with straight stripes on the shell, this could be due to the fact that random mating has occurred where these animals have had equal chance of mating with each other in the various habitat they were collected from. In the *A. marginata* snail, 5(10%) had 2+ 1 suture, 17 (34%) 3 + 1, 2(4%) 3 + 2, 1 (2%) 3 + 3, 21 (42%) 4 +1 and 4 (8%) with 4 + 2 while in the *A. achatina*, there is 0 (0%) 2 +1 suture, 2 (5%) had 3 + 1, 14 (40%) 3 + 2, none (0%) 3 + 3, 7 (17.5%) 4 + 1 and 15 (37.5%) 4 + 2 (Table 3). In the 3 + 1 suture animals, the mean distance between the second to fourth sutures ($0.67 \pm 0.07\text{cm}$, $0.26 \pm 0.03\text{cm}$) in *A. achatina* is longer than those of *A. marginata* (0.70cm , 0.03 cm) but the first and second suture is longer in *A. marginata* ($1.49 \pm 0.09 \text{ cm}$) than (1.4cm) *A. achatina*. In the 3 + 2 suture animals, the mean distance between sutures in *A. Achatina* snail (1.48 ± 0.03 ; 0.69 ± 0.03 ; 0.27 ± 0.02) cm is longer than (0.95 ± 0.05 , 0.35 ± 0.05 , 0.10 ± 0.00) cm *A. marginata*. In animals with 4 + 1 and 4 + 2 suture, the mean DBS of *A. marginata* snail is longer than the *A. achatina* snail (Table 3). The fact that within the 3 + 1 suture and 3 + 2 suture, the suture of *A. achatina* snail is bigger than *A. marginata* snail may be due to age difference between the animals. In this research, it was observed that the mean DBS decreases towards the apex suture. Among the breeds, the distance between first and second suture, second and third suture of *A. marginata* respectively (1.69 ± 0.08 , 0.79 ± 0.04) cm is longer than (1.59 ± 0.03 , 0.78 ± 0.02) cm that of *A. achatina* (Table 4). It was observed in this study that the longer the distance between the suture the more conspicuous the stripes and colour on that specific suture. The mean DBS of the first and second, third and fourth, fourth and fifth sutures of *A. marginata* is not significantly different from *A. achatina* at $P>0.01$. The mean of the distance between the second and third suture of *A. marginata* snail is significantly different from *A. achatina* snail at $P<0.01$ (Table 5). The SLW is highly positively correlated with Distance between 1st and 2nd suture (0.889), 3rd and 4th (0.879), 2nd and 3rd (0.851), 4th and 5th (0.846) sutures in *A. marginata* and 2nd and 3rd suture (0.718), 1st and 2nd sutures (0.697) in *A. achatina* (Table 6). The parameter that could most significantly ($P<0.01$) determine SLW is the distance between 4th and 5th suture in *A. marginata*. The result was negative and highly not significant $P>0.01$) for the

132 distance between sutures in *A. achatina* although the distance between the 3rd and 4th suture
133 could still accurately determine SLW (Table 7).

134

135 **Conclusion and Application**

136 This study has shed more light on the Mendelian (Qualitative) characters of *A. marginata*
137 and *A. achatina*. A strain of *A. marginata* has been identified in this study. The longer the
138 distance between the suture, the more conspicuous the stripes and colour on that specific
139 suture. The snail live weight is highly positively correlated with the distance between 1st
140 and 2nd suture, 2nd and 3rd suture in *A. marginata* and *A. achatina*. The prediction power is
141 high for the distance between 4th and 5th suture in *A. marginata*.

142

143

144

145 **References**

- 146 1. Ajetunmobi, A. and Olayemi, A. (2002). Animal Agriculture, monogastric and
 147 mini livestock farming, Ade positive horizon associates. Alografiks
 148 Communication Company, Pp2-99.
- 149
- 150 2. Akinnusi, O. (1997). Snail farming, Low investimet, high profit business livestock.
 151 Echo. April – June, 1997. Pp. 14-16.
- 152
- 153 3. Akinnusi, O. (1998). A practical approach to background Snail farming. Nig. J.
 154 Anim. Prod. vol.55, Pp 193-197.
- 155
- 156 4. Akinnusi, O. (2004). Introduction to Snail and Snail Farming. Triolas exquisite
 157 ventures Publisher. Abeokuta, Nigeria, Edition 111. Pp1-44.
- 158
- 159 5. Food and agriculture Organization (1986). Farming Snails by FAO. Better
 160 Farming Series 3/33. Rome. Italy.
- 161
- 162 6. National Research Council (1991). Macro livestock. Little known small animals
 163 with a promising economic future. National Economic press, Washington D. C.
- 164
- 165 7. Omole, A. J. , Taiwo, A. A. and Amusan, J. A. (2007). Technical Guide/Bulletin.
 166 Practical Snail Farming Ibadan, Nigeria. Institute of Agricultural Research and
 167 Training, Moor Plantation.
- 168
- 169 8. Onakomaiya, S. O., Oyesiku, K. A and Jegede, S. J. (1992): Ogun State in Maps.
 170 Rex Charles Publications, Ibadan Pp.7-8.
- 171
- 172 9. Thompson, R. and Cheney, S. (1996). Raising Snail. The alternative farming
 173 system information centre, National Agricultural Library.
- 174

175

176 **Table 1: QUALITATIVE CHARACTERS OF *Archachatina marginata* and**
 177 ***Archatina achatina***

CHARACTERS	A.M (%)	A.A (%)
SHELL COLOUR		
Brown with light brown yellow	n 18 36.0	40 100
Brown	n 32 64.0	0 0
Total	n 50	40
FOOT COLOUR		
Black	n 11 22.0	0 0.0
Brown	n 32 64.0	0 0.0
Brown black	n 7 14.0	40 100.0
Total	n 50 100.0	40 100.0
SHAPE OF SHELL		
Broad-wider apex shell	n 47 94.0	0 0
Narrow-wider apex shell	n 0 0.0	40 100.0
Pointed apex shell	n 3 6.0	0 0
Total	n 50 100	40 100
TYPE OF SHAPES ON SHELL		
Straight lines with thick brown lines on shell	n 40 80	0 0
Spotted stripes with straight stripes on shell	n 10 20	0 0
Ziz zag thick brown lines on shell	n 0 0	40 100.0
Total	n 80 100.0	40 100.0

178

179 **Table 2: Chi square for the qualitative characters**

VARIABLES	χ^2	<i>df</i>	Sig
SHELL COLOUR	39.724	1	0.000
FOOT COLOUR	84.072	2	0.000
SHAPE OF SHELL	90.000	2	0.000
TYPES OF STRIPES	90.000	2	0.000
ON SHELL			

180

181 *Chi square – Pearson Chi square*

182 *df – Degree of freedom*

183 *sig – significance*

184

185

186

187

188

189

190 **Table 3: Individual whorl population number and individual distance between sutures of archachatina marginata and**
 191 **Achatina achatina**

No. of sutures		D.B.S 1&2	D.B.S 2&3	D.B.S 3&4	D.B.S 4&5	SLW	D.B.S 1&2		D.B.S 2&3	D.B.S 3&4	D.B.S 4&5	Total	SLW	
2+1	n	5	5	0	0	0	N	0	0	0	0	5	45.900	
	%	10	10				%							
	X	0.660	0.320			17.460	X							
	S.E	0.0510	0.583			3.7993	S.E							
3+1	n	17	17	17			N	2	2	2	0	19		
	%	34	34	34			%	5	5	5	0+			
	X	1.485	0.674	0.262		69.565	X	1.400	0.700	0.300				
	S.E	0.0862	0.0725	0.0259		5.9332	S.E							
3+2	n	2	2	2	0		N	16	16	16	0	18		
	%	4	4	4	0		%	40	40	40	0			
	X	0.950	0.350	0.100		19.100	X	1.481	0.688	0.269				
	S.E	0.0500	0.0500	0.000		1.2000	S.E	0.0332	0.0287	0.0218				
3+3	n	1	1	1	1		n					50.986 3.5163 77.163 7.7347		
	%	2	2	2	2		%							
	X	1.000	0.500	0.200	0.200	20.900	X							
	S.E						S.E							
4+1	n	21	21	21	21		n	7	7	7	7			28
	%	42	42	42	42		%	17.5	17.5	17.5	17.5			
	X	2.167	1.029	0.476	0.148	142.771	X	1.586	0.771	0.343	0.100			
	S.E	0.0433	0.0332	0.0206	0.0206	9.9099	S.E	0.0459	0.0360	0.0202	0.0000			
4+2	n	4	4	4	4		n	15	15	15	15			19
	%	8	8	8	8		%	37.5	37.5	37.5	37.5			
	X	1.975	0.938	0.488	0.163	132.900	X	1.719	0.869	0.381	0.119			
	S.E	0.1250	0.0375	0.0125	0.0239	11.6427	S.E	0.0600	0.0405	0.0164	0.0101			
Total	n	50	50	50	50		n	40	40	40	40			

* +1, +2,+3 are the tiny sutures

194 **Table 4: Mean of Distance between sutures**

VARIABLES	A.M	A.A
Distance between 1 st and 2 nd suture	50	40
	1.693	1.593
S.E	0.0801	0.0331
Distance between 2 nd and 3 rd suture	50	40
	0.792	0.775
S.E	0.0455	0.0242
Distance between 3 rd and 4 th suture	50	40
	0.336	0.328
S.E	0.0260	0.0139
Distance between 4 th and 5 th suture	50	40
	0.075	0.065
	0.0121	0.0098

195

196

197

198 **Table 5: Test of Significance between mean of distance between Suture of**
 199 ***Archachatina marginata* and *Achatina achatina***

VARIABLES	<i>Levens test of sig</i>	<i>Variance assumption</i>	T-test of Sig
D.B 1 st and 2 nd suture	0.000	N.E.V	0.232
D.B 2 nd and 3 rd suture	0.000	N.E.V	0.742
D.B 3 rd and 4 th suture	0.000	N.E.V	0.774
D.B 4 th and 5 th suture	0.010	N.E.V	0.524

200

201

202

203

204

205

206 **Table 6: CORRELATION ANALYSIS OF *Archachatina marginata* and *Achatina***
 207 ***achatina***

	SLW	DBS1&2	DBS2&3	DBS3&4	DBS4&5
SLW		0.697**	0.718**	0.460**	0.557**
DBS	0.889**		0.882**	0.710**	0.629**
1&2					
DBS	0.851**	0.887**		0.721**	0.632**
2&3					
DBS	0.879**	0.933**	0.867**		0.604**
3&4					
DBS	0.846**	0.751**	0.706**	0.811**	
4&5					

208

209

210

211


212 **Table 7: REGRESSION ANALYSIS OF *Archachatina marginata* and *Achatina***
 213 ***achatina***

Distance between suture		B	S.E	Sig
Distance between 1 st and 2 nd suture	A.m	11.532	14.620	0.436
	A.a	-9.715	11.018	0.387
Distance between 2 nd and 3 rd suture	A.m	5.724	12.373	0.647
	A.a	-7.042	15.153	0.647
Distance between 3 rd and 4 th suture	A.m	-40.576	39.547	0.313
	A.a	-1.585	14.129	0.912
Distance between 4 th and 5 th suture	A.m	138.400	45.275	0.005
	A.a	-8.451	17.375	0.632

214

215

216



230

231

232 **Fig. 1:** The diagram of the snail showing the whorl, spire, aperture and distance
233 between suture

234