

Composition and dynamics of some nutrient levels in Jabi dam, Lower Usuma dam and Gwagwalada river in Federal Capital Territory, Abuja, Nigeria.

ABSTRACT

Jabi dam, Lower Usuma dam and Gwagwalada river in the Federal Capital Territory were studied for a period of eighteen months (June 2005 – January 2007) to determine their nutrient composition and dynamics. Samples were collected from these water bodies on monthly basis. The concentration of some nutrient parameters such as Phosphate-phosphorus ($\text{PO}_4\text{-P}$), Nitrate-Nitrogen ($\text{NO}_4\text{-N}$), Total Dissolved Solids (TDS), Calcium (Ca), Potassium (K) and Sodium (Na) were assessed and determined. The results obtained were statistically analysed to show any relationship amongst them. The mean values for $\text{NO}_4\text{-N}$, Ca and K were not significantly different ($P > 0.05$) between the three water bodies. While the mean value for $\text{PO}_4\text{-P}$ were significantly different ($P < 0.05$), with Jabi dam having the highest mean value of 11.61 mg/l, followed by Gwagwalada river, 9.96 mg/l and Lower Usuma dam had least mean value of 2.46 mg/l. Jabi dam, Lower Usuma dam and Gwagwalada river had mean TDS of 95.0 mg/l, 88.5 mg/l and 104.5 mg/l respectively, Gwagwalada river being significantly different ($P < 0.05$) from Jabi and Lower Usuma dams. Similarly, Jabi dam, Lower Usuma dam and Gwagwalada river had mean Sodium values of 2.42 mg/l, 2.85 mg/l and 3.73 mg/l respectively, Gwagwalada river being also significantly different ($P < 0.05$) from Jabi and Lower Usuma dams. $\text{PO}_4\text{-P}$ and $\text{NO}_4\text{-N}$ showed no significant ($P > 0.05$) monthly variations in the three water bodies. However, TDS, Ca, K, and Na showed significant ($P < 0.05$) monthly variations in the three water bodies. TDS, Ca, K, and Na had all their wet season mean values higher than their dry season mean values in all the three water bodies studied, while $\text{PO}_4\text{-P}$ and $\text{NO}_4\text{-N}$ had their wet season mean values lower than the dry season mean values in the three water bodies. The study revealed that the nutrient parameters measured were slightly lower than the World Health Organization (WHO) recommended values for aquatic life.

Keywords: Jabi dam, Lower Usuma dam, Gwagwalada river, Nutrients levels and variations.

INTRODUCTION

The trends of the study of water resources in the time past has been tailored towards the engineering aspects (e.g. generation of electricity and irrigation purposes), and also its specific need i.e. providing potable drinking water for man. However, only little work has been done in the ecological aspect of these water bodies. This neglect has generated threats to the organisms in such water body and a potential public health hazard to the user of such water resources and its ecosystem. Water is fundamental to life. It is capable of dissolving many substances including nutrients more than any other known solvent; hence it is called the universal solvent.

Increase in the rate of industrialization and consequently urbanization always exacerbates its effect on the environment (Abumere, 1994, Asonye *et al*, 2007, Balogun *et al*, 2011). It is a known fact that inadequate sanitary facilities in most rural and urban settlements have turned many surface waters into open sewers, thus most urban rivers and streams are dumping grounds for human wastes as found in Rivers Kaduna, Niger, Ogunpa, Ogun and Chanchaga (Ojutiku and Kolo 2008).

Nutrients such as phosphorus and nitrogen are essential for the growth of algae and other plants. Aquatic life is dependent upon these photo synthesizers, which usually occur in shallow surface water (Nortcliff, 2012). Excessive concentration of nutrients however can over-stimulate aquatic plants and algae growth thus affecting water quality and primary productivity potentials (Silva and Ronal, 1987; Adakole *et al* 2004). Over availability of nutrients in a stagnant environment can lead to consequent effect of eutrophication and its attendant consequences (Fagade *et al*, 1993; Mitchel and McDonald, 1995).

Nutrient balances which consider the inputs and outputs from the system have been used to estimate the magnitude and extent of nutrient mining. Inputs are fertilizers, organic residues, manures, nitrogen fixation and sediments. Outputs are through uptake and export, erosion, leaching and volatilization (Nortcliff, 2012). Also in his study of soil and food security he observed nutrient mining as a threat impacting on food security strongly in Africa where insufficient nutrients are returned to the soil after crop production.

Abuja as a growing city and also being the Federal Capital Territory of Nigeria with its growing population and increased demand of water as a resource and its other

potentials calls for study and research work that will help guide and give a baseline data on the primary productivity potentials of some of the water bodies in Abuja, thus the objective of research is to assess the nutrient composition and dynamics of Jabi dam, Lower Usuma dam and Gwagwalada river all within the Federal Capital Territory of Nigeria.

The water nutrient parameters assessed in this study include Nitrate-Nitrogen, Phosphate-Phosphorus, Total dissolved solids, Calcium, Potassium and Sodium.

MATERIALS AND METHODS

The Study Area

The Nigerian Federal Capital Territory (FCT) was established by decree No. 6 of 1976 under the Federal Capital Development Authority (FCDA). Abuja itself has been said to be the fastest growing city in Africa. Growth in this sense encompasses both the tangible and intangible developmental activities going on in the FCT, including huge construction activities, as well as other economic activities.

The detailed plan of the research involved monthly sampling for the water nutrient parameters at Jabi dam, Lower Usuma dam and Gwagwalada River, The sampling period spanned between June 2005 and January 2007.


Station 1 is the Jabi dam reservoir which used to supply treated water to Jabi village and it's environ but now as a recreation center.

Station 2 is the Lower Usuma dam reservoir which is 24 kilometers to the Central area of Abuja and supply treated water to the whole of FCT.

Station 3 is the Gwagwalada River at the mini water treatment plant constructed at Gwagwalada River. The sampling site is near the Gwagwalada township bridge.

Abuja F.C.T.

(Major Rivers & Streams)



0 20 km
Scale

LEGEND	
	Study Boundary
	Rivers
	Gwagwalada River
	Dams

MAP OF THE STUDY AREA SHOWING SAMPLE STATIONS

Determination of nutrient elements

Calcium (Ca) was determined by Atomic Absorption Spectrometer (AAS) using a computer controlled SOLAAR 969 Atomic Absorption Spectrometer using nitrous oxide-acetylene flame. The potassium and sodium levels were determined by flame photometer. Phosphate-Phosphorus was determined using the Ascorbic acid method, Nitrate-Nitrogen was determined using the phenoldisulphuric acid and Total dissolved solids all as described in standard methods (APHA, 1995).

RESULTS

The results of the concentration of nutrients and cations measured at Jabi dam, Lower Usuma dam and Gwagwalada river and their monthly variations are shown in Tables 1- 3 and figures 1 – 6 respectively.

The monthly variations of phosphate-phosphorus for the three water bodies showed sharp contrasts and fluctuations (fig 1). The differences are however significant ($P < 0.05$) and Jabi dam had the highest mean value of 16.38 mg/l followed by Gwagwalada river 9.96 mg/l and Lower Usuma dam 2.46 mg/l (Table 1). In the seasonal variations the dry season mean values were higher than the wet season mean values (Table 3).

The monthly variations of nitrate-nitrogen concentration in the different water bodies showed wide fluctuations (Fig.2). There were no significant differences ($P > 0.05$) in nitrate concentrations in the three water bodies, however, Lower Usuma dam had the highest mean value of 4.30 mg/l, while Jabi dam had the least mean value of 3.25 mg/l (Table 1). The seasonal variations showed no sharp distinction between the wet and dry seasons (Table 3).

Gwagwalada river had the highest total dissolved solids mean value of 104.5 mg/l followed by Jabi dam, which had a value of 95.0 mg/l, while Lower Usuma dam recorded the least mean value of 88.5 mg/l (Table 1). There were significant differences ($P < 0.05$) (Table 2) in the monthly variations of total dissolved solids (fig.3).

Jabi dam had highest calcium mean value of 28.39 mg/l, while Gwagwalada river and Lower Usuma dam had 27.86 mg/l and 27.48 mg/l respectively (Table 1). There were monthly variations (fig.4), which were significant at ($P < 0.05$) (Table 2).

The monthly mean concentration of potassium in the three water bodies shows fluctuations. The lowest concentration of potassium was obtained in lower Usuma dam in June and July (fig 5). There were no significant variations ($P>0.05$) in the potassium concentrations in the three water bodies, though Gwagwalada river had the highest potassium mean value of 4.64 mg/l while Jabi dam had the least mean value of 3.64 mg/l. (Table 1).

The monthly variations of sodium concentration showed that the three water bodies had the lowest values in July (fig. 6). Gwagwalada River showed the highest mean value (3.73 mg/l) which was significantly different ($P<0.05$) from Jabi and Lower usuma dams which had 2.42 mg/l and 2.85 mg/l respectively (Table 1).

In all the three water bodies studied, TDS, Ca, K, and Na had their wet season mean values higher than the dry season mean values (Table 3).

Table 1: Mean concentrations of some nutrient elements and cations measured at Jabi dam, Lower Usuma dam and Gwagwalada river in FCT, Abuja.

Parameters	Jabi dam	Usuma dam	Gwagwalada river.	SE
Phosphate (mg/l)	11.61 ^c ± 19.73	2.46 ^{ab} ± 3.21	9.96 ^b ± 13.31	2.27
Nitrate (mg/l)	3.25 ^a ± 6.59	4.30 ^a ± 6.72	3.99 ^a ± 4.23	0.44
TDS (mg/l)	95.0 ^a ± 135.1	88.5 ^a ± 125.5	104.5 ^{ab} ± 152.2	3.31
Calcium	28.39 ^a ± 26.15	27.48 ^a ± 27.25	27.86 ^a ± 27.60	4.53
Potassium	3.64 ^a ± 1.78	4.19 ^a ± 3.10	4.64 ^a ± 2.46	0.37
Sodium	2.42 ^a ± 0.84	2.85 ^a ± 1.12	3.725 ^{ab} ± 1.52	0.20

Mean values on the same row carrying the same superscript are not significantly different at $P>0.05$

168 Table 2: Monthly concentrations (mg/l) in the nutrient elements and cations in Jabi dam, Lower Usuma dam and Gwagwalada river in
 169 FCT, Abuja.

Parameter	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
PO ₄ -P (mg/l)	0.73 ^a ±0.15	1.10 ^a ±0.35	1.53 ^a ±0.66	1.33 ^a ±0.49	1.98 ^a ±2.15	12.76 ^a ±19.31	12.29 ^a ±14.40	14.99 ^a ±23.86	6.63 ^a ±7.49	3.55 ^a ±4.16	1.92 ^a ±3.39	1.92 ^a ±3.35
NO ₄ -N (mg/l)	3.52 ^a ±5.43	1.74 ^a ±2.44	0.21 ^a ±0.07	0.97 ^a ±1.19	0.33 ^a ±0.31	3.14 ^a ±3.95	2.07 ^a ±3.34	3.41 ^a ±3.43	3.02 ^a ±4.83	7.94 ^a ±7.7	8.08 ^a ±7.46	5.27 ^a ±6.0
TDS(NTU)	3.15 ^a ±0.72	256.00 ^{bc} ±15.06	270.00 ^c ±12.30	275.00 ^c ±18.29	261.50 ^{bc} ±19.84	1.78 ^a ±1.07	1.86 ^a ±0.97	1.81 ^a ±1.05	1.85 ^a ±1.01	141.05 ^b ±148.06	159.56 ^{bc} ±174.66	2.85 ^a ±0.65
Ca	42.69 ^a ±12.83	31.44 ^{ab} ± 6.37	23.18 ^{ab} ±5.34	28.06 ^{ab} ± 3.47	39.61 ^{ab} ± 2.49	16.08 ^{ab} ±15.69	4.55 ^b ±2.64	14.26 ^{bc} ± 9.27	16.77 ^b ± 6.92	30.26 ^b ± 5.03	59.9 ^c ±59.28	41.31 ^{bc} ±11.41
K	4.30 ^a ±1.63	7.85 ^b ±3.17	3.80 ^a ±2.07	5.93 ^{ab} ±2.41	3.41 ^{ab} ±0.95	3.23 ^{ab} ±1.52	1.76 ^a ±0.52	4.09 ^{ab} ±2.94	3.71 ^{ab} ±2.42	3.59 ^{ab} ±2.09	5.11 ^{ab} ±1.08	3.00 ^{ab} ±1.11
Na	5.01 ^a ±1.34	3.65 ^{ab} ±0.8	3.07 ^{ab} ±0.97	3.35 ^{ab} ±0.84	4.21 ^{ab} ±1.02	2.41 ^b ±0.95	1.68 ^b ±0.50	2.89 ^b ±1.61	3.10 ^b ±1.62	2.76 ^b ±1.40	2.88 ^{ab} ±0.81	4.16 ^{ab} ±1.31

170 Data on the same row carrying the same superscript are not significantly different at P>0.05

Table 3: Seasonal mean values of nutrient elements and Cations measured at Jabi dam, Lower Usuma dam and Gwagwalada river in FCT, Abuja.

Parameter	Seasons	Jabi dam	Usuma dam	Gwagwaladariver
PO ₄ -P (mg/l)	Wet	1.33	0.68	0.76
	Dry	16.38	2.79	13.95
NO ₄ -N (mg/l)	Wet	1.63	4.85	2.86
	Dry	1.33	8.43	3.37
TDS (mg/l)	Wet	161.52	153.43	180.12
	Dry	71.52	65.95	72.11
Ca (mg/l)	Wet	37.26	40.74	39.62
	Dry	22.06	19.19	20.30
K (mg/l)	Wet	3.79	6.58	5.83
	Dry	3.31	2.79	3.9
Na (mg/l)	Wet	2.65	3.37	4.35
	Dry	2.30	2.64	3.52

173

174

175

176

177

Table 4 Comparison of WHO recommended nutrient Content and cations for Aquatic life and Research Findings.

Parameter	WHO Recommended value (mg/l)	Research Findings
Phosphate-Phosphorus (mg/l)	-	2.11–11.61
Nitrate-Nitrogen (mg/l)	-	0.97– 8.08
Total Dissolved Solids (TDS)	-	86.3 – 104.3
Calcium	1000	29.7 - 34.00
Potassium	10	3.5 - 5.8
Sodium	10	1.00 - 0.01

187

188

189

190

191

192

DISCUSSION

The higher dry season mean values of phosphate in all the water bodies studied is similar to the observed phenomenon by Ekom *et al* (1993) in some rivers in Cross-river State of Eastern Nigeria, where he observed higher dry season mean of phosphate. Ojutiku and Kolo (2008) also observed same in Chanchaga river. The higher dry season mean values observed in all the stations could be attributed to concentration effect as a result of reduced water volume; while the low wet season mean values recorded in all the stations could be due to run-off from the farmlands during the wet season into the dams which has dilution effect and also could be due to co-precipitation of dissolved phosphate with carbonates (CaCO_3) a phenomena which sometimes occurs in lakes (Kolo and Oladimeji, 2004). The higher mean phosphate value in Jabi dam and Gwagwalada river than Lower Usuma dam might be due to its shallow and lentic nature coupled with dry season irrigation farming activities. Adakole *et al*, (2008) also observed farming activities to contribute to phosphate value in Kubani dam. The phosphate levels recorded ranged from 2.11mg/l – 11.61.00mg/l. This is within the tolerable limit of 5,000 $\mu\text{g/l}$ recommended by WHO (1993)

The effluent discharges of fertilizer application to farm lands which washes into the water bodies may be responsible for the high concentration of nitrate in soil and water. The higher dry season mean value observed in the water bodies could be due to concentration effect during the dry season as a result of reduced water volume. There were no significant difference ($P>0.05$) between the nitrate values in the three water bodies studied. The higher nitrate mean value in Lower Usuma dam suggests autochthonous origin as a result of mineralization and also might be due to large catchment area. The range value of 0.2mg/l to 13mg/l observed during the study period is relatively low compared 24.46 mg/l to 56.60 mg/l recorded by Kolo *et al*, (2009) in Dan-Zaria dam. This might be because of the shallow nature of Dan-Zaria dam coupled with farming activities around the dam and constant visitation of the dam by cattle for the purpose of drinking which in the process defecates and urinates into the water.

Higher total dissolved solids observed in Gwagwalada river could be due to erosion of sand particles as a result of swift water movement which characterizes lotic water bodies. The relatively high level at Jabi dam might be due to construction activities that encourages movement of dissolved solids in to the dam during raining season as a result of run offs.

Similarly, the higher wet season mean values in all the water bodies obviously could be as a result of run off prevalent at that season which bring into the water bodies allochthonous materials. Contrary to Adakole *et al*, (2008), the lower wet season mean value observed in man-made lake was attributed probably to dilution effect.

The range of 29.7 - 34.0 mg/l of Calcium mean value recorded in all the stations fall far below the recommended value of 1000mg/l for drinking water for the livestock (WHO, 1993). However the above range is higher than the range of 2.2 to 18.3 mg/l recorded by Ojutiku and Kolo (2008) in River Chanchaga and Minna portable water. The higher wet season mean value observed in all the water bodies could be as a result of rains and flood water influx.

The potassium range of 3.5 - 5.8 mg/l compares well with the range recorded in River Chanchaga and Minna portable water by Ojutiku and Kolo (2008). The relatively higher value of potassium recorded in Gwagwalada river though not significantly different ($P>0.05$) from other water bodies could be due to its lotic nature. Gwagwalada River could be as a result of peasant and dry season farming with their application of potassium based fertilizers.

The none significant difference ($P>0.05$) of sodium in Jabi dam and Lower Usuma dam could be because of their lacustrine nature while the higher mean value recorded in Gwagwalada river might be due to washing activities around the river with sodium based detergent.

REFERENCES

- Abumere, S (1994) "Urbanization in Nigeria" In J.D. Tarver (ed) "Urbanization in Africa", NewYork, Greendwood press, 1994.
- Adakole, J.A., Abulode, D.S. and Balarabe, M.L (2008). Assessment of water quality of a man-made lake in Zaria, Nigeria. In proceedings of Taal2007: The 12th World Lake Conference: 1373 – 1382.
- Adakole, J.A., Mbah, C.E., Odeje, S.C. and Balarabe, M.L (2004). Primary productivity in relation to water quality in two man-made lakes in Zaria, Nigeria. Nigerian Society for Experimental Biology Journal. 4(1): 37 – 42.
- American public Health Association (APHA) (1995). Standard methods for the examination of water and waste water. 4th edition AHWa - WPUC Washington D.C.PP 1007 – 1151.
- Asonye, C.C, Okolie N.P.,Okenwa,E.E.and Iwanyanwu, U.G. (2007) Some physic-chemical

- characteristics and heavy metal profiles of Nigerian rivers, streams and waterways. *African Journal of Biotechnology*. 6(5): 617 – 624.
- Balogun, K.J.; Ladigbolu, I.A. and Ariyo, A.A (2011) Ecological assessment of a coastal shallow lagoon in Lagos, Nigeria: A bio-indicator approach. *Journal of Applied Sciences and Environmental management* 15 (1) 41 – 46.
- Ekom, R.; Akpan and John, O. Offem (1993). Seasonal variation in water quality of the Cross River, Nigeria. *Rev. Hydrobiologia, Tropical*. 26(2)95-103.
- Fagade, S.O., Adebisi, A.A, Kolo, R.J., Elemi, B.F., Adeosun, A.A., and Ja’afu, A. (1993) Urbanisation and degradation of aquatic resources: The Ibadan experience. Proceedings of the National conference on Conservation of Aquatic Resources. (ed) Egborge, A.B.M.; Omoloyin, O.J.; Olojede, A and Manu, S.B. Natural Resources Conservation Council, pp208 – 219.
- Kolo, R.J.; Gashau, M. and Ojutiku R.O. (2009). Water quality characteristics of Dan-Zaria dam, Niger State, Nigeria. *Continental Journal of Fisheries and Aquatic Science* 3: 56 – 65.
- Kolo, R.J. and Oladimeji, A.A. (2004). Water quality and some nutrient levels in Shiroro lake, Niger State, Nigeria. *Journal of Aquatic Sciences* 19(2):99-106.
- Mitchell, C. and McDonald, A.T. (1995). Catchment characterization as a tool for upland water quality management. *Journal of Environmental management*.44:83-95.
- Nortcliff, Stephen (2012) Soil and food security. Nigerian Journal of Technological Research. Vol. 7: 1 -12.
- Ojutiku, R.O. and Kolo, R.J. (2008). Water quality and some nutrient levels in River Chanchaga and Portable water, Niger State, Nigeria. *International Journal of Bioscience*. 3(2):12–21.
- Silva, E.I.L. and Ronald, W.D. (1987). The seasonality of monsoonal primary productivity in Sri-Lanka; *Hydrobiologia*, 150:165-175.
- Thornton. J.A (1986) Nutrient in African lake ecosystems: Do we know all? *Journal of Limnological Society of Southern Africa*. 12: 6 – 21.
- WHO, (1993) International Standard for Drinking water. 4th Ed. World Health Organization: <http://www.lenntech.com/applications/drinking/standards/who’s drinking-water-standards.htm>.

306

307




Figure 1: Monthly variations in phosphate of 3 contrasting water bodies in Abuja


321

322

323

324

325



341

Figure 2: Monthly variations in nitrate of 3 contrasting water bodies in Abuja




Figure 2: Monthly variations in Total Dissolved Solid of 3 contrasting water bodies in Abuja




Figure 4: Monthly variations in Calcium of 3 contrasting water bodies in Abuja




Figure 5: Monthly variations in Potassium of 3 contrasting water bodies in Abuja

369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397




Figure 6: Monthly variations in Sodium of 3 contrasting water bodies in Abuja

398
399
400
401
402
403
404