

Original Research Article**COLLOIDAL STABILITY AND POTENTIAL STRUCTURAL DEFORMATION INDEX OF
FOUR NIGERIAN SOILS****ABSTRACT**

This study was undertaken to evaluate the effect of continuous cultivation on the colloidal stability and potential structural deformation index (PSDI) of four Nigerian soils. The four soils used for the study were; Entisol from Nsukka Hill, Ultisol from Nsukka poultry site, Inceptisol from Eha-Amufu and Inceptisol from Ikem, in Nsukka area of south eastern, Nigeria. The land use types considered were fallow and cultivated. The soils collected from 0-25cm depth were separated into five aggregate fractions, 5-2 mm, 2-1 mm, 1-0.5 mm, 0.5-0.25 mm and 0.25 mm, and changes in their colloidal stability and PSDI properties due to cultivation were determined for both dry and wet sieved fractions. The result of the study showed that the aggregates in the cultivated soils were less stable than those in the fallow soils. The colloidal stability result show less structural stability of the aggregates by the high DR values obtained. The correlation result between silt+clay and dispersion ratio (DR) for all the land use types showed that the contribution of silt+clay to the stability of these soils was low. There was non-significant correlation between OC concentration and PSDI values and between silt + clay and total N, available P and OC in all the soils. The potential of the aggregates to disintegrate upon contact with water was more in the order of, Inceptisol at Eha-Amufu cultivated, IEh (C) 84.4% > Ultisol at Nsukka cultivated, UNsk (C) 74.4% > Ultisol at Nsukka fallow, UNsk (F) 72.0% > Inceptisol at Eha-Amufu fallow, IEh (F) 68.3% > Inceptisol at Ikem cultivated, Iik (C) 65.5% > Entisol at Nsukka fallow, ENSk (F) 65.2% > Inceptisol at Ikem fallow, Iik (F) 64.7% > Entisol at Nsukka cultivated ENSk (C) 61.8%. From the results of this study it is evident that the colloidal stability and PSDI of dry aggregates and wet-aggregates differ substantially even though their diameters are the same. Hence they will have differing impact on the soils.

Key word: Colloidal stability; PSDI; Silt; Clay, Fallow, Cultivated,

INTRODUCTION

Soil aggregation command a lot of complex processes in soil ecosystem of which many soil parameters and activities in soil such as organic matter content, composition of soil minerals, soil organisms etc play very

much important role in soil aggregation. According to Degens, (1997), and Beare *et al*; (1997), some of the most important processes in soil aggregation are flocculation of clay, cementation of dispersed clay or Fe and Al-oxy hydroxides, gluing effect of binding agents, such as polysaccharides' from plants, soil fauna and microbes, and cross-linking and enmeshment by fungal hyphae and plant roots.

Hence soil particles aggregate as a result of gluing by bacterially produced polysaccharides' and physical-chemical interactions between silicate clay surfaces and functional groups of partially decomposed organic matter (Aspiras *et al*; 1977, Aringhieri and Segui 1978, Tisdall and Oades, 1979, Sorensen, 1981,). The labile fractions of soil such as carbohydrates was also reported by Tisdall and Oades (1982) to be involved in the aggregation of soil particles due to their binding and cementing actions. The physical enmeshing of soil particles and micro aggregates by mycorrhiza fungal hyphae, fine roots and root derived particulate organic carbon can as well stabilize soil aggregates (Tisdall and Oades 1982; Golchin *et al* 1997). Golchin *et al*; (1997) emphasized that microbial decomposition of carbohydrates associated with aggregates and the resultant production of binding agents of mucilage and metabolites would promote the stability of soil aggregates.

The clay and silt fractions in the soil were found to be associated with organic matter concentrations in the different aggregate fractions (Christensen, 1986) and there were significant correlations between aggregate stability and clay and silt contents. At the microscopic level, clay and organic matter act as bridges between the sand and silt particles, producing micro aggregates of soil (McLaren and Cameron 1996). Iron and aluminum oxide were found to be well correlated with aggregate stability (Panayiotopoulos and Kostopoulou, 1989). Reid and Gross (1981) reported that roots and hyphae stabilize macro aggregates, defined as > 250mm diameter while the water-stability of micro-aggregates depend on the persistent organic binding agents which appear to be a characteristic of soil independent of management. This view was upheld by Mbagwu (1989a), when he argued that micro-aggregate < 0.20mm were held together by persistent binding agents like humus substances, and oxides of iron and aluminum. Macro aggregate however have their own life cycle repeating formation and break down depending on biological activity and fresh residue inputs (Denef *et al*;2001, Six *et al*; 2000 and Bidisha, 2010). Nevertheless, the effectiveness of various binding agents at different stages in the structural organization of aggregates forms the basis of a model which illustrates the architecture of an aggregate (Reid and Gross, 1989).

The organic binding agents have been classified into

- ❖ Transient, mainly polysaccharides
- ❖ Temporary, roots and fungal hyphae and
- ❖ Persistent, resistant aromatic components associated with polyvalent metal cat ions and strongly sobbed polymers (Tisdall and Oades, 1982).

The growth of soil fungi can produce sufficient mycelia to aggregate soil particles (Swaby, 1949; Bond and Harris, 1964), but this mechanical binding effect may be transient because the hyphae themselves are susceptible to microbial decomposition. More so, many soil microbes produce polysaccharides (Hepper, 1975) and it is generally agreed that such compounds are important in binding soil particles into larger units (Harris *et al*; 1966; Martin, 1971). This might be the reason why Skinner (1979) stressed that vigorous root and microbial activity in pasture soil may well explain the large proportion of highly stable aggregates usually found under grass. Earlier in their studies, Harris *et al*., (1966), maintained that the formation of soil aggregates that are stable to the action of water or mechanical forces depend on many factors, some of which are microbiological. Furthermore, the introduction of grasses in structural improvement and the increase is believed to be a result of physical enmeshment by roots and hyphae (Tisdall and Oades, 1979),

and cementing action of periodate-tetraborate and pyrophosphate sensitive materials (Raid and Gross, 1981, Anger and Mehuys 1989).

Many works by some researchers have also helped to x-ray and reassess the effect of organic matter and microbial processes in binding, cementing and aggregating of soil particles. For instance, it was discovered that organic materials act as transient, temporary or permanently binding agents at different levels of aggregation (Hades, 1987; Dexter, 1988; Perfect and Kay, 1991; Tisdall and Waters, 1991; Piccolo 1996). While the location of the organic matter within the aggregates influenced their accessibility for microbial decomposition (Elliot, 1986), the distribution of the microbial biomass within the aggregates varied with the intensity of land use (Gupta and Germida, 1988, Kandeler and Murer, 1993). Miller and Jastrow, (1990), and Monrozier *et al*; (1991) reported that the stability of the aggregates is influenced by the contents of the microbial biomass and root and fungi hyphae within the aggregates. Piccolo and Mbagwu (1990) found organic matter very effective in stabilizing silt and clay particles within dry micro aggregates with organic waste amended Italian soils. Though the stability of the silt-plus-clay particles correlated significantly with the humic acid content, there was no significant correlation with the total organic matter (Mbagwu and Piccolo, 1998).

Laboratory studies using synthetic oxide precipitates have shown that Al and Fe-oxy hydroxides promote aggregation (EL-Swaify and Emerson, 1975; Oades, 1984), though the effect of oxides on soil properties are variable with respect to the nature and amount of oxides and the environment of their pedogenic formation (Schnert-Mann and Taylor, 1989). Fe-oxy-hydroxide was found to be less effective compared to Al (OH)₃, polymers in maintaining the stability of soil aggregates (EL-Swaify and Emerson, 1975; Hsu, 1989), probably due to the fact that the Fe-oxy hydroxides make stronger tendency than Al (OH)₃ to crystallize, thus having lesser cementing effect except in soils frequently alternating between oxidized and reduced states (Frankel and Sheinberg, 1980). Poor aggregates, according to Hsu, (1989), do not necessarily indicate that they do not affect aggregation. Angers and Mehuys, (1989), and Haynes's *et al*; (1991), stressed that water extractable and acid-extractable carbohydrate carbon are more strongly correlated with the stability of macro-aggregates than total organic carbon or total extractable carbohydrate.

Nonetheless, the stability of soil aggregates can decline or decrease with the reduction in carbohydrate levels following microbial decomposition. The production of mucilage and metabolite glues will equally reduce (Golchin *et al*, 1997). Soil structural stability and soil organic carbon content usually decrease and changes the distribution of soil aggregates with cultivation (Six *et al*; 2000, Eynard *et al*; 2004). While Young and young (2001) reported that soil structural stability will decrease under most forms of cultivation. Soil aggregate formation can as well be inhibited by tillage, traffic with agricultural machines and rainfall that accelerate aggregate breakdown (Bronick and Lal 2005) they also found temporal and spatial heterogeneity and complexity in soil continuum that make the soil difficult directly to measure the aggregation and carbon storage in soil. Most of the works cited above on the stability of soil aggregates on different soil types and management were mainly in the temperate region where the soils are very different from those in the tropics. The result got on the tropical soils revealed the need for more work and studies to be done on the tropical soils. Hence the essence of this research, which was undertaken to study the colloidal stability and potential structural deformation index of four Nigerian soils.

Materials and Methods

Field methods

Soil samples from the 0-25cm depth were collected from cultivated and adjacent fallow lands in four different locations in Nsukka area of south eastern Nigeria. Care was taken to minimize disturbance during sampling and transportation. The area has a rainforest savannah type of vegetation with a mean annual temperature of 24^oc. The area lies within latitude 06^o 61¹N and longitude 07^o 25¹E of Nigeria. The soils sampled for the study are classified according to soil taxonomy as an Ultisol, belonging to the sub-group, Typic Kandistult (Nkpologu series), Entisol belonging to Lithic Ustorthent (Uvuru series), while the other two soils belong to Vertic Inceptisol (SSS, 1992). These soils have been under cultivation for over and about 8years while fallow soils varied from 3 to 4years old. The cultural practices adopted in these soils were slashing and burning and tillage is merely by the use of Hoe. Fertilizer application is basal and it is mainly NPK.

Table 1 Location, Classification and Land use type

Location	Classification	Treatment symbol	Land use type
Nsukka Hill Site	Lithic Ustorthent	ENsk (F)	Fallow
	(Uvuru series)	ENsk (C)	Cultivated
Nsukka Poultry Site	Typic Kandistult	UNsk (F)	Fallow
	(Nkpologu series)	UNsk (C)	Cultivated
Eha-Amufu Site	Inceptisol	IEh (F)	Fallow
	(With vertic properties)	IEh (C)	Cultivated
Ikem site	Inceptisol	Iik (F)	Fallow
	(With vertic properties)	Iik (C)	Cultivated

Laboratory method

The soil samples were air-dried at room temperature and then sieved through a 5.00mm sieve. Clods were carefully crushed by hand along lines of natural cleavages to pass the sieve. Two hundred and fifty grams (250g) of the sieved sample, at a time, were transferred to the upper most of a nest sieve of sizes 2, 1, 0.5 and 0.25mm. They were shaken mechanically for 10 minutes. Further sieving by hand was done where necessary. This procedure, similar to that described by Kemper and Chepil (1965), resulted in the separation of the following aggregate fractions 5-2, 2-1, 1-0.5, 0.5-0.25 and < 0.25mm. The separation continued until enough quantity of each fraction was collected for further analysis.

The mean weight diameters of dry aggregates (MWDD) was determined by the method of Kemper and Chepil (1965), while the method of Kemper (1965), was used to determine the mean-weight diameter of wet aggregates (MWDW). To assess the relative susceptibility of the soils to disintegrate upon wet-sieving, an index, the potential structural deformation index, PSDI (Mbagwu and Bazzoffi, 1989) was computed thus:

$$\varepsilon = \sum_{i=1}^n (\theta_i w_i)$$

$$\text{PSDI} = \left\{ 1 - \frac{\sum_{i=1}^n \theta_i d_i}{\sum_{i=1}^n w_i} \right\} \times 100 \quad (1)$$

Where:

θ_i = Mean diameter of the i th size Fraction (mm).

w_i = Proportion of the total mass retained in the i th size fraction (from wet-sieving) gg^{-1} .

d_i = Proportion of the total mass retained in the i th size fraction (from dry sieving) gg^{-1} .

n = Number of sieve (size) fraction used.

The PSDI was used to normalize the changes in distribution of dry and wet aggregates to enable a realistic comparison of the relative effects of land use across the different soils to be made.

Colloidal stability:

The procedure of Middleton (1930) was used. The procedure involved determining the percent silt + clay in water-dispersed samples (A) and that in sodium resin-dispersed samples (B). The index $(A/B \times 100)$ known as the dispersion ratio DR, is a measure of the potential of the individual aggregates to resist break down upon contact with water molecules. In this study, 20g sample from each aggregate fraction was soaked in either distilled water or sodium hexametaphosphate solution for 24 hours, followed by end-over-end shaking on a mechanical shaker for 30 minutes at 10 revolutions per minute. The dispersed silt + clay were determined by the hydrometer method.

Data Analysis:

Little and Hills (1972) simple linear regression and correlation analysis was used to analyze the data obtained from this work in order to identify which intrinsic and dynamic properties most influenced the stability of the aggregates most. While least significant difference (LSD) at 5% was used to compare the treatment means obtained in PSDI and DR. The cumulative aggregate distribution from dry and wet sieving procedures of each aggregate size was later cartographed to show a vivid picture of PSDI distribution of each soil type.

Result and Discussion

Colloidal Stability

The values of dispersion ratio (DR) given in Table 2, show significant ($p=0.05$) differences among the soil types and land use systems. Higher dispersion ratio was observed in lik (F), while the least among the soil types and land use system was obtained from UNsk (F). The DR result also show lower values in cultivated soils compared to the fallow soils with the exception of UNsk where the cultivated value was higher than that of the fallow soil. The value obtained from IEh however, indicated that cultivation had no effect on the dispersion ratio. The nature of the result so obtained from these soil types and land use systems probably suggests that there was no decrease in the aggregate stability of these soils due to cultivation. The higher DR values show less structural stability of the aggregates.

198

199 Table 2: Aggregate stability of the fallow and cultivated soils as evaluated by two different indices

200	SOILS	PSDI	DR
201	ENsk (F)	65.16	72.73
202	ENsk (C)	61.84	66.80
203	UNsk (F)	71.98	55.55
204	UNsk (C)	74.41	62.50
205	IEh (F)	68.33	76.47
206	IEh (C)	84.41	76.47
207	lik (F)	64.68	87.50
208	lik (C)	65.51	73.33
209	Mean	69.54	71.40
210	LSD (0.05)	3.64	4.57

211 PSDI = Potential structure deformation index, DR= Dispersion ratio, LSD = Least significant difference, F = Fallow, C =
 212 Cultivated, ENSK = Entisol at Nsukka, UNSK = Ultisol at Nsukka, IEh = Inceptisol at Eha-Amufu, lik = Inceptisol at Ikem.

213 The non-significant correlation between silt + clay and DR for all the land use types (Table 3) showed that
 214 the contribution of silt + clay to the stability of these soils was low. Allison, (1973) observed that clay
 215 stabilizes soil aggregates. Thus the quantity of clay present may be an important factor in stabilizing soil
 216 aggregates. Also Heinonen (1955) suggested that a positive and significant correlation between aggregate
 217 stability and clay content should be expected in soils containing more than 30% (W/W) clay. The result
 218 could also be attributed to the nature of microbes, organic matter, or clay, Fe and Al oxide contents of these
 219 soils. Hepper (1975) observed that many soil microbes produce polysaccharides and it is generally agreed
 220 that such compounds are important in binding soil particles into larger units (Martin, 1971). Miquel and
 221 Norton (1994) reported that aggregate stability is largely dependent upon OM, clay, Fe and Al oxide
 222 contents. Also Panayiotopoulos and Kostopoulou (1989) found Fe and Al oxide to be well correlated with
 223 aggregate stability.

224

225

226

227

228

229

230

231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260

Table 3 Correlation of silt + clay (%) and OC % with other Soil Properties (N=8)

Independent Variable	Dependent Variable	Correlation Coefficient (r)
Silt + clay %	DR %	- 0.824 ^{ns}
	Total N %	- 0.146 ^{ns}
	Avail. P (Mgkg ⁻¹)	- 0.644 ^{ns}
	OC %	- 0.213 ^{ns}
OC %	PSDI %	- 0.364 ^{ns}
	DR %	- 0.438 ^{ns}
	Total N %	- 0.949 *
	Avail. P (Mgkg ⁻¹)	- 0.364 ^{ns}

* Significant at P = 0.05 ns = not significant.

Potential structure deformation index:

The PSDI values result given in Table 2 show higher values among the soil types and land use system, highest PSDI value among the soils was obtained from IEh (C) and this was significantly (p=0.05) different among the soil types and land use systems. The potential structural deformation index (PSDI) is used to normalize aggregate distribution between the dry and wet sieving procedures, as to obtain valid comparisons between land use types. The lower the percentage, the more resistant is the soil to disruptive forces. The cumulative aggregate distributions from dry-and wet sieving procedures for the ENsk are given in Fig 1, UNsk in Fig 2, and IEh in Fig 3 and Iik in Fig 4. (See Table 1 for the explanation of the symbols).




FIG 1: Distribution of aggregate obtained from dry and wet sieving procedures in an Entisol (Nsukka Hill Site) under different lands uses. (PSDI=Potential structure deformation index)




FIG 2: Distribution of aggregate obtained from dry and wet sieving procedures in an Ultisol (Nsukka Poultry site) under different land uses. (PSDI=Potential structural deformation index)




FIG 3: Distribution of aggregate obtained from dry and wet sieving procedures in an Inceptisol (Eha-Amufu site) under different land uses. (PSDI=Potential structural deformation index)




FIG 4: Distribution of aggregates obtained from dry and wet sieving procedures in an inceptisol (Ikem site) under different uses. (P SDI= Potential structural deformation index)

The area under the two curves indicates the relative resistance of the dry aggregates to disruption by water. The smaller the area the more resistant is the soil to the disruptive forces. This showed that the aggregates of the fallow soils were more resistant to disruptive forces compared to those of the cultivated soils. This statement is only true for the PSDI of IEh given in Fig 3.

The correlation coefficient between OC concentration and PSDI values of these soils were small and non-significant (Table 3). This indicates that the contributions of organic matter to the stability of the aggregates of these soils were small. Mbagwu and Bazzoffi (1989) made similar observation on Italian soils. Generally, the soils are not resistant to deformation because the PSDI values are large.

Nonetheless, the PSDI showed an order of vulnerability to disintegration upon contact with water molecules of IEh (C) 84.4% > UNsk (C) 74.4% > UNsk (F) 72.0% > IEh (F) 68.3% > Iik (C) 65.5% > ENsk (F) 65.2% > Iik (F) 64.7% > ENsk (C) 61.8%. In all the soils a significant correlation between OC and total N was established (Table 3). This indicates that the source of total N in the soils is from OM. The non-significant correlation between silt + clay and total N, available P and OC (Table 3) in all the soils, signifies that the sources of these nutrients are not from silt + clay contents in the soils, while the aggregates evaluated by dispersion ratio showed that the amount of soil to be lost from stimulated rainfall will be high in wet-sieved samples compared to the dry-sieved ones.

Conclusion

From the study, it was found out that cultivated soils in some of the soils studied did not have less stable aggregates nor disintegrated faster than those of the fallow soils when they came in contact with water. The PSDI values were found to be large, indicating that the soils are not resistant to deformation. There has to be a land use strategy to maintain stable aggregation but at the same time allow an ordered agricultural production of crops.

References

- Allison, F. E. (1973). Soil organic matter and its role in crop production Elsevier Scientific publishing company, Amsterdam, 637pp
- Angers, D. A, and Mehuys, G. R. (1989). Effects of cropping in Agriculture, Perth, Western Australian, Paper No SEAG 981022
- Aringhieri, R and Sequi, P. (1978). The arrangement of soil organic matter in a soil crumb 145-150pp In: W.W. Emerson et al (Eds) Modification of soil structure. John Wiley and Sons Inc. New York.
- Aspiras, R. B, Allen, O. N., Chester, G. and Harris, R. F. (1977). Chemical and physical stability of microbial stabilized aggregates. Soil Sci. Am. Proc. 35. 283-286
- Beare, M. H., Hu, S., Coleman. D.C., Hendrix, P.F. (1997) Influence of mycelia fungi on soil aggregate and organic matter storage in conventional and no-tillage soil Appl. Soil Ecol, 5,211-219
- Bidisha M., Joera, R., Yakov, K. (2010). Effect of aggregation process on distribution of aggregates size fraction and organic C content of a long-term fertilized soil European journal of soil biology (in press). 125,156, 200
- Bond, R. D. and Harris, T. R. (1964). The influence of micro flora on the physical properties of soils 1 Effects associated with filamentous algae and fungi J. Soil Res. 2: 111-122.
- Bronick, C. J. and Lal, R. (2005). Soil structure and management a review Geoderma 124:3-22.
- Christensen, B. T. (1986). Straw in corporation and soil organic matter in macro-aggregates and particle sizes separates. J. Soil Sci. 31:123-135.

- 415 Deneff, K. J., Six, J., Paustain, K., Merckx, R. (2001) importance of macro-aggregates dynamics in
416 controlling soil carbon stabilization in short-term effects of physical disturbance inducing by
417 dry-wet cycle. Soil biology and biochemistry 33:2145-2153.
- 418 Dexter, A. R. (1988). Advances in characterization of soil structure Soil Tillage Res. 11:199-238.
- 419 Elliot, E. T. (1986). Aggregate structure and cultivated soils. Soil Sci. Soc. Am. J. 50:627-633.
- 420 El-Swaify, S. A. and Emerson, W. W. (1975). Changes in the physical properties of soil clays due to
421 precipitated aluminum and iron hydroxides.1 Swelling and aggregate stability after drying Soil
422 Sci. Soc. Am. J. 99:1056-1063
- 423 Eynard, A., Schumacher, T. E., Lindstrom, M. J., Malo, D. D. and Kahl, R. A. (2004). Wet ability of soil
424 aggregates from cultivated and uncultivated Ustolls and Usterns. Aust. J. soil Res. 42:163-170.
- 425 Frankel, H and Sheinberg, I. (1980). The effect of hydroxyl-Al and hydroxyl-Fe polymers on
426 montmorillonite particle size Soil Sci. Soc. Am. J. 44: 626-629.
- 427 Golchin, A. P. and Dengiz, O. (1997). Effect of modifying land cover and long-term agricultural practices
428 on the soil characteristics in native forest land J. Environ. Biol. 29:667-682.
- 429 Gupta, V.V S. R. and Germida, J. J. (1988). Distribution of microbial biomass and its activity in different
430 soil aggregate size classes as affected by cultivation, Soil Biol. Biochem. 20:777-786.
- 431 Hadas, A. (1987). Long-term tillage practice effects on soil aggregation modes and strength. Soil Sci: Soc.
432 Am. J. 51:191-197.
- 433 Harris, R. F., Chester, G. and Allen, O. N. (1966). Dynamics of soil aggregation, Adv. Agron. 18:107-169.
- 434 Heinonen, R. (1955). Soil aggregation in relation to texture and organic matter, Agrogeol. Julk. 64:17
435 Abstracts Soils and Fertilizers 19, 43, 1956.
- 436 Hepper, C.M. (1975). Extracellular polysaccharides of soil bacteria In: Soil microbiology, Walker. (Ed)
437 London, Butterworth's.
- 438 Hsu, P. H. (1989). Aluminum hydroxides and oxy-hydroxides In: minerals in soil environments 2nd edn J. B.
439 Dixon and S.B. Weed (eds), SSSA Book Series 1, Madison, WI. 331-378
- 440 Kandeler, E. and Murer, E. (1993). Aggregate stability and soil microbial processes in a soil with different
441 cultivation Geoderma 56:503-513.
- 442 Kemper, W.D. (1965). Aggregate stability In: C. A. Black *et al* (Eds) Methods of soil Analysis part 1 No. 9,
443 511-519. Am. Soc. Agron. Madison, Wisconsin.
- 444 Kemper, W.D. and Chepil, W. S. (1965). Size distribution of aggregates In: C. A. Black *et al* (Eds) part 1.
445 Methods of soil analysis No 9: 499-510. Am. Soc. Agron. Madison, Wisconsin.
- 446 Little, T. M. and Hills, F. J. (1972). Statistical methods in Agricultural Research Univ. of California, Davies,
447 242pp
- 448 Martin, J. P. (1971). Decomposition and binding action of polysaccharides in soil Soil Biol. Biochem. 3:33-
449 41.

- 450 Mbagwu, J. S.C (1989a). Effects of organic amendment on some physical properties of a tropical Ultisol.
451 Boil. Wastes 28:1-23.
- 452 Mbagwu J. S. C. and Bazzoffi, P. (1989). Properties of soil aggregate as influenced by tillage practices. Soil
453 use and Management. 5:180-188.
- 454 Mbagwu, J. S.C. and Piccolo, A (1998). Water dispensable clay in aggregate of forest and cultivated soils in
455 Southern Nigeria in relation to organic matter constituents. In: Carbon and Nutrient Dynamics
456 in Natural and Agricultural Tropical Ecosystems (Eds) L. Bergstrom. And H. Kirchmen. 71-
457 83pp. CAB International
- 458 McLaren, R. G and Cameron, K.C. (1996). Soil Science 2nd edition Oxford University press. London.
- 459 Middleton, H. E (1930). Properties of soil which influence erosion, U.S. Dept. Agric. Tech. Bull. 178pp.
- 460 Miller, R. M. and Jastrow, J. D. (1990). Hierarchy of root and mycorrhizal fungi interaction with soil
461 aggregation. Soil Biol. Biochem. 22:579-581.
- 462 Miquel, R. J. and Norton, D. L. (1994). Aggregate stability and rain impacted sheet erosion of air in dried
463 and pre-wetted clayey surface soils under intense rain. Soil Sci.158:321-330.
- 464 Monrozier, L. T., Ladd, J. N., Fitzpatrick, R. W., Foster, R. C. and Raupach, M. (1991). Components and
465 microbial biomass contents of size fraction in soils of contrasting aggregation. Geoderma
466 49:37-62.
- 467 Oades, J. M. (1984). Soil organic matter and structural stability: mechanisms and implications for
468 management. Plant and Soil 76:319-337.
- 469 Panayiotopoulos, K. P. and Kostopoulou, T. (1989). Aggregate stability dependence on size, cultivation and
470 various soil constituents in red Mediterranean soils (Alfisols) Soil Tech. 2:59-69.
- 471 Perfect, E. and Kay, B. D. (1991). Fractal theory applied to soil aggregation. Soil Sci: Soc. Am. J. 55:152-
472 158.
- 473 Piccolo, A. (1996). Humus and soil conservation In: Piccolo A. (ed.), Organic Matter in Terrestrial
474 Ecosystem. Elsevier Science Publishers, Amsterdam 83-98.
- 475 Piccolo, A. and Mbagwu J. S. C. (1990). Effect of different organic wastes amendments on soil micro-
476 aggregates stability and molecular sizes of humic substances Plant and soil 123:27-37
- 477 Reid, J. B. and Gross, M. J. (1981). Effect of living roots of different plant species on the aggregate stability
478 of two arable soils J. Soil Sci. 32:521-541.
- 479 Schnert-Mann, U. and Taylor, R. M. (1989). Iron oxides. In: Minerals in Soil Environments 2nd ed. J. B.
480 Dixon and S.B. Weed (Eds).SSSA Book Series 1, Madison, WI 379-438pp.
- 481 Six, J., Elliot, E. T., Paustian, K. and Comb rink, C. (2000). Soil structure and organic matter 1 Distribution
482 of aggregate size classes and aggregate associated carbon Soil Sci. Soc. Am. J. 64:681-687.
- 483 Skinner, F. A. (1979). Effect of incubation on soil aggregate stability J. soil Sci. 30:473-481.
- 484 Soil Survey Staff (1992). Soil Taxonomy: A basic system of soil classification for making and interpreting
485 soil surveys. USDA-SCs Agric. Hand book 436. U.S. Govt. Printing Office, Washington D.C

- 486 Sorensen, L. H. (1981). Carbon-nitrogen relationships during the humification of cellulose in soils
487 containing different amounts of clay Soil Biol. Biochem. 13:313-321.
- 488 Swaby, R. T. (1949). The relationship between micro-organism and soil aggregation, J. Gen. Microbial.
489 3:236-254.
- 490 Tisdall, J. m. and Oades, J. M. (1979). Stabilization of soil aggregates by the root system of ryegrass. Aust.
491 J. soil Res. 17:429-441.
- 492 Tisdall, J. M. and Oades, J. M. (1982). Organic matter and water stable aggregates in soils. J. Soil Sci.
493 33:141-163.
- 494 Tisdall, J. M. and Waters, A. G. (1991). Aggregate hierarchy in soils Aust. J. Soil Res. 29:815-828.
- 495 Young, A. and Young, R. (2001). Soils in the Australian landscape Oxford University Press, Melbourne.
- 496
- 497
- 498
- 499
- 500