

Original Research Article

Evaluation of Sustainable Production Practices for Asian Vegetables (Luffa and Bitter Gourd) and their Mineral Nutrient Analysis in a Piedmont Soil of North Carolina

ABSTRACT

Aims: To reduce the inorganic fertilizers used to grow Asian vegetables (Luffa and Bitter gourds) by incorporating cover crops without compromising yield in piedmont soils of North Carolina

Study design: A split-split plot design was used in this study with two main plot treatments and four rates of fertilizer as subplot treatments replicated four times.

Place and Duration of Study: The study was conducted at the North Carolina A&T State University Farm, Greensboro, Guilford County, NC. from Fall, 2007 to Summer, 2008.

Methodology: Luffa (*Luffa acutangula* 'Rama') and bitter gourd (*Momordica charantia* 'Comet') were grown in a Mecklenburg Sandy Loam (fine, mixed, thermic ultic Hapludalfs) soil. Two cover crop treatments (Cover crop and no cover crop) and four fertilizer treatments (T1: 0-0-0, T2: 56-28-112, T3: 84-56-168, and T4: 168-112-224 N-P-K kg/ha) were studied to determine the yield of Asian vegetables in the Piedmont of North Carolina. The vegetables were hand harvested at weekly intervals for 9 weeks.

Results: Both luffa and bitter gourd yields increased with increase in rate of fertilizer. Cover crop treatment produced higher yields of luffa and bitter gourd compared to no cover crop treatment for all four fertilizer treatments. Statistical analysis has shown no significance difference in yield in treatment T1 – T3 but the yields were significantly higher ($p=0.05$) in T4 treatment for both luffa and bitter gourd. Cover crop residue incorporated in to the soil decomposed and released nitrogen which was utilized by the vegetable crops and the effect was evident in increased yields in cover crop treatment plots.

Conclusion: Asian vegetables (luffa and bitter gourd) were grown in piedmont soils of North Carolina and the study showed that T4 treatment (168-112-224 N-P-K kg/ha) resulted in higher yields than all other treatments. This study provides evidence that Asian vegetables – such as luffa and bitter gourd – can be grown successfully in the piedmont region of North Carolina.

Keywords: Asian vegetables, luffa, bitter gourd, cover crop, sustainable, fertilizer, Piedmont soil, North Carolina

1. INTRODUCTION

Luffa/Ridge gourd/Sponge gourd/Chinese okra (*Luffa acutangula* (L.) ROXB.) and bitter gourd/bitter melon (*Momordica charantia* L.) are two very popular asian vegetables consumed by people from Asian and other cultures. Luffa is consumed at young stage as a vegetable. Mature luffa fruit does not have the same palatability as the immature fruits. Bitter

gourd is also harvested and consumed at young stage as mature fruits are more bitter in taste with seeds becoming hard to chew. Both luffa and bitter gourd have alternate uses than their consumption as vegetables.

Luffa consumed as a vegetable can supply some antioxidant constituents to the human body [1]. Mature luffa is very fibrous and used for sponge production [2]. Luffa sponge has fibrous vascular system with continuous network system with a pore size range of 10 – 20 µm. Sponge from luffa has been used as supporting material for the cell culture of human hepatocyte cells [3]. Luffa sponge has also been used as absorbent of industrial dyes, such as Malachite Green (MG), which when released into aquatic systems are detrimental to aquatic life [4]. Phosphate, sulfate and nitrate anions were successfully exchanged by converting luffa sponge into an anion exchanger by etherification using 2-chloroethyldiethylamine hydrochloride (CEDA-HCl) followed by crosslinking using epichlorohydrin [5]. Luffa fiber both in its natural form and modified with NaOH were efficient and effective in absorbing lead (Pb^{2+}) from contaminated waters in batch experiments [6]. Sliced luffa sponge loaded with both flocculating and non-flocculating cells are used for continuous ethanol production in column-type bioreactors [7]. L-Lactic acid, an important chemical product used in food, medical and polymer industries has been produced from cassava starch using circulating loop bioreactors with cells immobilized in luffa sponge [8]. Clavulanic acid is a powerful antibiotic and can be produced by immobilizing *Streptomyces clavuligerus* on luffa sponge [9]. Luffa seeds are high in protein (39%) and fat (44%) content and oleic and linoleic fatty acid content is 68 % in the kernel oil [10].

Extracts from both luffa and bitter melon are found to have Free radical scavenging activity (FRSA) [11]. Bitter gourd has medicinal value such as its hypoglycemic nature [12, 13, 14]. Juice extracted out of the bitter gourd fruit and young shoot and leaves has a number of medicinal properties, such as blood cleansing, immune boost, etc. Bitter gourd consumption in the diet can help decrease blood glucose levels as well as some antioxidant properties [15, 16]. Chemical compounds in bitter gourd have the ability to reduce the amount of glucose that is released into the blood by inhibiting the enzymes that breakdown disaccharides (e.g. Sucrose) into monosaccharides (e.g. glucose and fructose) [17, 18]. Bitter gourd can regulate the transport channels for glucose transfer thereby reducing glucose transport into the blood [19]. This regulatory action is very significant in treating both Type I and Type II diabetic patients and helps to prevent high blood sugar levels after a meal. In an experiment on mice with Type I diabetes, bitter gourd proved an effective treatment by increasing secretion of insulin from pancreas [20, 21]. HIV inhibitor protein MRK 29 extracted from Thai bitter gourd fruit and seeds has shown to reduce the expression of viral core protein (p24) in HIV infected cells by 82% [22].

These Asian vegetables are popular and multipurpose and hence, successful and sustainable growing practices must be developed. The city of Greensboro NC and surrounding areas have large Asian communities and hence a good market for Asian vegetables exists. Growing these vegetables on small farms can help the small farms generate income where the cash crop tobacco had been grown previously. Keeping the needs of the small farms, existing market demand and sustainable growing practices in mind, this study was conducted at North Carolina A&T State University Agricultural Farm to both evaluate and demonstrate the cultivation practices of these vegetables. The goal of this study was to reduce the use of inorganic fertilizers by incorporating cover crops to grow Asian vegetables without compromising yield.

2. MATERIAL AND METHODS

2.1 Experimental location

The experiment was conducted at the North Carolina A&T State University Research Farm in Guilford County, North Carolina. The landscape at the experimental location was representative of many regions in the Piedmont and southeastern United States. The soil was a Mecklenburg Sandy Loam (fine, mixed, thermic ultic Hapludalfs). The plots were set up as 4 replications under cover crop (CC) and no-cover crop (NC) systems. Four different fertilizer treatments (T1: 0-0-0, T2: 56-28-112, T3: 84-56-168, and T4: 168-112-224 N-P-K kg/ha). A split-split plot design was used. Crimson clover (*Trifolium incarnatum*) and cereal rye (*Secale cereale*) were planted in the plots (cover crop system) in Fall, 07 and were incorporated into the soil in Spring, 08. In each treatment 50% of the fertilizer was applied and incorporated in to the soil with a disc plough and raised beds were prepared. The remaining 50 % fertilizer was supplied by fertigation. Raised beds 2' (W) and 30' (L) covered with black plastic and lined with drip tape for irrigation and fertigation were prepared prior to seed planting. Seeds were planted in the beds in the Spring 2008. Harvesting of the vegetables was done between Aug 8, 2008 and Oct, 9, 2008 for a total of 9 times at weekly intervals when the vegetables were at marketable stage (54 days after planting).

Cover crop treatments plots were studied to determine the rate of decomposition and release of plant nutrients in to the soil. The cover crop mixture (Crimson clover + cereal rye) was collected before incorporating it in to the soil. A 40g (9.2 g crimson clover + 30.8 g cereal rye) sample of the over crop mixture of was placed in a fiber glass mesh bag. Five bags per each fertilizer treatment plot within the cover crop treatment were placed at 3" depth and 24" apart. A total of 80 mesh bags (5 bags per treatment X 4 fertilizer treatments X 4 replications) were placed in the cover crop treatment plots. The mesh bags were collected at 3 week intervals during the luffa and bitter gourd growth period. Each time the mesh bags were collected, they were washed carefully to get rid of the soil without losing any cover crop residues. Washed cover crop residues were dried at 65°C for 48 hrs and their dry weights were recorded. Dried cover crop residues were then ground using a Wiley Laboratory Mill Model 4 (Thomas Scientific, Philadelphia) in to 1mm size particles. Samples from the sieved plant material was analyzed for total C and N by dry combustion method.

2.2. Mineral nutrient analysis

Luffa and bitter gourd tissues were analyzed to determine the nutrient content by Dry ashing procedure [23] given in SSSA book series:3. A 0.5 g of dried 20-mesh plant tissue was taken in a 15ml porcelain crucible and heated at 500°C for 8 hrs in a muffle furnace. The crucibles were taken outside of the furnace and cooled down to room temperature. The ash was dissolved by adding 10 ml of dilute acid (300 ml of HCl and 100 ml of HNO₃ in 1 L of pure water) and allowing the mixture to settle down at the bottom of the crucible. The supernatant was filtered and diluted to 50 ml for elemental assay.

2.3. Experimental design and statistical analysis

A split-split-plot design was used in this study with two cover crop treatments as main plots and four fertilizer treatments replicated four times (32 experimental units in total). Experimental units consisted of 3m x 10m subplots. Data for the experiments were subjected to analysis of variance using Statistical Analysis System, Version 9.2 [24].

3. RESULTS AND DISCUSSION

Luffa and bitter gourd were harvested 9 times between August and October, 2008. Total yield obtained with each fertilizer as well as cover crop treatments are presented in Table 1. Both luffa and bitter gourd produced consistently high yields with increase in fertilizer, T4 was the highest yield treatment in both crops for cover as well as no cover crop treatments.

3.1 Luffa yield

Luffa yield increased consistently with increase in fertilizer treatment in both cover and no cover crop treatments. Yields were consistently higher in cover crop plots compared to the no cover crop plots, except for T3 (Table 1). Duncan grouping of the yield data has shown that significantly higher yield ($P = 0.05$) was observed in luffa and bitter gourd from the plots applied with 168 kg N/ha compared to the other treatment conditions.

Table 1. Yield data for luffa and bitter gourd at different fertilizer treatments under cover and no-cover crop systems

N-P-K Rate (kg/ha)	Luffa (t/ha)		Bitter Gourd (t/ha)	
	No Cover crop (NC)	Cover crop (CC)	No Cover crop (NC)	Cover crop (CC)
0-0-0	25.9 ^C	27.0 ^C	5.5 ^B	8.5 ^B
56-28-112	40.3 ^B	44.9 ^B	8.1 ^B	11.7 ^B
84-56-168	56.8 ^{BA}	41.2 ^{BA}	10.9 ^{BA}	12.5 ^{BA}
168-112-224	57.3 ^A	61.1 ^A	15.9 ^A	16.6 ^A

NC – No cover crop; CC – Cover crop

Treatment means with same letter are not significantly different

Statistical analysis has shown that luffa yield was significantly affected by fertilizer treatments (Table 2) but not by cover crop treatment.

Table 2. SAS analysis of variance results showing the significance of treatments on luffa and bitter gourd yield

Luffa (t/ha)					
Factor	DF	Sum of Squares	Mean Square	F	P
Fertilizer Rate (FR)	3	4542.46	1514.15	9.16	0.04
Cover Crop (CC)	1	19.84	19.84	0.12	0.73
(FR) x (CC)	3	541.9	180.63	1.09	0.4
CV=29.02					
Bittergourd (t/ha)					
Factor	DF	Sum of Squares	Mean Square	F	P
Fertilizer Rate (FR)	3	363.55	121.18	4.73	0.03
Cover Crop (CC)	1	39.82	39.82	1.55	0.24
(FR) x (CC)	3	10.15	3.38	0.13	0.93

CV=45.03

159
160
161

3.2 Bitter gourd yield

162 Bitter gourd yield was higher with T4 treatment in both cover crop treatments (Table 1).
163 Similar to luffa, bitter gourd also showed a significant effect on yield with fertilizer treatments
164 but not with cover crop treatments (Table 2).


165
166

3.3 Cover crop vs no cover crop treatment in luffa and bitter gourd

167 Luffa and bitter gourd produced higher yields with the inclusion of cover crop residues.
168 Among the cover and no cover crop treatments, yield in cover crop incorporated plots was
169 higher for both the vegetables. The increase in yield can be attributed to the decomposition
170 and release of extra nitrogen and other nutrients that are utilized by plants. The
171 decomposition and release of nitrogen is supported by the data collected from another study
172 conducted in the same plots [25], according to the study 13.33 – 18.32g/kg of nitrogen was
173 released from the cover crop residue that was buried in the soil for decomposition. Bitter
174 gourd yield was comparatively higher in the cover crop plots than in the no cover crop plots
175 for all N-fertilizer treatments. The yield data of luffa and bitter gourd in cover crop treatment
176 plots indicates that the nitrogen released from the cover crop residue is utilized by the
177 vegetable crops and thereby increased the yield over the no cover crop treatment yields.

178 Temperature (Fig 1) during the luffa and bitter gourd growing period was on the
179 decline after the month of august when the first harvest occurred. Since Luffa and bitter
180 gourd are Asian vegetable, they tend to perform better and produce higher yields at warmer
181 temperatures. Cover crop was incorporated for the first time in these plots during the
182 growing period and hence the release of nitrogen from the crop residues was in low
183 quantities.

184



185

Fig 1. Average monthly temperatures in Guilford county area of North Carolina during the study conducted on luffa and bitter gourd.

Rainfall (Fig 2) was not uniform during the growing period. August and September months have received rainfall close to 6 and 5 inches respectively and October month has received only an inch of rainfall which might have affected the yield of luffa and bitter gourd. These two factors might have contributed to the slower decomposition of the cover crop residues incorporated in to the soil and there by releasing less nitrogen. Cover crop residue incorporation to improve crop yield is a long term process because of the slow decomposition rate of the plant residues and release of nutrients.




Fig 2. Monthly average precipitation during the year 2008 in Guilford county area of North Carolina

3.4 Mineral nutrient analysis

Luffa and bitter gourd tissue analysis (Table 3) for nutrient elemental concentrations has shown that both the vegetables have good amounts of all the elements that were analyzed.

Table 3. Mean concentration of mineral nutrient content of luffa and bitter gourd

Element	Luffa (mg/g)	Bitter Gourd (mg/g)
Ca	2.27 A	1.64 B
Cu	0.001B	0.01A
Fe	0.02A	0.02A

K	12.8 B	20.48 A
Mg	2.24B	2.57A
Mn	0.01B	0.02A
Na	1.66 A	1.60A
P	4.14A	4.30 A
S	0.94A	0.67B
Zn	0.03A	0.02B

Means values for each nutrient with the same letter are not significantly different

Bitter gourd has significantly high concentrations of potassium (K) which when consumed as vegetable can help control high blood pressure. In addition to potassium bitter gourd is also significantly high in copper, magnesium and manganese (Table 3). Luffa has significantly high concentration of calcium, sulfur and zinc. Nutritionally, both vegetables are good source of mineral nutrients. Bitter gourd has phytochemicals that control blood glucose level (glycemic index) and antioxidant components [15,16].

4. CONCLUSION

Results indicate that both luffa and bitter gourd produced higher yields in the cover crop treatment plots than in the no-cover crop plots. Significant yield differences ($P=0.05$) were observed in the luffa vegetable yield from the plots applied with treatment T4:168-112-224 N-P-K kg/ha than from the other treatment conditions for both the vegetables. Although yields were not significantly different between the cover crop and no-cover crop treatments, we may recommend growing luffa and bitter gourd under cover crop treatment and T4 fertilizer application. The consistently higher yields observed in case of both luffa and bitter gourd suggest that cover crop residue incorporation in to the soil will relatively increase the yields, even in short term. Cover crop residue inclusion in to the soil helps in building the organic matter of the soil and in general improve soil health for greater productivity in the long run. Small farms will be encouraged to incorporate cover crops in to their crop rotations in order to obtain higher yields as well as improve the soil conditions. Bitter gourd is a good source of potassium to include in the diet of patients with high blood pressure and to control blood glucose level.

238 **COMPETING INTERESTS**

239

240 No competing interests exist.

241

242

243

244 **CONSENT (WHERE EVER APPLICABLE)**

245

246 **N/A**

247

248

249 **ETHICAL APPROVAL (WHERE EVER APPLICABLE)**

250

251 **N/A**

252

253 **REFERENCES**

254

255

256 1. Du Q, Xu Y, Li L, Zhao Y, Jerz G, Winterhalter P. Antioxidant Constituents in the
257 Fruits of *Luffa cylindrica* (L.) Roem. J. Agric. Food Chem. 2006;54:4186-4190.

258 2. Davis J. DeCourley CD. Luffa sponge gourds: a potential crop for small farms. In:
259 Janick J, Simon JE, editors, New crops. Wiley, New York; 1993.

260 3. Chen JP, Yu SC, Hsu BR, Fu SH, Liu HS. Loofa sponge as a scaffold for the culture
261 of human hepatocyte cell line. Biotechnol. Prog. 2003;19:522-527.

262 4. Altınışık A, Gür E, Seki Y. A natural sorbent, *Luffa cylindrica* for the removal of a
263 model basic dye. Journal of Hazardous Materials. 2010;179(1-3):658 – 664.

264 5. Hassan LM. Quaternization and Anion Exchange Capacity of Sponge Gourd (*Luffa*
265 *cylindrica*). Journal of Applied Polymer Science. 2006;101:2495–2503.

266 6. Saueprasearsit P, Nuanjaraen M, Chinlapa M. Biosorption of lead (Pb²⁺) by luffa
267 cylindrical fiber. Env. Res. J. 2010;4(1):157-166.

268 7. Ogbonna JC, Tomiyama S, Liu Y, Tanaka H. Efficient production of ethanol by cells
269 immobilized in loofa (*Luffa cylindrica*) sponge. Journal of Fermentation and
270 Bioengineering. 1997;84(3):271–274.

271 8. Roble ND, Ogbonna JC, Tanaka H. L-Lactic acid production from raw cassava
272 starch in a circulating loop bioreactor with cells immobilized in loofa (*Luffa*
273 *cylindrica*). Biotechnology Letters. 2003;25(13):1093-1098.

274 9. Saudagar PS, Shaligram NS, Singhal RS. Immobilization of *Streptomyces*
275 *clavuligerus* on loofah sponge for the production of clavulanic acid. Bioresource
276 Technology. 2008;99(7):2250-2253.

277 10. Kamel BS, Blackman B. Nutritional and oil characteristics of the seeds of angled
278 Luffa (*Luffa acutangula*). Food Chemistry. 1982;9(4):277-282.

279 11. Ansari NM, Houlihan L, Hussain B, Pieroni A. Antioxidant activity of five vegetables
280 traditionally consumed by south-Asian migrants in Bradford, Yorkshire, UK.
281 Phytotherapy research. 2005;19(10):907 – 911.

282 12. Perl M. The biochemical basis of the hypoglycemic effects of some plant extracts. In:
283 Craker LE, Simon JE, editors. Herbs, spices and medicinal plants: Recent advances
284 in botany, horticulture, and pharmacology. Oryx Press, Phoenix, AZ; 1988.

285 13. Raman A, Lau C. Anti-Diabetic Properties and Phytochemistry Momordica charantia
286 L. (Cucurbitaceae). Phytomedicine. 1998;2:349-362.

287 14. Subratty AH, Gurib-Fakim A, Mahomoodally F. Bitter melon: an exotic vegetable
288 with medicinal values, Nutrition & Food Science. 2005;35(3):143 – 147.

- 289 15. Sathishsekar D, Subramanian S. Beneficial effects of *Momordica charantia* seeds in
290 the treatment of STZ-induced diabetes in experimental rats. Biol Pharm Bull.
291 2005;28(6):978-983.
- 292 16. Klomann, SD, Mueller AS, Pallauf J, Krawinkel MB. Antidiabetic effects of bitter
293 gourd extracts in insulin resistant db/db mice. Br J Nutr. 2010;104(11):1613-20.
- 294 17. Oishi Y, Sakamoto T, Udagawa H, Taniguchi H, Kobayashi-Hattori K, Ozawa Y, et
295 al. Inhibition of increases in blood glucose and serum neutral fat by *Momordica*
296 *charantia* saponin fraction. Biosci Biotechnol Biochem. 2007;71(3):735-740.
- 297 18. Kumar Shetty A, Suresh Kumar G, Veerayya Salimath P. Bitter gourd (*Momordica*
298 *charantia*) modulates activities of intestinal and renal disaccharidases in
299 streptozotocin-induced diabetic rats. Mol Nutr Food Res. 2005;49(8):791-796.
- 300 19. Singh J, Adeghate E, Cummings E, Giannikipolous C, Sharma AK, Ahmed I.
301 Beneficial effects and mechanism of action of *Momordica charantia* juice in the
302 treatment of streptozotocin-induced diabetes mellitus in rat. Mol Cell Biochem.
303 2004;261(1-2):63-70.
- 304 20. Yibchok-Anun S, Adisakwattana S, Yao CY, Sangvanich P, Roengsumran S, Hsu
305 WH. Slow acting protein extract from fruit pulp of *Momordica charantia* with insulin
306 secretagogue and insulinomimetic activities. Biol Pharm Bull. 2006;29(6):1126—
307 1131.
- 308 21. Fernandes NP, Lagishetty CV, Panda VS, Naik SR. An experimental evaluation of
309 the antidiabetic and antilipidemic properties of a standardized *Momordica*
310 *charantia* fruit extract. BMC Complement Altern Med. 2007;7(29):7-29.
- 311 22. Jiratchariyakul W, Wiwat C, Vongsakul M, Somanabandhu A, Leelamanit W, Fujii I,
312 et al. HIV Inhibitor from Thai Bitter Gourd. Planta Med. 2001;67(4):350-353.
- 313 23. Jones B. J. Jr, Case V.W. Soil testing and plant analysis. Soil Science Society of
314 America Book Series : 3; 1990.
- 315 24. SAS. SAS System for Windows. SAS Institute, Inc., Cary, NC; 2013.
- 316 25. Kurt TO, Reddy MR, Ravella R, Clarke M. Cover crop residue decomposition and
317 the release of nitrogen to Asian vegetables. Poster presented at Southern Agronomy
318 Branch Annual Conference, 2009; Atlanta, GA.