

# **NITROGEN-USE EFFICIENCY AND ASSIMILATE PARTITIONING FOR GRAIN YIELD IN SELECTED MAIZE INBRED LINES (*Zea mays*. L.) GENOTYPES**

## **ABSTRACT**

Breeding for high grain yield in maize under natural soil Nitrogen condition is of high importance to reduce the heavy usage of chemical fertilizer to enhance productivity and encourage safety of the environment and consumer health. Nine maize inbred lines viz; 1393, KU1414, 5057, 5012, 9030, 9450, SW1, SW6 and SW4, were characterized in a field experiment under natural soil nitrogen 1.81 g/kg. Data were collected on plant growth parameters (plant height, number of leaves, leaf area and stem girth); phenology (percentage emergence, day to anthesis, day to silking and day to green harvest); total nitrogen uptake and yield (plant wet weight, plant dry matter, cob weight and total seed weight). Parameters were paired and correlated to determine their relationships patterns. Positive relationships were obtained from all paired intra-parameter correlations from plant growth, phenology and yield, and from paired inter-parameter correlations between plant growth and yield. Negative relationships were obtained for paired inter-parameter correlations between phenology and growth and between phenology and yield. Total nitrogen uptake gave positive correlations with all parameters. Relative performance grouping analysis revealed that maize genotype 1393 had high N-uptake, high assimilate partitioning and high performance. KU1414, 5057 and 9030 had high N-uptake, low assimilate partitioning and low performance. 5012 had low N-uptake, low partitioning and high performance. 9450 had low N-uptake, low assimilate partitioning and low performance; and SW1, SW6 and SW4 had low N-uptake, high assimilate partitioning and high performance. Hence, 5012 and SW genotypes expressed efficient Nitrogen-use for high grain yield and better growth and phenological performances.

Key words: Nitrogen uptake, Maize, Genotype, Inbred line, Phenology, Correlation analysis, Performance.

---

37

## 38 INTRODUCTION

39 Crop varieties are grown under a wide range of climatic, weather and edaphic conditions (Kling  
40 et al., 1997). Nitrogen nutrient as a component of edaphic factors within an agro-ecology plays a  
41 vital role in the growth, development and yield of agricultural plants. Nitrogen nutrient has been  
42 reported to be a major limiting factor in maize production and grain yield (Below, 2011). This  
43 nutrient shortage reduces crop performance in that it is an essential component of plants nucleic  
44 acids (DNA & RNA) and a major component of chlorophyll, the photosynthetic structure for  
45 carbohydrate formation and an important factor for dry matter production (Doubley, 1995).  
46 Though Nitrogen is relatively abundant in nature, about 78% in the atmosphere and relatively  
47 small proportion of this is readily incorporated into the soil through atmospheric and biological  
48 fixation. Soil gains more Nitrogen from decaying matters and decomposition of domestic and  
49 industrial wastes. Nitrogen is deficient within the agro-ecology owing to its instability and rapid  
50 loss caused by huge transformations (N-cycle) involving mineralization, nitrification,  
51 denitrification, volatilization, run-off and leaching (Below, 2011). N problem is also caused by its  
52 limited availability for plant uptake only in form of Ammonium ( $\text{NH}_4^+$ ) and Nitrate ( $\text{NO}_3^-$ ) out of  
53 other forms such as nitrogen gas ( $\text{N}_2$ ), ammonia ( $\text{NH}_3$ ) and nitrite ( $\text{NO}_2^-$ ). Research revealed N-  
54 nutrient depletion as a serious threat to crop yield and for profitable maize production (Vyn and  
55 Stewart, 1994; Vyn, 2001). Therefore optimum yield is hampered, because poor Nitrogen supply  
56 decreases dry matter accumulation, distribution and mobilization for economic yield (Ciampitti *et*  
57 *al.*, 2012; Ciampitti and Vyn, 2012; 2013). However, supplementary use of chemical fertilizers to  
58 enhance maize productivity is not environment-friendly and high content residue in maize  
59 harvests is not safe for human health. Hence, breeding cultivars with high grain yield under  
60 natural soil Nitrogen condition is necessary. Therefore, this study focused on: (i) assessment of  
61 selected maize inbred lines on Nitrogen use for growth and development; (ii) characterization of  
62 maize lines on source-sink capacity for Nitrogen utilization; (iii) identification of maize lines with  
63 good N-utilization efficiency and better assimilates partitioning for high grain yield.

64

## 65 MATERIALS AND METHODS

66 The research field experiment was carried out on Practical Year Training Programme Farm,  
67 Faculty of Agriculture and Forestry, University of Ibadan (Lat.  $7^{\circ}27'1.7''$  North,  $3^{\circ}53'35.8''$  East;  
68 average max temp.  $33^{\circ}\text{C}$  and min temp.  $28^{\circ}\text{C}$ ; relative humidity 78%; annual rainfall  $>1000\text{mm}$ ).  
69 Soil samples were collected for total Nitrogen content determination using Kjeldahl method after  
70 land clearing at surface and sub-surface at depth range 0-15cm and 15-30cm respectively. Nine  
71 inbred lines as shown in table 1 were planted out in four replicate of Randomized Complete  
72 Block Design and proper agronomic practices were observed. Each line was observed on  
73 varietal responses to N-uptake, mobilization and partitioning and yield performance according to  
74 the laboratory procedure of Kjeldahl.

75 Data were collected on parameters of growth (plant height, number of leaves, leaf area and  
76 stem girth); phenology (percentage emergence, day to anthesis, day to silking and day to green

harvest); total nitrogen uptake and yield (plant wet weight, plant dry matter, cob weight and total seed weight) (Table 3) and subjected to analysis of variance using Statistical Analysis System (SAS, 19.0), the means were separated using Least Significant Difference (LSD) test at 0.05 degree of freedom. The parameters were paired for correlation analysis ( $r = 1/n-1 \sum (X-\mu_x)(Y-\mu_y) / \sigma_X \sigma_Y$ ) to determine their relationship patterns. Relative performance score analysis was used to group inbred lines genotypes based on N-uptake, assimilate partitioning and performance.

## RESULTS AND DISCUSSION

Anthesis, is one of the most important periods for the diagnosis of nutritional status of maize in respect to nitrogen and other mineral nutrients that determines yield generating effect of nitrogen. This came up from the fact that plant nutritional status at this growth stage affects the photosynthetic activity of leaves after anthesis, and consequently, the length of the grain filling stage (Rajcan and Tollenaar, 1999).

From the result of the experiments in Table 2, it is revealed that Line L1 (1393) has the highest stem nitrogen percentage (5.23%) at anthesis followed by line L2 (KU1414) with stem nitrogen percentage (4.47 %) and the least percentage value found in line L8 (SW6) (1.65 %). Line L2 had the highest leaf nitrogen percentage (13.50 %) followed by line L1 (11.72 %) while line L9 (SW4) exhibited the least leaf nitrogen percentage (6.45 %). This indicates that high nitrogen is being partitioned and accumulated in the leaf than in the stem at anthesis, and L1, L2, L3 and L5 has the ability to accumulate nitrogen and partitioned it than other lines tested. Hence, leaf serves as high assimilate bank to remobilize for grain filling at plant productive phase, therefore it is more a factor that determine seed yield in maize plant (Ciampitti and Vyn, 2013). Also, plant yield parameters taken during anthesis and at harvest gave a mean value of 168.89g and 122.6g respectively for all the lines. This revealed that plants had high plant wet weight at anthesis than at harvest; which may be ascribed to high moisture content in the plants due to active physiological processes during anthesis than at harvest stage. Also, plant dry matter at anthesis is relatively low with mean value of 25.00g for all the lines, compared to plant dry matter recorded after harvest with mean value of 56.85g. This was ascribed to high assimilate accumulation in the plants than moisture content at harvest stage, and the trait was expressed in each line with their respective genetic potentials. Plant total seed weight taken after harvesting gave a mean value of 45.99g for the lines, SW4 had highest value of 71.00g and 9450 had least value of 20.66g. This wide gap obtained from the lines indicated a clear cut for individual line genetic capacity to efficiently mobilize assimilates within plant system for grain yield. Correlation analysis as shown in Table 3 revealed that intra-parameter correlations from plant growth, phenology and yield gave positive relationships and significant correlations at  $p=0.05$  as indicated between respective plant height and number of leaf with coefficient value  $r=0.5185$  (Fig. 1), days to silking and days to anthesis with coefficient value  $r=0.7094$  (Fig. 2) and plant wet weight and cob weight with coefficient value  $r=0.7797$  (Fig. 3). Inter-parameter correlations from plant growth and phenology gave negative relationships as indicated between leaf area and days to silking with coefficient value  $r=-0.7819$  (Fig. 4); inter-parameter correlations

from plant phenology and yield gave negative relationships as indicated between days to anthesis vs plant dry matter with coefficient value  $r=-0.3269$  (Fig. 5), and inter-parameter correlations from plant growth and yield gave positive relationships as indicated between cob weight vs plant height with coefficient value  $r=0.5106$  (Fig. 6). Correlation analyses revealed an increase in value of one paired of intra-parameters from growth, phenology and yield, and between one paired of inter-parameters from growth and yield will leads to proportional increase in the value of the other. Thus, negative and inverse relationships between pairs of inter-parameters from phenology and growth, and from phenology and yield indicated that increase in value of one paired parameters will leads to proportional decrease in the value of the other.

Relative performance score analysis for genotypes characterization (Table 4) revealed that genotype 1393 had high N-uptake, high assimilate partitioning and high performance values. This indicated a disadvantage(d) at high nitrogen demand. KU1414, 5057 and 9030 had high N-uptake, low assimilate partitioning and low performances. This indicated advantage at low nitrogen demand and disadvantage at poor performances. Genotype 9450 had low N-uptake, low partitioning and low performance,this indicated poor performances. Genotype 5012 had low N-uptake, low partitioning and high performance,this indicated advantage at low nitrogen demand and good performance. SW1, SW6 and SW4 had low N-uptake, high partitioning and high performance,this indicates a low nitrogen demand and good performance.

## CONCLUSION

Breeding and characterization of genotypes for high grain yield under natural soil nitrogen condition is a feasible procedure and promising development to avert supplementary use of chemical fertilizers to enhance maize productivity and profitable production. This study identified inbred lines 5012, SW1, SW6 and SW6 as good genotypes with efficient Nitrogen-use for high grain yield and good performance. These genotypes are recommended for use at improving farmers cultivated varieties to encourage maize cultivation with zero use of chemical fertilizers in order to preserve our environment and safeguard consumers' health.

## REFERENCES

- Below, M. I., 2011.The Quest for 20-Ton Maize.'Buzios, Brazil. Sept. 19, 2011. XIX
- Below, M. I., 2011.7 Ways to Push YourYield Barrier.' Barrie, Ontario, Canada. Jan. 4-6, 2012. XIII Brazilian Congress of Plant Physiology.
- Ciampitti, I.A., and Vyn ,T.J. (2013). [Grain nitrogen sources over time in maize: a review. Crop Sci. 53: 366-377.](#)
- Ciampitti, I. A., H. Hao, P. Friedemann, and Vyn,T.J. (2012) . [Potential physiological frameworks for mid-season field phenotyping of final plant N uptake, N use efficiency and grain yield in maize.](#) Crop Science 52: 2728-2742.

- 156  
157 Ciampitti, I.A., and Vyn,T.J. (2012). Physiological perspectives of changes over time in  
158 maize yield dependency on nitrogen uptake and associated nitrogen efficiencies: A  
159 review. Field Crops Res. 133: 48-67.  
160 <http://www.sciencedirect.com/science/article/pii/S0378429012001013>  
161  
162 Doebley, J. F., Stec, A., and Gustus, C. (1995) .The molecular genetics of crop  
163 domestication.Genetics 141 (1), 333 461.
- 164 Kling, J. G. and Gregory, E. (1997). Morphology and growth of maize. IITA/CIMMYT Research  
165 Guide 9.Int'l Institute of Tropical Agriculture.[On-Line}. Available at  
166 [http://www.iita.org/cms/details/trn\\_mat/irg9/irg9.htm](http://www.iita.org/cms/details/trn_mat/irg9/irg9.htm) [URL accessed July2010].
- 167 Rajcan, I. and Tollenaar, M. (1999). Source: sink ratio and leaf senescence in maize: II.Nitrogen  
168 metabolism during grain filling. Field Crops Res., 60: 255-265.
- 169  
170 Stanton, W. R. (1961). Foodcrop preferences: their significance to plant breeding programs and  
171 agricultural research policy. W. Afr. J. Biol. Chemistry 4, 84-89.
- 172 Vyn, T.J. (2001).Nutrient placement and high yield management in corn.InfoAg 2001  
173 Conference Proceedings, Aug. 7-9, Indianapolis, IN.
- 174 Vyn, T.J. and Stewart. G.A. (1994). Cover crop nitrogen availability to corn with conservation  
175 tillage. Pages 813-818 in Proceedings of 13 th International Soil Tillage Research Organization  
176 Conference, Aalborg, Denmark, July 24-29.

177

178

179

180

181

182

183

184

185

186

187

188 Table 1. List of the maize inbred line seeds and their description

| 189 | Lines | Source  | Name   | Type   |
|-----|-------|---------|--------|--------|
| 190 | L1    | IITA    | 1393   | White  |
| 191 | L2    | IITA    | KU1414 | Yellow |
| 192 | L3    | IITA    | 5057   | White  |
| 193 | L4    | IITA    | 5012   | White  |
| 194 | L5    | IITA    | 9030   | White  |
| 195 | L6    | IITA    | 9450   | Yellow |
| 196 | L7    | LAUTECH | SW1    | Yellow |
| 197 | L8    | LAUTECH | SW6    | White  |
| 198 | L9    | LAUTECH | SW4    | White  |

199

200

201

202

203

204

205

Table 2. Percentage Nitrogen uptake at anthesis and yield parameters of the maize Inbred lines taken at anthesis and after green harvest.

| Lines | At Anthesis |       |       |        |       | At Harvest |       |        |       |       |       |
|-------|-------------|-------|-------|--------|-------|------------|-------|--------|-------|-------|-------|
|       | SN%         | LN%   | TNU   | PWW    | PDM   | PWW        | PDM   | CBW    | CBL   | CBD   | TSW   |
| L1    | 5.23        | 11.72 | 16.95 | 159    | 23.00 | 208.00     | 92.00 | 106.00 | 16.20 | 16.20 | 57.00 |
| L2    | 4.47        | 13.50 | 17.97 | 170    | 25.00 | 117.60     | 48.50 | 65.20  | 14.30 | 14.30 | 41.33 |
| L3    | 3.67        | 10.04 | 13.71 | 150    | 20.00 | 72.00      | 34.66 | 44.00  | 14.60 | 14.60 | 24.66 |
| L4    | 3.75        | 8.26  | 12.01 | 155    | 30.00 | 88.80      | 44.66 | 69.00  | 14.60 | 14.60 | 58.66 |
| L5    | 2.54        | 10.81 | 13.35 | 151    | 18.00 | 89.30      | 39.33 | 21.00  | 9.10  | 9.10  | 31.00 |
| L6    | 2.26        | 9.98  | 12.24 | 160    | 26.00 | 99.70      | 34.50 | 53.30  | 14.50 | 14.50 | 20.66 |
| L7    | 1.96        | 8.68  | 10.64 | 200    | 35.00 | 158.80     | 74.00 | 80.40  | 16.00 | 16.00 | 56.66 |
| L8    | 1.65        | 6.66  | 8.31  | 165    | 24.00 | 119.50     | 64.00 | 97.50  | 14.30 | 14.30 | 52.66 |
| L9    | 1.87        | 6.45  | 8.33  | 165    | 24.00 | 150.00     | 80.00 | 103.6  | 16.30 | 16.30 | 71.33 |
| Mean  | 3.04        | 9.56  | 12.61 | 163.89 | 25.00 | 122.60     | 56.85 | 71.11  | 14.40 | 14.40 | 45.99 |

SN%-Stem nitrogen percentage      LN%-Leaf nitrogen percentage  
PWW- Plant wet weight      PDM- Plant dry matter  
CBW- Cob weight      CDM- Cob drymatter  
CBL-Cob length      TSW-Total seed weight

239

240

241 Table 3: Correlation matrix of growth, phenology, nitrogen uptake and yield parameters  
242 for the lines.

| 243 |     | PHT    | LA      | LN     | CL     | DA     | DS     | DH     | PWW    | PDM   | CW     | TN    | SW |
|-----|-----|--------|---------|--------|--------|--------|--------|--------|--------|-------|--------|-------|----|
| 244 | PHT | -      |         |        |        |        |        |        |        |       |        |       |    |
| 245 | LA  | .5736  | -       |        |        |        |        |        |        |       |        |       |    |
| 246 | LN  | .5185  | .3208   | -      |        |        |        |        |        |       |        |       |    |
| 247 | CL  | .5106  | .3506   | .0051  | -      |        |        |        |        |       |        |       |    |
| 248 | DA  | -.5267 | -.502   | .0003  | -.2773 | -      |        |        |        |       |        |       |    |
| 249 | DS  | -.4205 | -.7818* | -.1582 | -.1158 | .7094* | -      |        |        |       |        |       |    |
| 250 | DH  | -.4055 | -.3048  | .0038  | .0246  | .5941  | .4223  | -      |        |       |        |       |    |
| 251 | PWW | .1291  | .4620   | .2274  | .3214  | -.0174 | -.1098 | .0019  | -      |       |        |       |    |
| 252 | PDM | .5387  | .4060   | .0038  | .2950  | -.3269 | -.0364 | -.2319 | .0701  | -     |        |       |    |
| 253 | CW  | .5106  | .8987*  | .0951  | .6242  | -.3649 | -.3895 | .0071  | .7797* | .1300 | -      |       |    |
| 254 | TN  | .4506  | .4763   | .0424  | .1276  | .7602* | .6009  | .6948* | .1009  | .2478 | .2709  | -     |    |
| 255 | SW  | .7875* | .9269*  | .6426  | .5299  | .6170  | .8611* | .2958  | .6282  | .4276 | .8122* | .3593 | -  |

256 \* Correlation is significant at p=0.05 level

257

258

259

260

261 Table 4.Relative Performance Score obtained from lines characterization on plant growth,  
262 phenology, Nitrogen content and yield parameters.

263

| Lines | PHT | LN  | LA | SGT | DAT | DSK | DGH | EM% | PWA | PDA | PWH | CBW | CBH | SN% |     |   |     |
|-------|-----|-----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---|-----|
| LN%   | TSW | RMS |    |     |     |     |     |     |     |     |     |     |     |     |     |   |     |
| L1    | -   | +   | +  | +   | -   | +   | -   | -   | -   | +   | +   | +   | +   | +   | 10* |   |     |
| L2    | -   | +   | -  | +   | -   | -   | -   | +   | +   | -   | -   | -   | -   | +   | +   | - | 6   |
| L3    | -   | -   | -  | -   | -   | -   | -   | -   | -   | -   | -   | -   | +   | +   | +   | - | 3   |
| L4    | +   | -   | +  | +   | +   | +   | +   | -   | -   | +   | -   | -   | +   | +   | -   | - | 9*  |
| L5    | -   | +   | -  | +   | -   | -   | +   | -   | -   | -   | -   | -   | -   | -   | +   | - | 4   |
| L6    | -   | -   | -  | -   | +   | -   | +   | -   | -   | +   | -   | -   | +   | -   | +   | - | 5   |
| L7    | +   | +   | +  | -   | +   | -   | +   | +   | +   | +   | +   | +   | -   | -   | -   | + | 12* |
| L8    | +   | -   | +  | -   | +   | +   | +   | +   | +   | -   | -   | +   | -   | -   | -   | + | 9*  |
| L9    | +   | +   | +  | +   | +   | +   | +   | -   | +   | +   | -   | +   | +   | -   | -   | + | 12* |

264 **PHT**-Plant height, **LN**-Leaf number, **LA**-Leaf area, **SGT**-Stem girth, **DAT**-Days to anthesis, **DSK**-

265 Days to silking, **DGH**-Days to green harvest, **EM%**-Percentage emergence, **PWA**-Plant weight

266 at anthesis, **PDA**-Plant dry matter at anthesis, **PWH**-Plant weight after harvesting, **CBW**-Cob

267 weight, **CBH**-Cob height, **SN%**-Stem nitrogen percentage, **LN%**-Leaf nitrogen percentage,

268 **TSW**-Total seed weight

269 + represents values above mean (8.78)

270 - represents values below the mean

271 \* represents performance of lines above general average

272




**Fig. 1:** Correlation graph obtained from plant height and leaf number parameters for the lines.  
 $r$  = correlation coefficient

277




**Fig. 2:**Correlation graph obtained from days to anthesis and days to silking parameters for the lines.**r** = correlation coefficient

281




**Fig. 3:** Correlation graph obtained from plant wet weight and cob weight parameters for the lines.  
 $r$  = correlation coefficient




**Fig. 4:** Correlation graph obtained from leaf area and days to silking parameters for the lines.  
 $r$  = correlation coefficient




**Fig. 5:**Correlation graph obtained from day to anthesis and plant dry matter parameters for the lines.  
**r** = correlation coefficient




**Fig. 6:**Correlation graph obtained from plant height and cob weight parameters for the lines.  
**r** = correlation coefficient