

**The effect of macro and micronutrient fertilizers on the growth and yield of rice under
calcareous soils**

Majid Qasempour Alamdari

Department of agronomy, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran.

Postal Address: Flat No. 22, Ahmadi Nezhad Alley, Komeyl Street, Bahre-Eram Sharghi,
Babol, Mazandaran, Iran.

Telephone and Mobile No: +98 111 2253228 , +98 911 117 3013

Email address of the corresponding author: ghasempour3m@yahoo.com

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62

Abstract

In order to examine the effect of macro and micronutrient fertilizers on the growth and yield of rice (*Oryza sativa* L.) under calcareous soils, the experiment carried out as split- split plot design in randomized complete block with three replications. Macro, sulfur powder and micronutrient fertilizers have chosen as main, sub and sub-sub plots, respectively. The amount of lime and pH of the experimental soil is 16.7% and 7.54, respectively that categorized as calcareous soils. The aim of the present study is decrease soil pH at the five stages of rice plant growth and investigation of vegetative growth and rice grain yield. The results showed that soil pH decreased by using NPK+S+Zn fertilizers (0.32) compared to the control plot. With decreasing soil pH all the agronomic characteristics of rice plant and its grain yield increased significantly, except 1000-grain weight. The number of filled grain (119) and total number of seed per panicle (127) and grain yield (4376 kg ha^{-1}) reached at the maximum values when NPK+S+Zn fertilizers applied together. Rice grain yield was increased 1483 kg ha^{-1} by this treatment compared to the control plot (2893 kg ha^{-1}). Thus, this treatment can be resulted as the best treatment under calcareous soils conditions with high soil pH.

Keywords: Macro and micronutrient, Calcareous soil, Growth, Yield, Rice.

63

64 **1. Introduction**

65 The two greatest problems faced by the third world countries, at present, are the exponential
66 increase in population and the basic need for providing adequate food. With the increasing
67 population each year, the demand for foodstuff also increases and it will have to be balanced
68 obviously by a qualitative and quantitative increase in food production. The food production
69 in turn can be improved only by improving the conditions of arable soil and the correct use of
70 fertilizers. More than half of arable soils of the world are belonged to alkaline or calcareous
71 soils. The presence of excess amount of lime or Na element in these soils disorders the
72 nutrition uptake especially micronutrient elements and finally injures plant growth [2 and 14].
73 On the other hand, the anomalous use of nitrogen and phosphorous fertilizers are common for
74 growing high yield varieties of crops but it has been found to decrease the uptake capability
75 of plants for micronutrients. There is deficiency of micronutrients such as Zn, Cu, Mn and Fe
76 in agronomic soils of most arable soils on the world because of the excessive use of nitrogen
77 and phosphate fertilizers and the existence of calcareous and alkaline soils. Some of the
78 micronutrients such as Zn, Cu and Mn, are key elements to remedy of human and animal
79 diseases. The deficiency of these micronutrients can cause chronic malnutrition in developing
80 countries where malnutrition is a major problem. Micronutrient malnutrition has become a
81 global problem of immense proportions. Incredibly, over 3 billion people are afflicted with
82 micronutrient malnutrition and the numbers are increasing [10]. Furthermore, certain
83 micronutrient deficiencies are an issue even in developed countries such as the United States.
84 The uptake of Zn, Cu, Mn and Fe elements by plants is retarded if the soil is alkaline and is
85 limy [15]. Generally, alkaline soils are deficient in micronutrients compared to the acidic
86 soils. About 30% of the agricultural soils of the world face the lack of these elements. In
87 addition to the frequent use of super-phosphate fertilizers, the plantation at short intervals

also causes a decrease in the density of micronutrients in arable soils. The plant growth is also affected in soils deficient in micronutrients even the use of N, P and K fertilizers cannot help in increasing the grain yield. The rice plant is special to the tropical countries, and care should be taken in choosing the fertilizer and the method of its use. Amongst the recognized macronutrients, like N, P, K, Ca, S and Mg the rice plant needs more N, P and K. This plant is susceptible to Zn and Cu deficiency, more than other micronutrients.

Next to wheat most of arable soil of the world is used for rice growing. Suitable growth of this plant is under the acidic soil pH (5.5 to 6.5) condition, where most nutrients can be taken up by the plant especially micronutrients [6]. Fajeria et al [3] reported that the best rice yield is obtained in the soil that has pH about 6 and it decreases in higher pH. At that pH (6.0), the absorption of P, Fe, Zn and Mn by rice plant also increased [9]. The pH of soil decreases by addition of sulfurous and nitrogen fertilizers [1]. The enzyme urease rapidly hydrolyzes the commonly used nitrogen fertilizers like urea. Acidification occurs when nitrifying bacteria oxidizes the NH_4^+ and NH_3 . The NO_3^- and H^+ ions produced cause decrease in the soil pH. The addition of 100 kg ha^{-1} of ZnCl_2 also showed an increase in the rice yield [18]. Other experiments carried out by Malakoti and Tehrani [8] and Fallah and Saadati [4] who reported that with using Zn and Cu fertilizers grain yield increases. The main objective of the present study is to examine the effect of macronutrients such as N, P and K and the effect of sulfur powder on soil pH and micronutrients such as Zn, Cu and Mn and its relationship with soil pH on growth and the yield and yield components of rice.

2. Material and Methods

The present experimental soil has adequate Ca^{2+} available in the form of CaCO_3 and has been categorized as calcareous soil containing sufficient lime (16.7%) with soil pH (7.62). E.C, S.A.R, E.S.P, CEC, OC and OM % of this soil and P, K, Zn, Cu, Mn and Fe content in the experimental soil are shown in table 1. Fe content of the experimental soil was more than

critical level, hence it has not been added in the soil. Texturally it is sandy-loam course textured with low clay content. The experiment was carried out as split-split plot design in randomized complete block with three replications. There were 3 factors in this experiment which are Macro-elements as main plot in two levels (1-control and 2-NPK-fertilizers as basic fertilizer) and sulfur as sub plot in two levels (1-control and 2-S-powder) and Micro-elements as sub-sub plot in five levels (1-control, 2-Zn, 3-Cu, 4-Mn and 5-Zn,Cu,Mn-sulfate fertilizers). Thus, 20 treatments were located in each plot and each of these, were replicated three times (60 plots). Basic fertilizer (NPK-fertilizers), microelement fertilizers and sulfur powder were applied to the plots area as recommended by routine soil analysis [7]. Each plot trampled and mixed with fertilizers in to the soil for balance and perfect distribution of fertilizers. N, P and K-fertilizers were as Urea (50 kg ha^{-1}), Tripl super- phosphate (50 kg ha^{-1}) and potassium sulfate (25 kg ha^{-1}), respectively. Zn, Cu and Mn-fertilizers were as Zn sulfate (30 kg ha^{-1}), Cu sulfate (10 kg ha^{-1}) and Mn sulfate (20 kg ha^{-1}), respectively. Sulfur powder has also been chosen at the rate of 100 kg ha^{-1} . The rice plant seeds sowed on the nursery field after germinating. Afterwards, the rice seedling was transplanted on the selected plots when it was at 15 cm height. Distance of transplanting hill was $25 \times 25 \text{ cm}$ and was cultivated 4 stems in each hill. The type of rice variety was Taroom with growth period 90 days, which belongs to the high quality types as Basmati varieties. The field was irrigated and flooded four days after transplanting and other agronomic operations were done by routine and general methods.

During the rice plant growth the soil pH was measured at five stages; one week after transplanting, at the maximum tillering stage (30 days after transplanting), at the leaf tube formation stage, at the panicle initiation stage and at the seed maturity stage. To measure soil pH, 5-soil samples were collected per plots and mixed and 1 kg of these has chosen to measure soil pH. To determine soil pH, 100 cc distilled water was added to 10 gm of soil

sample and mixed. After 24 hours, the pH of soil solution was recorded with pH-meter having temperature compensation adjustment. The agronomic characteristic of rice plant were also measured during the rice plant growth that are the number of tillers per hill at the maximum tillering stage (30 days after transplanting), plant height at panicle initiation stage and seed maturity stage, panicle length, the number of filled grain per panicle, the total number of seed per panicle. After reaching the physiological maturity, the plants were harvested within the central 2 meters of each plot to avoid contamination from the adjacent plot. The harvesting was done at 50 cm height above the soil surface. The combine machine separated the grains, and then 1000-grain weight was calculated. Afterwards, the grain humidity was also measured and brought to the level of 14% humidity and its yield /ha was calculated. Data were analyzed by the analysis of variance technique (ANOVA) and the mean differences were adjudged by Duncans' new multiple range tests (DNMRT); [5]. Microsoft Excel software has been used for statistical analysis of data.

Table 1. Soil analysis of experimental field at 0-30 cm depth

pH	EC (mmohs/cm)	SAR (%)	ESP (%)	CEC (meq/lit)	OC (%)	OM (%)
7.62	1.12	12.1	9.5	21.9	2.43	4.17
P (ppm)	K (ppm)	Zn (ppm)	Cu (ppm)	Mn (ppm)	Fe (ppm)	Soil Texture
16	240	4.8	0.5	21.4	92.2	Sandy-Loam

3. Results and Discussion

The results showed that the macronutrient fertilizers affected on the plant height at both stages (panicle initiation and seed maturity) and panicle length very significantly. It also influenced on the other agronomic characteristics of rice plant significantly (Table 2). Sulfur powder had very significant influence on total mean soil pH, plant height at panicle initiation and grain yield. It had also significant effect on the plant height at seed maturity stage, the number of filled grain per panicle and the total number of seed per panicle. Table 2 shows

that the effect of micronutrient fertilizers was very significant on all the agronomic traits of rice plant except panicle length, total number of seed per panicle and 1000-grain weight. Sulfur powder and micronutrient fertilizers interaction was very significant on mean soil pH, total number of tillers and grain yield. An interaction effect of macro, micronutrient and sulfur powder fertilizers was also significant on plant height at both stages and grain yield. Sulfur powder, macronutrient fertilizers and their interaction affected on the grain yield significantly, while sulfur powder, micronutrient fertilizers and their interaction affected on the grain yield very significantly. According to the obtained results by analysis of variance of data, the most agronomic characteristics of rice plant and its grain yield were statistically influenced using macro, micronutrient and sulfur powder fertilizers, separately or together. It shows that adding the sulfur or sulfate fertilizers into the calcareous soils is effective to reach lower soil pH, higher total number of grain per panicle and finally more rice grain yield.

UNDER PEER REVIEW

Table 2. Analysis of variance of macro and micronutrient fertilizers on the measured characteristics at the present study.

Treatments	df	Total mean soil pH at five stages	Total number of tillers	Plant height at panicle initiation	Plant height at seed maturity	Panicle length	Filled grain per panicle	Total seed per panicle	1000 grain weight	Grain yield
Replication	2	0.003 ^{ns}	2.1462 ^{ns}	3.5167 ^{ns}	3.6932 ^{ns}	0.0347 ^{ns}	25.867 ^{ns}	53.117 ^{ns}	0.5962 ^{ns}	24043 ^{ns}
NPK(a)	1	0.0564*	175.79*	660.02**	734.3**	37.131**	806.67*	576.6*	10.004*	4E+06*
E (a)	2	0.0021	9.2562	0.2167	2.7062	0.3307	22.067	6.45	0.4432	185673
S(b)	1	0.1179**	9.6802 ^{ns}	170.02**	15.302*	1.35 ^{ns}	166.67*	135*	1.5682 ^{ns}	2E+06**
NPK*S	1	0.0003 ^{ns}	0.0002 ^{ns}	88.817**	47.348**	0.096 ^{ns}	96.267*	19.267 ^{ns}	1.3202 ^{ns}	219736*
E(b)	4	0.0028	4.6662	0.2667	1.8073	0.382	8.7667	9.6833	1.6807	18182
Zn,Cu,Mn (c)	4	0.012**	2.9639**	20.375**	16.965**	1.2748 ^{ns}	37.9**	25.358 ^{ns}	0.4211 ^{ns}	194265**
NPK*Zn,Cu,Mn	4	0.0077**	0.9028 ^{ns}	2.5583*	9.0422*	0.4711 ^{ns}	9.0833 ^{ns}	8.0583 ^{ns}	0.2371 ^{ns}	32603 ^{ns}
S*Zn,Cu,Mn	4	0.0078**	3.2923**	0.3917 ^{ns}	3.9402 ^{ns}	0.1796 ^{ns}	5.4167 ^{ns}	4.4583 ^{ns}	0.1103 ^{ns}	107672**
NPK*S*Zn,Cu,Mn	4	0.01**	0.4747 ^{ns}	3.1083*	12.444*	0.3506 ^{ns}	7.1 ^{ns}	12.392 ^{ns}	0.1556 ^{ns}	74039*
E (c)	32	0.0017	0.6655	0.8583	3.275	0.5171	7.825	12.942	0.6066	19530
C.V(%)	---	1.12	13.64	3.66	2.75	3.99	4.7	4.2	3.6	10.05

* and ** are significant and very significant at the level of P=0.05 and P=0.01, respectively in F-test. Ns is not significant.

Table 3 shows that the all used treatments had significant difference on all the agronomic characteristic of rice plant. Mean comparison of data at the present study shows that soil pH at the five stages decreased using macro, sulfur powder and micronutrient fertilizers during plant growth period. This is in agreement with earlier workers by Slaton et al. [12] and Sharma and Swarup [11]. The minimum soil pH is related to treatment no.20 when macro (NPK), micronutrient (Zn,Cu and Mn) and sulfur powder fertilizers applied together. Soil pH decreased 0.35 unit by treatment no. 20 (NPK+S+Zn,Cu,Mn) and 0.32 unit by treatment no. 17 (NPK+S+Zn fertilizers) compared to the control plot. The maximum number of tillers (18.8), plant height (at panicle initiation 124cm and seed maturity stage 164cm) and panicle length (28cm) was obtained by treatment no.20. The all used treatments decreased 1000-grain weight. The minimum of its value was obtained by treatment no. 20 (23.5 g). The number of filled grain (118) and total number of seed per panicle (126) and grain yield (4376 kg ha^{-1}) were reached to the maximum values when macro (NPK), micronutrient (Zn) and sulfur powder fertilizers applied together (treatment no. 17). Rice grain yield was increased 1483 kg ha^{-1} by this treatment compared to the control plot (2893 kg ha^{-1}). It is because of increasing the number of filled grain and total number of seed per panicle at this treatment. On the other hand, the number of tillers per hill and panicle length at treatment no. 17 is the similarly at the level of $P=0.05$ compared to treatment no. 20, statistically. It shows that all of the agronomic characteristics of the rice plant at treatment nos. 17 and 20 are the same values, statistically, except plant height. According to the obtained results, treatment no. 17 (use of NPK+ S + Zn-sulfate fertilizers), can be concluded as the best treatment. The production expense will be declined using this treatment because of non using of Cu and Mn fertilizers in it as compared to treatment no. 20.

Table 3. Mean comparison of the effect of macro and micronutrient fertilizers on the measured characteristics at the present study.

Treatments	Total mean soil pH	Number of tillers	Plant height at panicle initiation/cm	Plant height at seed maturity/cm	Panicle length/cm	Filled grain per panicle	Total seed per panicle	1000 grain weight/gr	Grain yield/kg ha ⁻¹
1-Witness (control)	7.54 ⁿ	12.5 ^o	110 ^{i-m}	148 ^l	24.6 ^{kl}	101 ^l	109 ^k	25.5 ^a	2893 ^s
2-Zn	7.37 ^m	14.5 ^{hij}	111 ^{h-l}	153 ^{e-i}	25.8 ^{ghi}	107 ^{f-k}	117 ^{e-i}	25.0 ^{ab}	3310 ^{pq}
3-Cu	7.29 ^{d-h}	13.9 ^{j-m}	110 ^{i-m}	152 ^{f-j}	25.7 ^{g-j}	108 ^{f-j}	116 ^{e-j}	24.5 ^{a-d}	3266 ^{pqr}
4-Mn	7.33 ^{g-l}	13.8 ^{j-n}	112 ^{g-k}	154 ^{d-h}	25.8 ^{ghi}	109 ^{e-i}	118 ^{d-h}	24.6 ^{abc}	3420 ^p
5-Zn,Cu,Mn	7.29 ^{d-h}	14.3 ^{h-k}	114 ^{f-i}	155 ^{d-g}	26.4 ^{gh}	110 ^{d-h}	118 ^{d-h}	24.3 ^{a-e}	3816 ^{e-l}
6-S	7.29 ^{d-h}	15.0 ^j	112 ^{g-k}	151 ^{g-k}	25.6 ^{g-k}	106 ^{g-l}	116 ^{e-j}	24.3 ^{a-e}	3766 ^{f-o}
7-S+Zn	7.24 ^{b-d}	15.5 ^h	113 ^{f-j}	152 ^{f-j}	25.8 ^{ghi}	107 ^{f-k}	118 ^{d-h}	24.3 ^{a-e}	3880 ^{c-i}
8-S+Cu	7.25 ^{c-e}	14.0 ^{i-l}	111 ^{h-l}	151 ^{g-k}	25.7 ^{g-j}	108 ^{f-j}	118 ^{d-h}	24.3 ^{a-e}	3840 ^{d-j}
9-S+Mn	7.28 ^{d-g}	14.0 ^{i-l}	113 ^{f-j}	152 ^{f-j}	25.5 ^{g-l}	107 ^{f-k}	117 ^{e-i}	23.8 ^{b-g}	3833 ^{d-k}
10-S+Zn,Cu,Mn	7.30 ^{e-i}	14.5 ^{hij}	114 ^{f-i}	153 ^{e-i}	26.7 ^{b-g}	111 ^{d-g}	119 ^{d-g}	24.0 ^{b-f}	3776 ^{f-n}
11-NPK	7.31 ^{f-j}	15.3 ^{hi}	115 ^{e-h}	159 ^{bc}	27.1 ^{a-f}	110 ^{d-h}	120 ^{c-f}	23.6 ^{b-i}	3800 ^{e-m}
12-NPK+Zn	7.32 ^{f-k}	17.8 ^{a-d}	115 ^{e-h}	157 ^{b-e}	27.2 ^{a-e}	113 ^{c-e}	121 ^{b-e}	23.8 ^{b-g}	4016 ^{b-g}
13-NPK+Cu	7.27 ^{c-f}	17.1 ^{b-g}	115 ^{e-h}	156 ^{c-f}	27.1 ^{a-f}	111 ^{d-g}	121 ^{b-e}	23.6 ^{b-i}	3966 ^{c-h}
14-NPK+Mn	7.28 ^{d-g}	17.7 ^{a-e}	116 ^{d-g}	158 ^{bcd}	27.1 ^{a-f}	112 ^{c-f}	121 ^{b-e}	23.6 ^{b-i}	4016 ^{b-g}
15-NPK+Zn,Cu,Mn	7.33 ^{g-l}	18.1 ^{abc}	117 ^{def}	157 ^{b-e}	27.2 ^{a-e}	114 ^{b-d}	121 ^{b-e}	23.7 ^{b-h}	4056 ^{b-f}
16-NPK+S	7.23 ^{bc}	18.2 ^{ab}	119 ^{cde}	159 ^{bc}	27.3 ^{a-d}	116 ^{abc}	123 ^{a-d}	23.6 ^{b-i}	4100 ^{b-e}
17-NPK+S+Zn	7.22 ^b	17.6^{a-f}	120 ^{bcd}	160 ^b	27.2^{a-e}	119^a	127^a	23.8 ^{b-g}	4376^a
18-NPK+S+Cu	7.22 ^b	17.8 ^{a-d}	122 ^{abc}	159 ^{bc}	27.5 ^{abc}	118 ^{ab}	123 ^{a-d}	23.6 ^{b-i}	4150 ^{abc}
19-NPK+S+Mn	7.23 ^{bc}	17.6 ^{a-f}	123 ^{ab}	159 ^{bc}	27.7 ^{ab}	118 ^{ab}	125 ^{abc}	23.6 ^{b-i}	4126 ^{a-d}
20-NPK+S+Zn,Cu,Mn	7.19^a	18.8^a	124^a	164^a	28.0^a	118 ^{ab}	126 ^{ab}	23.5 ^{c-j}	4283 ^{ab}

A given means per each column with the same letters, have not significant difference, statistically ($P=0.05$).

Table 4 shows that the use of Zn,Cu and Mn fertilizers increased grain yield 923 kg ha⁻¹. This observation is in agreement with earlier worker by Surajit and Datta [16]. The results showed that by adding only sulfur powder, grain yield increased 873 kg ha⁻¹ compared to the control plot. This observation is in agreement with the earlier workers [2 and 3]. Saadati [13] and Fallah and Saadati [4] showed that 200-400 kg ha⁻¹ of sulfur addition increased the rice yield by 500 kg ha⁻¹. Turner [17] has reported that plots treated with sulfuric acid out-yielded those treated with nitrogen and other nutrients. It seems that lowering of soil surface pH from 7.5 to 6.2 raised the rice crop yield by about 1000 kg ha⁻¹. Using NPK fertilizers also increased grain yield 907 kg ha⁻¹ compared to the control plot. When NPK and Zn,Cu,Mn fertilizers applied into the soil together, grain yield increased 1163 kg ha⁻¹ as compared with control plot and when NPK, S and Zn,Cu,Mn fertilizers applied together, grain yield increased 1400 together as compared with control plot. It shows that NPK, sulfur powder and Zn,Cu,Mn fertilizers increases grain yield separately, but when all of those use into the soil together, grain yield will be increased significantly. Surajit and Datta [16] had reported that use of NPK fertilizers without addition of Zn decreased the rice yield, while addition of zinc oxide or zinc sulfate increases the yield of rice. It was reported that yield increased by 850 kg ha⁻¹. Malakoti and Tehrani [8] and Fallah and Saadati [4] had also reported that the addition of potassium chloride (as macronutrient), zinc sulfate and copper sulfate as fertilizers revealed that the rice yield increases by 340, 1070 and 600 kg ha⁻¹, respectively.

Table 4. Mean comparison of macro, sulfur and micronutrient fertilizers on rice grain yield

Treatments	Grain yield (kg ha ⁻¹)	Increase relating witness
Witness	2893	-----
Zn,Cu,Mn	3816	923
S	3766	873
NPK	3800	907
S+Zn,Cu,Mn	3776	883
NPK+Zn,Cu,Mn	4056	1163
NPK+S	4100	1207
NPK+S+Zn,Cu,Mn	4283	1400

4. Conclusion

The obtained results showed that using macro, micronutrient and sulfur powder fertilizers into the calcareous soils with high soil pH and calcium carbonate content is effective. With decreasing soil pH all the agronomic characteristics of rice plant and its grain yield increased significantly, except 1000-grain weight. The number of filled grain and total number of seed per panicle and grain yield were reached at the maximum values when NPK+ S + Zn-sulfate fertilizers (treatment no. 17) applied together. Thus, this treatment can be resulted as the best treatment under calcareous soils conditions.

5. Acknowledgment

The author is thankful from all person or institutes who helped him to do the present study.

6. References

1. Benbi DK, Nieder R. Handbook of processes and modeling in the "Soil-Plant System". 1st ed. The Haeorth Press,Inc. USA; 2004.
2. Das DK. Introductory Soil Science. 1st ed. Kalyani Publ. New Delhi; 1996.
3. Fajerla NK, Baligar VC, Jones CA. Growth and crop plants nutrition. Translated by Fathi, Gh. Publ. Mashhad Jahad Uni; 1999. (In Persian)
4. Fallah V, Saadati N. Management of fertilizer using in paddy field of Mazandaran. Publ. Country Rice Research Inst. Mazandaran. Amol; 1997. (In Persian)

- 280 5. Gomez KA, Gomez AA. Statistical procedures for agricultural research. 2nd ed. John
281 Wiley sons, New York, Chichester, Bribane, Toronto, Singapore; 1984.
- 282 6. Qasempour AM, Khodabandeh N. Rice cultivation. Publication Islamic Azad University,
283 Qaemshahr, Iran; 2004. (In Persian).
- 284 7. Malakoti MJ, Gheybi MN. Determination of critical level of strategies agricultural
285 nutrients and correct recommendation of fertilizers in country. In: Proceeding of the 1st
286 National congress of reduction of poisons consumption and suitable use of chemical
287 fertilizers in agriculture. Agricultural Ministry, Karaj, Iran; 1997. (In Persian).
- 288 8. Malakoti MJ, Tehrani MM. Effects of Micronutrients on the yield and quality of
289 agricultural products. Micronutrients with Macro-Effects. Publ. of Tarbiat Modarres Uni.
290 Tehran, Iran; 1999. (In Persian)
- 291 9. Malakoti MJ, Nafisi M. Necessity of production and consumption of agricultural sulfur for
292 increasing of agricultural products quantity and quality. Publ. Agri. edu. Agri. Ministry. Iran;
293 1996. (In Persian)
- 294 10. Ross M Welch, Robin D Graham. Breeding for micronutrients in staple food crops from a
295 human nutrition perspective. J. Experimental Botany. 2004;55 (396):353-364.
- 296 11. Sharma P, Swarup A. Comparison of Pyrite varying in water-soluble sulfur with Gypsum
297 for the reclamation of alkali soils under a Rice-Wheat rotation. *Biology and Fertility of soil J.*
298 1996;24:96-101.
- 299 12. Slaton NA, Norman RJ, Gilmour JT. Oxidation Rates of Commercial Elemental Sulfur
300 Products Applied to an Alkaline Silt Loam from Arkansas. *Soil Sci. Soci. America J.*
301 2001;65:239-243.
- 302 13. Saadati N. Determination of using necessity of S, Mg, Zn, Cu and B in Mazandaran
303 irrigated field. Publ. Country Rice Research Inst. Mazandaran. Amol, Iran; 2001. (In Persian)

- 304 14. Somani LL, Kanthaliya PC. Soil and Fertilizers at a glance. 1st ed. Agrotech Publishing
305 Academy. Hiran Magri, UDAIPUR; 2004.
- 306 15. Soleymanpanah M. The effect of Zinc and Mn Sulfate on quantitative and qualitative
307 characteristics of two variety of Wheat in different factors of phosphorous. Proc. of the 6th
308 Iranian Congr. of crop production and plant breeding of Mazandaran Uni. Babolsar, Iran;
309 2000. (In Persian).
- 310 16. Surajit K, Datta D. Principals and practices of rice production. The International Rice
311 Research Institute (IRRI). Los Banos, Philippines; 1981.
- 312 17. Turner F. Precision diagnosis on farms. Soil scientist at the Texas Agricultural
313 Experiment station. 2000. Accessed April 2012. Available:
314 www.ricejournal.com/backissues/april2000/story1.asp.
- 315 18. Wild A. Soils, Land and Food (Managing the land during the twenty-first century). 1st
316 Published.

UNDER PEER REVIEW