UNDER PEER REVIEW

Original Research Article

Maize response to competition from speargrass *(Imperata cylindrica* (L.) Raeuschel) regrowth from planted rhizomes I: Species Growth relationship, Yield and Yield components

---

ABSTRACT

Field study was conducted to evaluate the effect of speargrass shoot regrowth from previously planted rhizomes on the growth and yield of subsequent maize crop.

This study was evaluated in a Randomized Complete Block Design with three replications. The study was conducted at the International Institute of Tropical Agriculture Ibadan, Nigeria, between September 2005 and September 2006.

In this study maize and speargrass rhizomes were previously planted in eight monoculture densities (4, 8, 12, 16, 20, 32, 48 and 64 plants per plot and eight total densities in a mixture of 1:1 ratio of maize and speargrass (2:2-32:32) per plot in May 2005 and May 2006. .The speargrass regrowth experiment was initiated in September 2005 and 2006 respectively after the previous maize harvest. All plot that had maize or speargrass either as monoculture or in mixture with each other were slashed and subsequently replanted with maize maintaining the previous maize densities earlier planted; while all plot previously planted to speargrass either in monoculture or in mixture with maize were allow to regrow from the previously planted densities.

.

UNDER PEER REVIEW

Our Results of this study suggest that both maize and speargrass competed for the same resources. However, maize was more competitive than speargrass in 1:1 mixture in this study. Intraspecific competition between maize plants was responsible for maize grain yield loss of about 28.4%, while the overall interspecific competition effect from speargrass regrowth densities was responsible for a yield of about 18%. There was high and significant negative correlation between speargrass parameters and maize grain yield ($r \geq -0.56 \leq -0.78$) and maize biomass per plant ($r \geq -0.49 \leq -0.67$). For speargrass, interspecific competition was greater than intraspecific competition causing a speargrass biomass loss of about 35% from maize competition; while intraspecific competition accounted for about 17.9% biomass reduction. Speargrass densities of 8-16 plants m^{-2}, in mixture with maize had enormous rhizome biomass, and hence caused a grain yield of 43% due to vigorous regrowth. Farmers should try to use optimum maize population that will give between 5 and 6 plants m^{-2}; this will reduce the effect of speargrass regrowth from the rhizome, especially where land preparation is by slashing. Such maize population density will maximize the relative competitive ability of maize, reduce intraspecific competition and minimizes the effect of speargrass regrowth after land preparation, particularly in areas where farmers cannot afford to purchase herbicides and apply them correctly.

Key words: *Competition, maize, yield, speargrass regrowth, rhizomes, biomass*

1. INTRODUCTION

Maize is an important food staple for than 1 billion people in Sub-Saharan Africa, and a preferred crop of 900 million of world poor. In most developing countries it is

.

UNDER PEER REVIEW

grown in mixture with other crops or as sole crop [1]. Aggregated production especially in West Africa has shown an increase of about 73% between the 1980s and 2000s [2]. However, this increase is due to expansion in the area under cultivation while productivity per unit area still very low (0.5-1.0 ton/ha) due to several agro ecological factors including weeds infestation. Maize is sensitive to weed competition especially in the first 3 weeks after emergence [3]. Although maize is a vigorous and tall growing plant, it is susceptible to competition from weeds, with losses greater than 30% commonly reported [4]. Maize is a high- risk crop, mainly due to the varying climatic conditions as well as inadequate management practices. Researchers indicate that maize plants are very susceptible to weed competition and yield losses and are estimated at 35% to complete crop failure [5]. To obtain high crop yields weed control is very important because weed compete with maize for nutrients soil moisture and light. Therefore, information on competitive interaction between *Imperata cylindrica* (L.) Raueschel (speargrass) and maize will be useful for developing and implementing effective management programs. Speargrass *[Imperata cylindrica* (L.) Rauesch.] is a rhizomatous, perennial grass weed, widely distributed throughout the tropics and in some warm areas of the temperate region [6]. It has become a major problem in the production of arable crops such as maize, soybean, and root and tuber crops in forest transition zone of West Africa [7] [8]. Most of the methods of control (hoe weeding, hand pulling and slashing) employed by rural farmers are not effective, because of its ability to infest, spread and colonize native vegetation [9, 10, 11,12]. Yield loses attributed to speargrass infestation in maize has been reported to be between 40 and 80% [13, 14, 8, 15]. The density of speargrass and the competitive ability of the crop influence the effect of speargrass competition on crop yield. Most experiments conducted on speargrass have been to explore the relative aggressiveness of speargrass and certain crops [15] and few have been accomplished through the experimental manipulation of population crop density, proportion, or spatial arrangement [16]. However, none have critically looked at the interaction between speargrass shoot regrowth and maize densities and proportion with the aim of evaluating effects of densities and proportions of speargrass shoot regrowth on maize under the field environment or conditions.

.

UNDER PEER REVIEW

.It is therefore the objective of this experiment to assess the competitive ability of maize against speargrass regrowth from manipulated densities

2. MATERIAL AND METHODS

2.1Description of study site

The experiment was conducted at the research farm of International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria (7° 30'N, 3° 54'E), in the forest/savanna transition zone. The zone is characterized by an annual rainfall that averaged 1250 and 1500 with a bimodal distribution pattern which peaks in July and September. . Before monitoring the effect of speargrass regrowth on maize, the site was under speargrass fallow for four months The soil type at the experimental site was loamy sand (Alfisol) with a pH of 6.7 and organic matter of <2%, 0.13% N, 3.08 mg/kg P (available), 0.34K (cmol), and soil texture of 85% sand, 5% clay, and 9% silt. The average annual temperature is 26°C and dominant soil type is Alfisol [17, 8]

2.2 Experimental procedure and design

The speargrass for this study was initially planted on the 25th of May 2005, and May 30th, 2006 in both addition and replacement series experiments [18] from sprouted rhizomes using a grid of 2 m by 2m with 64 quadrilles of 25 cm by 25 cm; and its competitive effect monitored on maize growth. This previous treatment consisted of three replicates of 16 established densities as monoculture, which included eight monocultures of maize and speargrass at the following densities (4, 8, 12, 16, 20, 32, 48, 64) and eight mixtures of maize: speargrass (2:2, 4:4, 6:6, 8:8, 10:10, 16:16, 24:24, 32:32) per plot of 2 m ×2 m. All the plots were slashed immediately after harvesting previous maize (approximately 4 months after planting the maize), and maize stover and speargrass shoots packed out of the plots on September 20, 2005

.

UNDER PEER REVIEW

and September 25, 2006 resprctively.. The same grid of 2 m × 2 m with 64 quadrilles of 25 × 25 cm was used to replant maize (cv. 'ACR 89-DMR-ESR-W') in all the plots, which included plots previously planted to sole maize and speargrass; and plots that had maize and speargrass in mixture on 13 September 25, ,2005 and September 30, in 2006 . The regrowth experiment was also conducted with three replicates in a randomized complete block design. Each replicate included the same 16 treatments: eight monocultures of maize (4:0, 8:0, 12:0, 16:0, 20:0, 32:0, 48:0, 64:0 per plot) replanted in the same plot they were planted earlier (May 25, 2005 and May 30, 2006), while speargrass monoculture plots were allowed to regrow from the previously planted densities(0:4, 0:8, 0:12, 0:16, 0:20, 0:32, 0:48, 0:64 per plot) in September 25, 2005 and September 30, 2006 For the eight mixtures of maize: speargrass, maize was also replanted into each plot that had the maize: speargrass mixture of various densities and proportions to simulate the treatment arrangement of the experiment earlier in the season. (2:2, 4:4, 6:6, 8:8, 10:10, 16:16, 24:24, 32:32 per plot).. Maize seedlings were thinned to one stand per hill one week after planting. All plots were kept free of other weeds that may interfere with competition between the target species by weekly hand pulling of weeds. Basal fertilizer was applied at a recommended rate of 45 kg ha^{-1} of N, P_20_5, and K_20 at 2 weeks after planting (WAP) on October 9, 2005 and October 14, 2006, while urea at 45 kg N ha^{-1} was applied at 6 WAP, on 30 October 2005 and October 30 2006. Both types of fertilizers were applied broadcast.

2.3 Data Collection

In both years data were collected on height, leaf area and photosynthetic active radiation (PAR) of both species at 50% maize silking (7 WAP), on. 15 November 2005 and 20 November 2006~~.~~ Leaf area of each plant or species was determined ~~at~~

.

UNDER PEER REVIEW

50% silking from an average value of five plants measured per plot with LI-3000 portable leaf area meter with head scanner PAM 1684, (LI-COR, Inc., P. O. Box 4425, Lincoln, N668504 USA). PAR interception was determined with 1-m long Decagon sunfleck ceptometer (Decagon Devices, Inc., P O Box 835, Pullman, WA 99163 USA) at ground level and above the maize canopy on a cloudless day between 12:00 and 14:00 h. at 50% maize silking (7 WAP). An average of four reading per treatment from each replicate was used to obtain treatment mean.

The percentage of PAR intercepted (X) by the maize canopy was calculated as:

$$X = [1 - (B/A)] \times 100 \ldots\ldots\ldots\ldots [1]$$

Where, B is the PAR, umol m^{-2} s^{-1}, measured below the maize canopy 10 cm above the ground, A is the above maize canopy PAR reading, made in the open.

Similarly data was collected on, total above ground biomass of both species by cutting at the soil surface, and below ground biomass of speargrass (rhizomes) on 27 December 2005 and on 30 December 2006. An area of $2m^{-2}$ of was harvested from each plot for the determination of total above biomass of maize and speargrass, and grain yield. Maize grain yield was adjusted to 12% moisture content using the Tri-grain moisture tester (Model 14998 with Serial number 1170, Dickey-John Corporation Auburn, IL, 62615 USA). Both maize and speargrass plant samples were oven dried in a Gallenkamp oven (OVE–300 plus Series) at 80 ^{o}C until constant mass was recorded with a digital balance (XD–4K B042809, Denver Instrument Company, USA).

2.4 Data Analysis

All the data collected was analyzed by year, except for correlation between maize and speargrass parameters. ANOVA was performed using the MIXED MODEL and general linear model (GLM) procedures in the Statistical Analysis Systems software

.

UNDER PEER REVIEW

(SAS) [19] [20]. In the mixed model procedure, years and replication were considered random effects in the model. Data were analyzed and presented by year. Least-square means of the individual treatment effects were separated using the contrast at P = 0.05 and standard error of the means in the LSMEANS output. Spearman's correlation analysis was performed to determine the relationship between speargrass biomass, speargrass shoot density and maize biomass, maize density, and maize grain yield. In addition, combined multiple correlation and regression analysis were performed to determine the relationship between speargrass and maize attributes perceived to be competitive indicators/characters.

3. RESULTS

3.1 Maize and Speargrass biomass

In 2005, maize shoot biomass was higher in mixture with speargrass than in monoculture, but the differences were not significant (P>0.05) (Table 1). However, maize total plant biomass yield was significantly higher for 4 (P=0.0194) and 32 (P=0.0060) plants per plot in monoculture compared to mixture yield at the same density (Table 1). Contrast estimate in 2006 indicated that maize biomass was significantly higher in mixture than in monoculture ($P \leq 0.05$) at all densities except at 6:6 plant proportions of both maize and speargrass, compared to 12:0 of maize: speargrass, (P=0.0104). However, maize plant biomass was higher in monoculture at the following densities: 4 ($P \leq 0.05$), 16 (P>0.05), and 32 (P ≤ 0.05) per plot compared to their 1:1 mixed ratio in 2006 (Table1). All the speargrass parameters considered via density; shoot weight, rhizome weight and total dry weight as well as maize density were negatively correlated with maize biomass, respectively(r = -0.49, r = -0.45, r = -0.67, r = -0.64 and r = -0.58) (Table 2). In both years, except at

.

UNDER PEER REVIEW

speargrass density of 12 plants in 2006, when compared to 6:6 maize: speargrass plant proportions (*P*=0.052) per plot; speargrass shoot biomass was significantly higher in monoculture than in mixture (Table 1) in replacement series. . The relationship between speargrass parameters and maize density was significant in both years except for maize density and speargrass shoot biomass in 2006 (*P*=0.0967) (Table 3). Similarly, the relationship between speargrass parameters and maize height was significant in 2005(*P* =0.001), and significant for only speargrass rhizome biomass in 2006(*P* =0.009) (Table 3).

Table 1. Effects of speargrass regrowth on maize and speargrass biomass

		Biomass per plant			
		Maize		Speargrass	
Proportions					
Maize	**Speargrass**	**2005**	**2006**	**2005**	**2006**
-----No. per plot---		<----------(g plant^{-1})---------- ---------->			
4	0	700.86	359.42	-	-
2	2	787.00	435.33	285.56	154.64
0	4	-	-	398.70	324.24
8	0	663.33	241.64	-	-
4	4	640.83	325.50	151.74	123.60
0	8	-	-	249.46	229.32
12	0	538.53	253.97	-	-
6	6	607.56	245.28	98.55	318.17
0	12	-	-	270.44	163.28
16	0	431.11	249.26	-	-
8	8	775.83	276.92	110.73	143.29
0	16	-	-	190.98	133.89
20	0	455.88	214.85	-	-
10	10	558.60	227.45	82.67	86.96
0	20	-	-	187.72	149.35
32	0	381.93	165.68	-	-
16	16	521.12	188.04	42.00	35.88
0	32	-	-	120.49	85.19
48	0	289.23	104.17	-	-
24	24	453.82	162.40	27.00	19.34
0	48	-	-	101.72	64.87
64	0	190.40	89.88	-	-
32	32	313.54	119.43	16.33	21.29
0	64	-	-	82.44	52.07
SE±		77.4	28.6	36.96	46.78

.

UNDER PEER REVIEW

Contrast	<----------Probability < F1---------->			
4:0 (0:4) vs 4:4	0.0194	0.0046	<.0001	0.0567
8:0 (0:8) vs 8:8	NS	NS	0.0645	NS
16:0(0:16)vs 16:16	NS	NS	NS	NS
32:0(0:32) vs 32:32	0.0060	0.0080	<.0001	0.0085
4:0 (0:4) vs 2:2	0.0323	0.0118	0.0006	NS
8:0 (0:8) vs 4:4	0.0296	0.0008	0.0051	NS
12:0(0:12) vs 6:6	0.0193	0.0104	0.0028	0.0521
16:0(0:16) vs 8:8	NS	0.0019	0.0022	NS
20:0 (0:20) vs 10:10	0.0211	0.0042	NS	NS
32:0 (0:32) vs 16:16	NS	0.0484	<.0001	0.0040
48:0 (0:48) vs 24:24	NS	<.0001	NS	0.0003
64:0(0:64) vs 32:32	0.0060	0.0080	<.0001	0.0085

[1]NS denotes not significant at the 5 % level of probability;
– indicates where maize or speargrass is not applicable

Table 2. Relationship among maize and speargrass parameters at Ibadan, (means of 2005 and 2006 combined)

Speargrass/maize density	Maize biomass		Maize grain yield		Maize density	
	r	P value	r	P value	r	P value
Shoot biomass.	-0.45	0012	-0.56	<.0001	-0.74	<.0001
Shoot density	-0.49	0.0004	-0.73	<.0001	0.16	0.2817
Rhizome biomass	-0.67	<.0001	-0.78	<.0001	-0.89	<.0001
Total speargrass dwt (shoot + rhizome)	-0.64	<.0001	-0.72	<.0001	-0.86	<.0001
Maize density	-0.58	<.0001	-0.62	<.0001	1.00	-

.

UNDER PEER REVIEW

Table 3. Estimates of regression parameter for speargrass attributes as a linear function of maize density and maize height in a maize-speargrass competition experiment

	Maize							
	2005				2006			
	B_0[a]	B_1[b]	B_0	B_2[c]	B_0	B_1	B_0	B_2
Shoot biomass								
Estimates	109.418	-0.962	-400.03	2.667	127.494	-1.049	178.53	-0.728
Standard error	13.506	0.212	127.700	0.731	38.868	0.605	236.05	1.638
Pr > |t|	<.0001	0.0002	0.0046	0.0014	0.0034	0.0967	0.4575	0.6609
R^2	0.48		0.38		0.12		0.01	
Rhizome biomass								
Estimates	74.752	-0.674	-262.68	1.756	67.286	-0.56	-109.99	1.038
Standard error	9.559	0.150	92.506	0.534	7.338	0.114	51.828	0.359
Pr > |t|	<.0001	0.0002	0.0095	0.0033	<.0001	<.0001	0.0453	0.0086
R^2	0.48		0.33		0.52		0.27	
Total speargrass dwt.								
Estimates	184.171	-1.636	-662.72	4.423	194.78	-1.611	68.546	0.310
Standard error	22.753	0.358	216.907	1.251	36.588	0.569	244.310	1.695
Pr > |t|	<.0001	0.0001	0.0058	0.0019	<.0001	0.0097	0.7817	0.8568
R^2	0.49		0.36		0.27		0.002	

[a]B_0 is an intercept .

[b]B_1 is density.

[c]B_2 is height.

.

UNDER PEER REVIEW

3.2 Plant leaf area and height

Leaf area of maize per plant in monoculture was not significantly different from maize leaf area grown in mixture with speargrass in both years (P>0.05), except at the density of 8:0 maize plant per plot in compared to both species at 8:8 mixtures in both years (P=0.033). The effect of species competition on speargrass leaf area was significant only in 2005 (P=0.055). Speargrass leaf area per plant was consistently higher at all densities in monoculture than in mixture with maize in both years, except at 8:0 and 16:0 speargrass density per plot (Table 4). Averaged over the years, speargrass leaf area was significantly higher only at 12:0 plants density in monoculture compared to 6:6 proportion in mixture (P=0.004), and higher at 8:8 plant densities proportion in mixture compared to 16:0 plant density in monoculture (P=0.022). Maize height did not differ in monoculture and in mixture with speargrass in 2005. However, maize grown in mixture with speargrass at various species proportions were shorter except at densities of 2:2 and at 24:24 species proportion where maize was taller than in monoculture (P=0.051) in 2005 (Table 4). Maize and speargrass at 8:8 and 24:24 plant proportions were significantly taller than maize at 8:0(P=0.031) and 48:0 plant densities and proportions in 2006 (P=0.043) (Table 4). The proportions of maize and speargrass densities in mixture affected speargrass height significantly, compared to speargrass height in pure stand or monoculture in both years (P=0.05).

UNDER PEER REVIEW

TABLE4. Effects of speargrass shoot regrowth from previous density and proportion on maize and speargrass leaf area and height at 7WAP in 2005 and 2006.

		Leaf area per plant				Plant height			
Proportions		**Maize**		**Imperata**		**Maize**		**Imperata**	
Maize	**Imperata**	**2005**	**2006**	**2005**	**2006**	**2005**	**2006**	**2005**	**2006**
-----No. per plot---		----------(Cm^2 $plant^{-1}$)----------				----------(Cm $plant^{-1}$)----------			
4	0	2516.1	1639.99		-	171.0	142.4	-	-
2	2	2312.6	1909.86	178.13	256.78	191.7	165.8	47.2	79.9
0	4	-	-	285.52	307.35	-	-	60.3	84.0
8	0	2094.7	1623.89	-	-	174.3	149.3	-	-
4	4	2892.8	1679.85	174.77	289.10	180.3	151.3	50.5	74.9
0	8	-	-	222.99	285.67	-	-	49.4	83.4
12	0	2032.5	1338.87	-	-	178.8	133.8	-	-
6	6	2017.3	1448.72	207.44	263.70	176.9	136.6	59.5	74.2
0	12	-	-	242.27	294.85	-	-	67.7	83.7
16	0	2042.3	1703.50	-	-	170.7	148.3	-	-
8	8	2066.2	1615.22	254.60	286.83	163.9	151.8	60.0	79.8
0	16	-	-	234.29	282.18	-	-	67.9	87.4
20	0	2034.3	1520.68	-	-	173.3	146.5	-	-
10	10	2270.7	1369.40	236.44	190.56	163.9	126.1	50.1	68.9
0	20	-	-	376.97	277.51	-	-	70.2	89.3
32	0	2003.7	1495.48	-	-	170.9	139.8	-	-
16	16	2627.6	1705.44	192.74	263.78	169.7	137.9	59.5	73.3
0	32	--	-	362.19	278.43	-	-	74.5	84.5
48	0	1984.0	1269.15	-	-	168.3	133.7	-	-
24	24	1814.5	1433.26	181.91	211.62	172.5	142.8	58.6	82.7
0	48	-	-	265.42	237.38	-	-	63.3	82.1
64	0	1723.7	1225.56	-	-	164.3	132.9	-	-
32	32	1669.3	1326.07	148.65	238.94	164.0	134.1	60.2	82.6
0	64		-	355.21	242.56	-	-	73.3	81.5
SE±		304.03	166.27	52.08	35.90	7.3	6.2	5.0	3.9
		<--_-----_----------Probability < F[1]----------->-							
Contrast									
4:0 (0:4) vs 4:4		NS	NS	NS	NS	NS	NS	NS	0.044
8:0 (0:8) vs 8:8		0.0328	0.0434	NS	NS	NS	0.031	0.007	NS
16:0(0:16)vs 16:16		NS	NS	0.0071	NS	NS	NS	0.001	0.059
32:0(0:32) vs 32:32		NS	NS	NS	NS	NS	NS	NS	NS
4:0 (0:4) vs 2:2		NS	NS	NS	0.0052	NS	NS	0.018	0.005
8:0 (0:8) vs 4:4		NS	NS	NS	NS	NS	NS	NS	0.007
12:0(0:12) vs 6:6		NS	NS	0.0053	NS	NS	NS	NS	NS
16:0(0:16) vs 8:8		NS	NS	0.0081	NS	NS	NS	0.0006	NS
20:0 (0:20) vs 10:10		NS	NS	NS	NS	NS	NS	NS	NS
32:0 (0:32) vs 16:16		NS	NS	NS	NS	NS	NS	NS	0.073
48:0 (0:48) vs 24:24		NS	NS	0.0301	NS	0.051	0.043	0.059	NS
64:0(0:64) vs 32:32		NS	NS	NS	NS	NS	NS	NS	NS

[1]NS denotes not significant at the 5 % level of probability
– indicates where maize or speargrass is not applicable

.

UNDER PEER REVIEW

3.3 Maize grain yield (kg ha^{-1})

Maize grain yield was significantly higher in monoculture at all densities, compared to yields in mixture at all proportions in 2005 (*P*=0.004). Yield in both monoculture and mixture increased with increasing species densities. Grain yield differences between monoculture and mixture were higher (≥ 41%) with low densities (2:2–8:8) proportion of species in mixture, and lower (≥ 15–29%) with high densities (16:16–32:32) proportion of species in mixture (Table 5). When the interaction between maize and speargrass is averaged over densities in monoculture and species proportion in mixture, maize and speargrass association resulted in 36% maize yield loss with replacement series analysis. Yield was also higher in monoculture, compared to mixture for the densities considered in addition series (*P*=0.001), and yield differences between monoculture and mixture were also higher (≥7–18%) at low densities, and lower (4%) at high densities (Table 5). Irrespective of the method of analysis, speargrass competition with maize in 2005 resulted in a 22% maize yield reduction. The effect of speargrass competition with maize on maize grain yield in 2006 followed the same trend as in 2005. In monoculture, maize yield increased with increasing density up to the density of 32:0 (8 plants m^{-2}) and started dropping from the density of 48:0 (12 plants m^{-2}) to 64:0 (16 plants m^{-2}) which is an indication of interference, possible resulting from neighbouring maize plants (intraspecies completion)(Table 5). Similarly, grain yield in mixture with speargrass increased with increasing species proportions in mixture, but dropped at 10:10 and 32:32 plant proportions in mixture. Maize grain yield was significantly higher in monoculture at 4:0, 12:0, 16:0, 20:0, 32:0, 64:0 (*P*=0.05) and 8:0 (P>0.05) compared in mixture with half of each species in 2006 ; and yields were lower by 5% in monoculture for 48:0 plant density compared to 24:24 proportion of the plant

.

UNDER PEER REVIEW

species in mixture (*P*=0.013) in 2006 (Table 5). Maize yield were 6.49 and 26% higher in 4:4 and 8:8 of the plants species in mixture compared to 4:0 (*P*=0.005) and 8:0 (*P*<0.001) plant densities in monoculture in 2006 (Table 5). However, yields at 16:0 and 32:0 plants in 2006 in monoculture, were significantly higher by 24% and 32% than yields in 16:16 and 32:32 of the plant species in mixture (*P*=0.05). The overall competition effect indicates a 16% grain yield reduction in maize resulting from speargrass interaction with maize in 2006.

Table 5. Effects of speargrass regrowth densities and proportions on maize grain yield at Ibadan in 2005 and 2006

Mixture proportion		Maize grain yield	
Maize	**Speargrass**	**2005**	**2006**
-----No. per plot---		**<----(Kg ha^{-1})----->**	
4	0	1499.61	944.76
2	2	882.78	654.60
8	0	2768.15	1425.76
4	4	1226.96	1010.35
12	0	3302.38	2479.37
6	6	1917.66	1228.48
16	0	3588.60	3002.78
8	8	2547.80	1926.63
20	0	4792.67	3170.40
10	10	2939.36	1614.49
32	0	5651.27	3446.82
16	16	3467.39	2295.99
48	0	5795.89	2632.43
24	24	4174.79	2755.67
64	0	6405.92	2448.74
32	32	5441.79	2333.29
SE±		420.56	254.17
Contrast		<---------Probability > F---------->	
4:0 (0:4) vs 4:4		< 0.0001	0.0049
8:0 (0:8) vs 8:8		0.0004	0.0001
16:0(0:16) vs 16:16		0.0060	0.0034
32:0(0:32) vs 32:32		0.0041	0.0193
4:0 (0:4) vs 2:2		< 0.0001	0.0403
8:0 (0:8) vs 4:4		< 0.0001	NS

.

UNDER PEER REVIEW

12:0 (0:12) vs 6:6	0.0004	< 0.0001
16:0(0:16) vs 8:8	0.0002	0.0006
20:0 (0:20) vs 10:10	0.0044	0.0396
32:0 (0:32) vs 16:16	0.0002	< 0.0001
48:0 (0:48) vs 24:24	0.0011	0.0286
64:0(0:64) vs 32:32	0.0041	0.0193

3.4 Light interception

Light interception increased with increasing species density and proportion in the mixture in both years (*P*=0.001, in 2005 and *P*=0.064, in 2006). Densities of 2-8 plants of each species in mixture intercepted ≤ 40% (13-40%) of the incident PAR in 2005, while the densities of 10-32 plants of each species in the mixture intercepted ≥ 46% (46-77%) of the incident PAR received by the canopy in 2005(Fig.1). However, in 2006, the PAR interception was more variable among the proportions compared to that of 2005. Maize-speargrass mixture at 2-8 plants of each species per plot intercepted between 35 and 53%, while for the densities at 10-32 plants of each species in mixture per plot, incident PAR interception by the canopy was between 37 and 76% in 2006 (Fig. 1). Averaged over the years, percentage PAR interception was significantly (*P*=0.05) different between the various densities in mixture.

.

UNDER PEER REVIEW



Figure 1: Light interception by maize and speargrass proportions at various densities in 2005 and 2006

UNDER PEER REVIEW

4. DISCUSSION

The effect of competition from speargrass shoot regrowth competition affected the growth and development of both species in pure or mixed cultures. However, maize also responded to density. The relationship between maize density and PAR in both years indicates that the slopes of the lines were positive. This means that as maize density increases, so does its ability to intercept PAR, and this in turn affects speargrass, being a shade-intolerant plant [8, 21]. Hence the negative slope observed in the relationship between maize density and speargrass biomass in this study. The result means that increasing maize density further will result in an average increase of PAR interception by 0.35 times , and this will translate into an average loss of speargrass shoot biomass by 1.01 times, rhizomes by 0.62 times, and total biomass by 1.62 times. Thus, the competitive ability of speargrass was enhanced at a lower density of maize; this is evident in the higher speargrass biomass (≥100 g per plant) at lower maize densities in mixture (2:2-8:8) compared to speargrass biomass at higher maize density (10:10-32:32) interaction in mixture (≤87 g per plant). The observed antagonistic effect between maize and speargrass is attributable to severe competition for light, which is evident by the intensity of competition at high maize densities. All speargrass attributes measured were negatively related to maize density due to competition for light. These results are in agreement with similar studies that report on nutsedge competition with maize and tomato [22, 23]. Similar competition for light resources between spring barley and weeds has been reported [24]. Since maize and speargrass have similar growth habits and life cycles, the two species may have been competing for the same resources and the success of speargrass in this case may be dependent on the

.

UNDER PEER REVIEW

extent of rhizome regrowth and interference. Although speargrass was established from presprouted rhizomes, maize grew taller and developed a closed canopy faster than speargrass. This may have induced shade over speargrass, and reduce PAR available to speargrass. Therefore, maize limited the light received by speargrass . Previous study reported that little in height differences of two plants in time of competition can cause a lot differences in competition [25]. Change in height difference between crop and weeds has been considered as one of the most important effects of weed competition that can be favorable for crop or weeds based on species and purposed conditions [26]. Maize height irrespective of culture or proportion with speargrass averaged 172cm and 142 cm at 7 WAP in 2005 and 2006 respectively; while speargrass height averaged 58.5 cm and 80.5 cm at 7 WAP in 2005 and 2006 respectively. Plant leaf area at 7 WAP averaged 2131.4 cm and 1519.1cm in 2005 and 2006 respectively for maize whether or not in mixture. Similarly speargrass leaf area at 7 WAP averaged 245.3 cm and 262.9 cm in 2005 and 2006 respectively. Plant height and leaf area have been reported as important attributes of plants that defines growth and competitive ability in mixtures [27, 28, 29, 30, 31]. The observed differences in the competitive ability of maize and speargrass from previous reports may be attributed to differential densities and proportion of the species in competition, as well as canopy relationship of the species. This is moreso as the density of maize may have been fixed across all level of speargrass infestation, and also under varied management strategies. Though maize was more competitive than speargrass in 1:1 mixture in this study; the fact that speargrass shoot density was negatively correlated with maize grain yield per plant ($r=-0.73$, $P<.0001$) and maize biomass per plant ($r=-0.49$, $P=0.0004$) is also an

.

UNDER PEER REVIEW

indication of speargrass competition with maize, and this can be more intense depending on the density and proportion in competition, and duration of infestation. This result conforms to earlier studies that have reported on the competitive relationships between maize and speargrass [8, 15].

5. CONCLUSION

The competitive interaction between maize and speargrass indicate that both compete for the same resources. Their competitive interaction and ability depends on the proximity, the regrowth stage, and the canopy relationship during growth association. Maize due to taller canopies had advantage for light competition over speargrass, and was more competitive than speargrass at low densities of speargrass. The greater competitiveness of speargrass, as we observed, may be apparent only under conditions of relatively high speargrass infestation and long duration of competition. Speargrass can compete strongly with maize for light especially at high densities, and competition for light must therefore be managed to its disadvantage. Speargrass competition can be intensified if disturbance due to slashing encouraged shoot regrowth from rhizomes, because rhizome competition was found to have a greater effect on the relative performance of speargrass. Thus, speargrass densities of 8-16 plant m^{-2} in mixture with maize had the greatest rhizome biomass and hence caused a grain yield reduction of up to 43%, due to vigorous regrowth. Slashing, often employed by farmers probably encourages higher speargrass rhizome activity, resulting in a more intense competition with associated crops. Speargrass is quite expensive to manage if effective control is required, farmers should try to use or integrate a maize seeding rate that will give

.

UNDER PEER REVIEW

the optimum population that will reduce the effect of speargrass shoot regrowth from rhizome after land preparation. Such maize population density will maximize the relative competitive ability of maize and minimizes the effect of speargrass, particularly in areas where farmers cannot afford to purchase herbicides and apply them correctly.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Badu-Apraku B, Fakorede M. A.B., and Ajala S.O. (2003) Promoting maize technology transfer in West and Central Africa: a networking approach. In: Maize revolution in West and Central Africa. Proceedings of a Regional Maize Workshop (Eds B Badu- Apraku, MAB Fakorede, M Ouedraogo, RJ Carsky& A Menkir), 14–18 May, 2001, IITA–Cotonou, Benin Republic. 45–65., WECAMAN/IITA, Ibadan, Nigeria.

2. Fakorede MAB, Badu-Apraku B, Kamara AY, Menkir A and Ajala SO (2003) Maize revolution in West and Central Africa: an overview. In: Maize revolution in West and Central Africa. Proceedings of a Regional Maize Workshop (eds B Badu-Apraku, B, MAB Fakorede, M Ouedraogo, RJ Carsky & A Menkir), 14–18 May, 2001, IITA– Cotonou, Benin Republic. 3–15. WECAMAN/IITA, Ibadan, Nigeria.

3. Olorunmaiye PM, and Olorunmaiye KS (2009). Effect of integrated weed management on weed control and yield components of maize and cassava intercrop in a southern Guinea savanna ecology of Nigeria. Aus J Crop Sci. 3: 129-136.

4. Rahman, A. 1985. Weed control in maize in New Zealand. Pp 37-45 In: Maize-Management to Market. (Eds.): H.A. Eagles and G.S. Wratt. Agron. Soc. N.Z., Special pub. No. 4, Palmerston North, New Zealand.

5. Sharma, V., D.R. Thakur and J.J. Sharma. 1998. Effect of metachlor and its combination with atrazine on weed control in maize (Zea mays). Indian J. Argon. 43: 677–80.

.

UNDER PEER REVIEW

6. Holm, L.G., Plucknett, D.L., Pancho, J.N. and Hherberger, J.P. 1977. Imperata cylindrica (L.) Beauv. In The world's worst weeds , pp 62–71.University Press of Hawaii, Honolulu. Hawaii.

7. Chikoye D , Ekeleme F and Ambe JT (1999) Survey of distribution and farmers' perceptions of speargrass [Imperatacylindrica (L.) Raeuschel] in cassava–based systems in West Africa. International Journal of Pest Management 45, 305–311.

8. Udensi, U. E., Akobundu, I. O., Ayeni, A. O. and Chikoye, D. (1999) Management of speargrass (Imperata cylindrica (L.) Raeuschel) with velvet bean (Mucuna pruriens var. utilis) and herbicides .Weed Technology 13: 201–208.

9. Dozier H, Gaffney JF, McDonald SK, Johnson RL and Shilling DG (1998) Cogongrass in the United States: history, ecology, impacts and management. Weed Technology 12, 737–743.

10. Akobundu, I. O. and Fagade S. O. 1978. Weed problems of African rice lands. In: Rice in Africa, pp 181-192, Buddenhagen, I W. and Persley, G. F.(eds) Academic Press, London..

11. Ogunyemi, S. (1977).The distribution of some weeds of arable crops in Southwestern Nigeria. In: Proceedings Seventh Conference of the Weed Science Society of Nigerian July 1977,pp 29–34, NRCRI, Umudike, Nigeria

12. Eussen, J. H. H., Slamet, S. and Soeroto, D (1976). Competition between alang-alang [Imperata cylindrica (L.) Beauv.] and some crop plants. Biotrop Bulletin 10: 24pp.

13. Lagoke, S.T.O, and A. A. Fayemi {1981}. Competition of speargras {Imperata cylindrica var. Africana {L.} Beauv} with maize. Indian Journal of Weed Science. 13 {2}: 83-94.

14. Koch W, Grobmann F, Weber A, Lutzeyer HJ and Akobundu, IO (1990) Weeds as components of maize/cassava cropping systems. In: Standortgemaesselandwirtschaft in West Africa, 283–298. UniversitaetHohenheim. Stuttgart, Germany.

15. Chikoye D, Ekeleme F and Udensi UE (2001) Imperata cylindrica suppression by intercropping cover crops in Zea mays /Manihot esculenta systems. Weed Science 49; 658−667

.

UNDER PEER REVIEW

16. Udensi Ekea Udensi and David Chikoye (2013). Effect of relative proportion and density on competition between Speargrass (*Imperata cylindrical* (L.) Raeuschel) and Maize (Z*ea mays* L.) in a moist savannah of South-western Nigeria

17. Okigbo, B. N. (1995). Major farming systems of the lowland savanna of sub-Saharan Africa and the potential for improvement. In: *Moist savannas of Africa*: potentials and constrains for crop production. B.T. Kang, I.O. Akobundu, V.M. Manyong, R.J. Carsky, N. Sanginga and E.A. Kueneman, eds. Proceedings of the International Workshop held at Cotonou, Republic of Benin, 19-23 September 1994, pp 31-48. Produced by IITA, Ibadan, Nigeria.

18. Radosevich SR (1987) Method to study interactions among crops and weeds. Weed Technology 1, 190–198.

19. SAS Institute Inc. 1995. SAS/STAT *User's Guide, Version 6 Fouth Edition. Vol. 2.* SAS Institute Inc.Carey, NC, USA.

20. Littell, R.C., Milliken, G.A., Stroup, W.W, and Wolfinger, R.D. (1996). SAS system for mixed models. SAS Institute Inc., Cary, NC, USA. 656 pp.

21. Chikoye, David, Ellis-Jones, Jim, Kormawa, Patrick, Udensi, Udensi E., Ibana, Simon E., and Avav,Ter-rum. 2006. Options for Cogongrass (Imperata cylindrica) Control in White Guinea Yam (Dioscorea rotundata and Cassava (Manihot esculenta). Weed Technology 20: 784-792.

22. Santos, B.M., T.A. Bewick and D. J. Shilling (1997b). Competitive interaction of tomato (*Lycopersicon esculentum*) and nutsedge (*Cyperus spp*.). *Weed Science* 45:229-233.

23. Tuor, F. A. and Froud-Williams, R. J. (2002). Interaction between purple nutsedge, maize and soybean. *International Journal of Pest Management* 48: 65-71

24. Petersen, J. (2003).Weed:Spring barley competition for applied nitrogen in pig slurry. Weed Research 43, 33-39.

25. Kropff MJ, Weaver SE, Smits MA (1992) Use of ecophysiological models for crop- weed interference: relations among weed density, relative time of weed emergence, relative leaf area and yield loss. Weed Sci. 40: 296-301.

26. Knezevic SZ, Weise SF, Swanton CJ (1994) Interference of redroot pigweed (Amaranthus retroflexus L.) in maize (Zea mays L.). Weed Sci. 42: 568-578.

UNDER PEER REVIEW

27. Baumann DT, Bastiaans, L and Kropff, MJ (2002) Intercropping system optimization for yield, quality, and weed suppression combing mechanistic and descriptive models. Agronomy Journal 94, 734–742.

28. Daugovish O, Thill, DC and Shafii B (2002) Competition between wild oat (*Avena fatua*) and yellow mustard (Sinapis alba) or canola (Brassica rapus). Weed Science 50, 344 587–594.

29. Lemerle D., Verbeck B., Cousen RD and Coombes NE (1996) The potential for selecting wheat varieties strongly competitive against weeds. Weed Research 36 , 505–513.

30. Roush ML and Radosevich SR (1985) Relationship between growth and competitiveness of four annual weeds. Journal of Applied Ecology 22 , 895–899.

31. Wicks GA, Ramsel RE, Nordquist, PT and Schmidt JW (1986) Impact of wheat cultivars on establishment and suppression of summer annual weeds. Agronomy Journal 78 , 59−62.

.