

Original Research Article

Effect of radiation process followed by traditional treatments on nutritional quality of high tannin sorghum cultivar

Ismat G. Abdalla¹, Khogali E. Ahmed¹ and Elfadil E. Babiker^{2*}

¹Department of Food Science and Technology, Faculty of Agriculture, University of
Khartoum, Khartoum North 13314, Shambat, Sudan

²Department of Food Science and Nutrition, College of Food and Agricultural Sciences,
King Saud University, P. O. Box 2460, Riyadh 11451, Kingdom of Saudi Arabia

Author's contribution

This work was carried out (Planning, experiments, analysis and paper writing) in
collaboration between all authors

ABSTRACT

Aims: To study the effect of radiation process on nutritional quality of raw and processed flour of high tannin sorghum cultivar (Karamaka).

Place and Duration of Study: Department of Food Science and Technology, Faculty of Agriculture, University of Khartoum, Sudan between December 2012 and March 2013

Methodology: The flour of the cultivar was radiated using gamma radiation at doses of 5, 10 and 15 kGy and thereafter the radiated flour was fermented and/or cooked. Antinutrients, protein digestibility, amino acid contents and score, total and extractable minerals were determined.

Results: Radiation of raw flour decreased the level of antinutrients with a concomitant increase in protein digestibility and minerals extractability. Moreover, radiation increased the level of some amino acids and decreased others. Among amino acids the most limiting ones are lysine, methionine plus cysteine and threonine. Fermentation and/or cooking of radiated and non-radiated flour improved the protein digestibility and minerals extractability as well as the level of some amino acids. Most of the amino acids were slightly stable against all treatments with few exceptions.

Conclusion: Radiation of the flour followed by traditional processing greatly improved the nutritional quality of sorghum flour.

Key words: Sorghum, tannin, radiation, traditional treatments, nutrients

*Correspondent: E-mail: ebabiker.c@ksu.edu.sa, Tel. +966565197408 Fax: +96614678394

1. INTRODUCTION

Plant foods such as cereals and legumes have consistently been listed as the major potential sources of dietary protein. However, with increasing interest in new food sources and in improved nutritional quality, the seeds of cereals are now receiving more attention. Most of the African people like to consume sorghum as fermented flat bread (Kisra), thick porridge (Aceda), thin fermented gruel (Nasha), boiled grains (Balella) and beverages (Abreh and Hulumur) as a source of protein and minerals [1]. Irradiation could replace or supplement chemical preservatives; in other cases it may have unique advantages whether dry or frozen foods [2].

In agricultural science and food technology, recent research has elucidated new potential applications for radiation. High doses of ionising radiation have been shown to inhibit growth of microbes [3, 4] and to reduce the level of antinutrients [5] which were reported to reduce the availability of proteins and minerals [6, 7, 8, 9, 10]. There are also many reports supporting the use of gamma irradiation as a fungicidal agent [11].

There are insufficient reports about possible effects of gamma radiation on nutritional value of the seeds associated with seed sterilizations. Seeds of different plants that are consumed as food have varying nutrient values which are dependent on the basic constituents of seed proteins. The amino group ($-NH_2$) is the most radiosensitive portion of the amino acids [12]. Extensive research showed that the macronutrients (carbohydrates, proteins and lipids) content are relatively stable against irradiation dose up to 10 kGy [13]. However, Lee *et al.* [14] reported that gamma irradiation affects proteins by causing conformational changes, oxidation of amino acids, rupturing of covalent bonds and formation of protein free radicals. Also, chemical changes in the proteins caused by gamma irradiation include fragmentation, cross-linking, aggregation and oxidation by oxygen radicals that are generated in the radiolysis of water.

Gamma irradiation has a slight effect on the amino acid profile at recommended doses to foods [13]. This effect could be related to the structure of each amino acid, simple amino acids increased upon irradiation, such as glycine, which undergo reductive deamination and decarboxylation [15]. In addition, aliphatic amino acids with increasing chain length provide additional C–H bonds for interaction with OH radicals which reduces the extent of oxidative deamination [15]. Wang and vonSonntage [16] reported that sulphur-containing as well as aromatic amino acids are, in general, the most sensitive to irradiation, while simple amino acids could be formed by destruction of other amino acids. Mohamed *et al.* [5] reported that millet flour had a severe problem during storage and it was observed to produce off-flavour and bitter taste which can be overcome by using radiation.

The literature has many reports demonstrating that thermal processing methods improve the nutritional quality of foods. However, there is a scarcity of information relating to the effects of processing with ionizing energy. Therefore, the present work was carried out to investigate the effect of radiation process followed by traditional treatments on nutritional quality of raw and processed high tannin sorghum cultivar.

2. MATERIALS AND METHODS

2.1 Materials

Grains of the sorghum cultivar (Karamaka) were collected from Department of Agronomy, Faculty of Agriculture, University of Khartoum, Sudan. Collected seeds (4 kg) of the cultivar were ground to pass a 0.4 mm screen. All chemicals used for the experiments were of reagent grade.

2.2 Radiation process

The flour with a moisture content of 5.45% was spread uniformly and stored in polythene bags of mass of 100 gm, gamma radiation process was conducted at Kaila irradiation processing unit, Sudanese Atomic Energy Corporation (SAEC). The flour was exposed to gamma rays generated by a cobalt-60 source (Gammacell 220, MDS Nordion, Ottawa, Canada). The flour was radiated at 0, 5, 10 and 15 kGy following the procedures described by Helinski *et al.* [17] with a dose rate of ca. 3.2 kGy/h at 24±1 °C and normal relative humidity. Double side irradiation (exposure to both sides) was performed for uniform dose delivery. A dosimetry system was used to measure the dose received by the

batch based on the Gafchromic HD-810 film (International Specialty Products, NJ, USA; FAO/IAEA/USDA 2003). Three dosimeters were included with each batch of flour and read after irradiation with a Radiachromics reader (Far West Technology Inc., CA, USA). All experiments were repeated 3 times and 3 replicates of each flour type were irradiated. Radiated and non-radiated flour of the cultivar was naturally fermented till the pH of the dough reached 4.50 and thereafter cooked for 20 min in a water bath and then dried and ground to pass a 0.4 mm screen for further analysis.

2.3 Tannin determination

Quantitative determination of tannins was carried out using the modified vanillin-HCl method according to Price and Butler [18] using 200 mg sample. A standard curve was prepared expressing the results as catechin equivalents, i.e amount of catechin (mg per ml) which gives a colour intensity equivalent to that given by tannins after correcting for blank.

2.4 Phytate determination

Phytic acid content was determined by the method described by Wheeler and Ferrel [19] using 2.0 gm dried sample. A standard curve was prepared expressing the results as $\text{Fe}(\text{NO}_3)_3$ equivalent. Phytate phosphorus was calculated from the standard curve assuming a 4:6 iron to phosphorus molar ratio.

2.5 *In vitro* protein digestibility (IVPD) determination

The IVPD of the samples was carried out using single enzyme (pepsin) digestion according to the method of Maliwal [20] in the manner described by Monjula and John [21]. Digestibility was calculated using the following equation:

$$\text{Protein digestibility (\%)} = \frac{\text{N in supernatant} - \text{N in pepsin}}{\text{N in sample}} \times 100$$

2.6 Total and extractable minerals determination

For total content, minerals were extracted from the samples by the dry ashing method described by Chapman and Pratt [22]. Minerals were determined using atomic absorption spectrophotometer (Perkin-Elmer 2380). Na and K contents were determined using flame photometer (corning 400). For extractability, minerals in the samples were extracted by

the method described by Chauhan and Mahjan [23] using 1.0 gm of the sample. HCl extractability (%) was determined as follows:

$$\text{Mineral extractability (\%)} = \frac{\text{Mineral extractable in 0:03NHCl (mg/100g)}}{\text{Total minerals (mg/100g)}} \times 100$$

2.7 Amino acid content and score determination

The amino acids composition of the samples was measured on hydrolysates using amino acids analyzer (Sykam-S7130, Tokyo, Japan) based on high performance liquid chromatography technique. Sample hydrolysates were prepared following the method of Moore and Stain [24]. About 200 mg of the sample was taken in a hydrolysis tube. Then 5 ml of 6 N HCl was added to the sample and the tube tightly closed and incubated at 110 °C for 24 h. After incubation, the solution was filtered and 200 ml of the filtrate was evaporated to dryness at 140 °C for 1 h. The hydrolysates after dryness were diluted with 1.0 ml of 0.12 N citrate buffer (pH 2.2). Aliquot of 150 µl of the sample hydrolysates was injected in an action separation column at 130 °C. Ninhydrin solution and an eluent buffer (solvent A, pH 3.45 and solvent B, pH 10.85) were delivered simultaneously into a high temperature reactor coil (16 m length) at a flow rate of 0.7 ml/min. The buffer/ninhydrin mixture was heated in the reactor at 130 °C for 2 min to accelerate chemical reaction of amino acids with ninhydrin. The products of the reaction mixture were detected at wavelengths of 570 and 440 nm on a dual channel photometer. The amino acids composition was calculated from the areas of standards obtained from the integrator and expressed as gm/100 gm protein.

The essential amino acid (EAA) score was determined by applying the formula:

$$\text{AAS score \%} = \frac{\text{gm of EAA in 100 gm test protein}}{\text{gm of EAA in 100 gm FAO/WHO reference pattern}} \times 100$$

2.8 Statistical analysis

Each determination was carried out on three separate samples and analysed in duplicate on dry weight basis; the figures were then averaged. Data were assessed by the analysis of variance [25]. Comparisons of means for treatments were made using Duncan's multiple range tests. Significance was accepted at $P \geq 0.05$.

3. RESULTS AND DISCUSSION

The effect of gamma irradiation and/or traditional processings on tannin and phytate contents and *in vitro* protein digestibility of sorghum cultivar (Karamaka) are shown in Table 1. Ten cultivars of sorghum were investigated for tannin content; Karamaka was found to contain higher amount of tannin (2.90%) and accordingly designated as a high tannin cultivar. Tannin content of raw flour significantly ($P \leq 0.05$) reduced after radiation and the level of reduction increased with increase in radiation dose as reported by Mohamed *et al.* [5] for millet cultivars. Traditional processing (fermentation and/or cooking) of raw flour significantly ($P \leq 0.05$) decreased the level of tannin which agree with that reported by Idris *et al.* [26], who stated that combination of fermentation and cooking of sorghum improved the nutritional quality and drastically reduced tannin content to safe levels than any other processing methods. Also Osman [27] reported that fermentation markedly reduced tannin content of sorghum cultivars. The reduction in tannin content due to fermentation is likely to be due to biochemical activity of fermenting organisms [28]. However, when traditional treatments were applied in combination with radiation process further reduction in tannin content was observed with a minimum value of 0.465% obtained for fermented and cooked seeds that radiated at a dose of 10 KGy. The results are in agreement with those reported by Brigide and Canniatti-Brazaca [29] who observed that tannin content was inversely correlated with the applied irradiation doses.

Radiation of raw flour significantly ($P \leq 0.05$) reduced the level of phytate and the rate of reduction increased with increase in radiation dose as reported by Mohamed *et al.* [5] for millet cultivars. Traditional processing (fermentation and/or cooking) significantly ($P \leq 0.05$) decreased the level of phytate as reported by Idris *et al.* [26], who found that combination of cooking and fermentation of sorghum significantly ($P \leq 0.05$) reduced the level of phytate. Also Osman [27] reported that fermentation markedly and significantly ($P \leq 0.05$) reduced phytate content of sorghum cultivars. The reduction in phytate content due to fermentation is likely due to biochemical activity of the microbial enzyme phytase which hydrolyzed phytate. However, when traditional treatments applied in combination with radiation process further reduction was observed with a minimum value of 105.40 mg/100gm obtained for fermented flour radiated at 15 KGy. Osman *et*

al. [30] reported that phytic acid and tannin contents of broad bean were significantly ($P \leq 0.05$) reduced as a result of radiation. El- Niely [31] stated that radiation after processing significantly ($P \leq 0.05$) decreased the level of phytic acid of legumes. The reduction in phytic acid during radiation process is likely to be due to chemical degradation of phytate to the lower inositol phosphates and inositol by the action of free radicals produced by radiation or might be due to cleavage of the phytate ring itself [12]. Both cereal and microbial phytases can contribute to a reduction in phytate during fermentation process [28]. Duodu *et al.* [32] reported that cooking did not decrease phytic acid in sorghum porridge, but cooking and irradiation caused a significant decrease (40%).

The effect of gamma irradiation on *in vitro* protein digestibility (IVPD) of Karamaka cultivar is shown in Table 1. Radiation of raw flour was found to be effective in improving the IVPD at low dose of 5 KGy but as the level of radiation increased the IVPD significantly ($P \leq 0.05$) decreased. Traditional processing (fermentation and/or cooking) of raw flour significantly ($P \leq 0.05$) increased the level of IVPD as reported by Abdelhaleem *et al.* [33]. When traditional treatments applied in combination with radiation process further improvement in IVPD was observed with a maximum value of 42.71% for fermented flour radiated at 15 KGy. The results obtained agrees with results obtained by El-Niely [31] who studied the influence of irradiation on *in vitro* protein digestibility of broad beans irradiated at 2.5, 5, 10 and 20 KGy, he observed that the *in vitro* protein digestibility improved by 4.5, 10, 16 and 20%, respectively. The improvement in IVPD of fermented flour before and after radiation is likely to be due to reduction in antinutrients especially phytate which degraded by phytase during fermentation as reported by Sokrab *et al.* [10]. On the other hand the reduction in IVPD may be due to disulphide formation resulting in disulphide cross-linked protein oligomers and polymers. These polymers form a coat reducing the accessibility of the protein bodies to enzymatic attack [32].

Table 2 shows the content and HCl extractability of some major minerals (Na, K and Mg) of high tannin sorghum cultivar (Karamaka) as affected by radiation and/or traditional processings. For the raw flour, Na content was found to be 22.60 mg/100g and out of this amount about 67.4% was found to be extractable; K was 470.10 mg/100g with

extractability of 50.00% and Mg was 50.10 mg/100g and out of this amount about 52.50% was extractable. Variations in extractability between Na, K and Mg is likely to be due to the binding ability of such minerals with antinutrients which are observed to form complexes with minerals and reduced their extractability as reported by Idris *et al.* [26]. Radiation of raw flour had no significant ($P \leq 0.05$) effect on total Na, K and Mg but significantly ($P \leq 0.05$) increased the extractability of such minerals. The increment in minerals extractability after radiation could be attributed to reduction in the antinutrients. Fermentation of raw flour significantly ($P \leq 0.05$) increased both total and extractable Na, K and Mg. Fermentation leads to lowering of pH as a consequence of bacterial production of lactic and acetic acids, which is favourable for phytase activities, resulting in lowering of phytate. The reduction in phytate resulted in an improvement in minerals extractability.

Cooking of raw flour decreased both total and extractable minerals. Cooking of raw fermented dough alleviates the effect of cooking and significantly ($P \leq 0.05$) increased minerals extractability. Fermentation of radiated flour significantly ($P \leq 0.05$) increased both total and extractable minerals. Radiation of the flour at 5, 10 or 15 KGy followed by cooking decreased minerals content but increased their extractability as compared to raw samples. Cooking of irradiated fermented dough caused a significant ($P \leq 0.05$) improvement in extractable minerals. Moreover, as the level of radiation increased the degree of improvement in minerals extractability was significant ($P \leq 0.05$). The results obtained showed that processing of radiated flour significantly ($P \leq 0.05$) increased the extractable Na, K and Mg with increase in radiation dose but had no significant ($P \leq 0.05$) effect on total minerals. Generally the increase in extractable minerals during traditional processing before and after radiation could be attributed to the reduction in the level of antinutrients as a result of such treatments [6, 9].

Table 3 shows the content and HCl extractability of some trace minerals (Zn, Cu, Mn and Co) of sorghum cultivar (Karamaka) as affected by radiation and/or traditional processings. The raw flour of the cultivar contained 2.54, 0.42, 3.35 and 0.15 mg/100g of Zn, Cu, Mn and Co, respectively and out of this amount about 49.29, 34.17, 39.42 and 62.07 were found to be extractable for the minerals, respectively. Variations in extractability between trace minerals are likely to be due to the binding ability of such

minerals with antinutrients as reported by Idris *et al.* [26]. Radiation of raw flour had no significant ($P \leq 0.05$) effect on total minerals but significantly ($P \leq 0.05$) increased the HCl extractability of such minerals. Fermentation of raw flour slightly increased total minerals and significantly ($P \leq 0.05$) increased the extractable ones. Cooking of raw flour had no significant ($P \leq 0.05$) effect on total and extractable minerals. Cooking of fermented dough alleviates the effect of cooking and significantly ($P \leq 0.05$) increased the extractability of trace minerals but did not significantly ($P \leq 0.05$) affect total minerals.

Radiation at all levels followed by fermentation of raw flour gave varying changes in total and extractable minerals for the cultivar but generally there is a significant ($P \leq 0.05$) increase in total as well as in extractable minerals compared to raw flour. However, cooking of radiated flour had no great effect on content but significantly ($P \leq 0.05$) increased extractability of trace minerals. Cooking of radiated/fermented dough caused a significant ($P \leq 0.05$) improvement in extractable minerals but did not affect total minerals. The results obtained showed that as the level of radiation increased the degree of improvement in minerals extractability was significant ($P \leq 0.05$). The results showed that processing of irradiated flour significantly ($P \leq 0.05$) increased the extractable trace minerals with increase in radiation dose but had no significant ($P \leq 0.05$) effect on total trace minerals. The increment in extractable minerals during traditional processing before and after radiation could be attributed to the reduction in the level of antinutrients as a result of such treatments.

Tables 4 shows the effect of radiation process on amino acids composition of raw and treated flour of Karamaka cultivar. Radiation of raw flour of the cultivar at 5, 10 and 15 kGy decreased the level of all amino acids except phenylalanine plus tyrosine with a concomitant decrease in amino acid score except that of phenylalanine + tyrosine. Joseph *et al.* [34] reported that with the exception of tyrosine the amino acids in cowpea (acidic, basic, bolar and non polar amino acids) were decreased significantly with increase in gamma radiation compared to their respective controls. Hooshmand and Klopfenstein [35] reported that lysine content decreased by 7 and 13% respectively, when maize and wheat flours were irradiated at 7.5 kGy. The observed increase in some free amino acid content due to exposure to ionizing radiation is in agreement with the findings

reported by Satter *et al.* [36], who documented increases in essential and nonessential amino acids of soybean when irradiated at a dose level of 10 kGy.

The precise effect of ionizing radiation on free amino acid content depends on various factors, such as sensitivity of the exposed system, the type of particular functional tissue and even other conditions, such as aqueous soaking after irradiation, as has been indicated by Siddhuraju *et al.* [12]. Wang and vonSonntage [16] reported that sulphur-containing as well as aromatic amino acids are, in general, the most sensitive to irradiation, while simple amino acids could be formed by destruction of other amino acids. Bhat *et al.* [37] reported that radiation is an efficient process in maintaining the nutritional potential of *Mucuna pruriens* seeds. Fermentation of raw flour of the cultivar increased the level of lysine and phenylalanine plus tyrosine (Table 4). It has been reported that fermentation of Sicklepod leaves significantly ($P \leq 0.05$) increased alanine, valine, cystine, isoleucine, leucine and ammonia contents while aspartic, threonine, serine, glutamic, tyrosine, histidine, lysine and arginine contents were significantly ($P \leq 0.05$) decreased [7]. Radiation at 5, 10 and 15 kGy followed by fermentation significantly ($P \leq 0.05$) increased the level of phenylalanine plus tyrosine of the cultivar compared to the respective control.

Cooking of raw flour of the cultivar slightly increased the level of leucin and significantly ($P \leq 0.05$) that of phenylalanine plus tyrosine. Fageer *et al.* [38] observed that cooking of corn in water increased lysine, valine, leucine and phenylalanine while threonine, methionine and isoleucine were decreased. Radiation at 5, 10 and 15 kGy followed by cooking significantly ($P \leq 0.05$) increased phenylalanine plus tyrosine of the cultivar. Cooking of fermented dough of radiated and non-radiated flour increased phenylalanine plus tyrosine. Osman *et al.* [7] reported that cooking of the fermented Sicklepod leaves significantly ($P \leq 0.05$) decreased aspartic, threonine, serine, glutamic acid, glycine, tyrosine, phenylalanine and arginine contents. The results obtained showed that the effect of radiation alone on amino acids composition was minor as both fermentation and cooking had negative effect on some amino acids. Moreover, radiation process is an ideal method in preserving material as reported for millet flour [5].

4. Conclusion

The observations about nutritional quality in the studied samples tend to suggest that radiation processing up to 15 kGy had little effects on their value and had slight effect on the amino acids of the flour whether raw or processed. Therefore, radiation can be applied to alleviate the severe problem of antinutrients. Moreover, when accompanied by traditional processing an improvement in protein and minerals availability was observed.

References

1. Abdelseed BH, Abdalla AH, Yagoub AA, Mohamed IA, Babiker EE. Some nutritional attributes of selected newly developed lines of sorghum after fermentation. J Agric Sci Tech. 2011; 13: 399-409.
2. Sendra E, Capellas M, Guamis B, Felipe X, Mor-Mur M, Pla R. Revisión: Irradiación de alimentos-aspectos generales/Review: Food irradiation-general aspects. Food Sci Tech Intern. 1996; 2: 1-11.
3. Mbarki R, Sadok S, Barkallah I. Influence of gamma irradiation on microbiological, biochemical, and textural properties of bonito (*Sarda sarda*) during chilled storage. Food Sci Tech Intern. 2008; 14: 367-373.
4. Bhavsar SP, Augustine SK, Kapadnis BP.. Effect of Physical and Chemical Treatments on *Campylobacter* Spiked into Food Samples. Food Sci Tech Intern. 2007; 13: 277-283.
5. Mohamed EA, Ali NA, Ahmed SH, Mohamed IA, Babiker EE. Effect of radiation process on anti-nutrients and HCl extractability of calcium, phosphorus and iron during processing and storage. Rad Phy Chem. 2010; 79: 791-796.
6. Idris WH, AbdelRahman SM, ELMaki HB, Babiker EE, EL Tinay AH. Effect of malt pretreatment on HCl extractability of calcium, phosphorus and iron of sorghum (*Sorghum bicolor*) cultivars. Intern J Food Sci Techn. 2007; 42: 194–199.
7. Osman NM, Mohamed AI, Babiker EE. Fermentation and cooking of Sicklepod (*Cassia obtusifolia*) leaves: Changes in chemical and amino acid composition, antinutrients and protein fractions and digestibility. Intern J Food Sci Techn. 2010; 45: 124-132.

8. Mohiedeen IE, Abdullahi HE, Abd Elmoneim OE, Babiker EE, Mallasy LO. Effect of fermentation and cooking on protein quality of maize (*Zea Mays Linnaeus*) cultivars. Intern J Food Sci Techn. 2010; 45: 1284-1290.
9. Mohamed Nour AA, MohamedAhmed IA, Babiker EE, Yagoub AA. Investigations on winter season Sudanese sorghum cultivars: effect of sprouting on the nutritional value. Intern J Food Sci Techn. 2010; 45: 884-890.
10. Sokrab AM, Isam AM, Babiker EE. Effect of malting and fermentation on antinutrients, and total and extractable minerals of high and low phytate corn genotypes. Intern J Food Sci Techn. 2012; 47: 1037-1043.
11. Aziz NH, El-Far FM, Shahin AAM, Roushy SM. Control of fusarium, moulds and fumonisin B₁ in seeds by gamma irradiation. Food Con. 2007; 18: 1337–1342.
12. Siddhuraju P, Makkar HPS, Becker K. The effect of ionizing radiation on antinutritional factors and the nutritional value of plant materials with reference to human and animal food. Food Chem. 2002; 78: 187–205.
13. WHO. High-dose Irradiation: Wholesomeness of Food Irradiated with Doses above 10 kGy. Geneva: WHO. 1999; WHO Technical Report-Series no, 890.
14. Lee S, Lee M, Song K. Effect of gamma-irradiation on the physicochemical properties of gluten films. Food Chem. 2005; 92: 621–925.
15. Erkan N, Ozden O. The changes of fatty acid and amino acid compositions in seabream (*Sparus aurata*) during irradiation process. Rad Phy Chem. 2007; 76: 1636–1641.
16. Wang D, von Sonntag C. Radiation induced oxidation of phenylalanine. In: Proceedings of the Workshop on Recent Advances on Detection of Irradiated Foods, BRC Information, Chem. Anal. Comm. of the Eur. Comm., Brussels. 1991; Pp. 212–217.
17. Helinski MEH, Hassan MM, El-Motasim WM, Malcolm CA, Knols BGJ, El-Sayed B. Methodology towards a sterile insect technique field release of *Anopheles arabiensis* mosquitoes in Sudan: irradiation, transportation, and field cage experimentation. Malaria J. 2008; 7: 1–10.

18. Price ML, Butler LG. Rapid visual estimation and spectrophotometric determination of tannin content of sorghum grain. *J Agric Food Chem.* 1978; 25: 1268–1273.
19. Wheeler EL, Ferrel RE. A method for phytic acid determination in wheat and wheat fractions. *Cer chem.* 1971; 48: 312–320.
20. Maliwal BP. *In vitro* method to assess the nutritive value of leaf concentrate. *J. Agric. Food Chem.* 1983; 31: 315–319.
21. Monjula S, John E. Biochemical changes and *in vitro* protein digestibility of the endosperm of germinating *Dolichos lablab*. *J Sci Food Agric.* 1991; 55: 229–238.
22. Chapman HD, Pratt PF. Method for the analysis of soil, plant and water 2nd. California University Agricultural Division, California, 1982; Pp. 170-178.
23. Chauhan BM, Mahjan L. Effect of natural fermentation on the extractability of minerals from pearl millet flour. *J Food Sci.* 1988; 53: 1576–1577.
24. Moore S, STAIN WH. Chromatographic amino acids determination by the use of automatic recording equipment methods. *Enzy.* 1963; 63: 819–831.
25. Snedecor GW, Cochran WG. Statistical Methods, 17th edn. Ames, IA: The Iowa State University Press. 1987; Pp. 221–222.
26. Idris WH, AbdelRahman S M, ELMaki HB, Babiker EE, EL Tinay AH. Effect of germination, fermentation and cooking on phytic acid and tannin contents and HCl-extractability of minerals of sorghum (*Sorghum biocolor*) cultivars. *J Food Tech.* 2005; 3: 410-416.
27. Osman M A. Changes in sorghum enzyme inhibitors, phytic acid, tannins and *in vitro* protein digestibility occurring during Khamir (local bread) fermentation. *Food Chem.* 2004; 88: 129-134.
28. Eltayeb MM, Hassan AB, Mashier SA, Babiker EE. Effect of processing followed by fermentation on antinutritional factors content of pearl millet (*Pennisetum glaucum* L.) cultivars. *P J Nut.* 2007; 6: 463-467.
29. Brigide P, Canniatti-Brazac A. Anti-nutrients and *in vitro* availability of iron in irradiated common beans (*Phaseolus vulgaris*). *Food Chem.* 2006; 98: 85-89.
30. Osman A M, Hassan AB, Osman GAM, Nagat M, Mohamed AH, Eiman ED, Babiker EE. Effects of gamma irradiation and/or cooking on nutritional quality of

- faba bean (*Vicia faba* L.) cultivars seeds. J Food Sci Techn. 2012; DOI10.1007/s13197-012-0662-7
31. El-Niely HFG. Effect of radiation processing on anti-nutrients, *in vitro* protein digestibility and protein efficiency ratio bioassay of legume seeds. Rad Phy Chem. 2007; 76: 1050-1057.
 32. Duodu KG, Minnaar A, Taylor JRN. Effect of cooking and irradiation on the labile vitamins and anti-nutrient content of a traditional African sorghum porridge and Spinach relish. Food Chem. 1999; 66: 21-27.
 33. AbdElhaleem WH, Mustafa AI, El Tinay AH, Babiker EE. Effect of fermentation, malt-pretreatment and cooking on antinutritional factors and *in vitro* protein digestibility of sorghum cultivars. P J Nutr. 2008; 7: 335-341.
 34. Joseph OA, Klasus M, Kwaku D, Amanda M. Functional properties of cowpea (*Vigna unguiculata* L. Walp) flours and pastes as affected by γ irradiation. Food Chem. 2005; 93: 103–111.
 35. Hooshmand H, Klopfenstein CF. Effect of gamma irradiation on the mycotoxin disappearance and amino acid contents on corn, wheat and soybeans with different moisture contents. Pl Food Hum Nutr. 1995; 47: 227-238.
 36. Satter A, Akhtar MA. Irradiation and germination effects on phytate, protein and amino acids of soybean. Pl Food Hum Nutr. 1990; 40: 185–195.
 37. Bhat R, Sridhar KR, Young C, Bhagwath AA, Ganesh S. Composition and functional properties of raw and electron beam-irradiated *Mucuna pruriens* seeds. Inter J Food Sci Techn. 2008; 43: 1338–1351.
 38. Fageer AM, Babiker EE, El Tinay AH. Effect of malt pretreatment and/or cooking on phytate and essential amino acids contents and *in vitro* protein digestibility of corn flour. Food Chem. 2004; 88: 261–265.

Table 1. Tannin (%), Phytate (mg/100g) and percent *in vitro* protein digestibility (IVPD) of sorghum cultivar Karamaka as influenced by radiation followed by traditional treatments

Radiation dose (KGy)	Treatment	Parameter		
		Tannin	Phytate	IVPD
0	Raw	2.90 ±0.23 ^a	272.00 ±0.16 ^a	35.20 ±0.17 ^h
	Fermented	1.752 ±0.07 ^e	129.20 ±1.98 ^m	40.69 ±0.10 ^d
	Cooked	1.137 ±0.00 ⁱ	217.60 ±0.23 ^d	14.12 ±0.09 ^p
	Fermented/cooked	0.586 ±0.32 ^m	160.480 ±0.1 ^h	20.49 ±0.16 ^l
5	Raw	2.572 ±0.22 ^b	236.37 ±2.13 ^b	37.75 ±1.86 ^g
	Fermented	1.657 ±0.01 ^f	112.53 ±0.11 ⁿ	41.51 ±1.85 ^c
	Cooked	1.050 ±0.02 ^j	191.92 ±0.21 ^f	15.01 ±1.06 ^o
	Fermented/cooked	0.528 ±0.03 ⁿ	138.98 ±0.13 ⁱ	21.56 ±0.01 ^k
10	Raw	2.242 ±0.01 ^c	226.58 ±0.05 ^c	38.47 ±0.09 ^f
	Fermented	1.352 ±2.91 ^h	132.30 ±1.95 ^k	42.18 ±1.82 ^b
	Cooked	0.913 ±0.05 ^k	109.69 ±0.11 ^o	15.56 ±2.91 ⁿ
	Fermented/cooked	0.465 ±0.15 ^o	136.41 ±0.84 ^j	21.70 ±0.01 ^j
15	Raw	2.227 ±0.020 ^d	217.33 ±0.04 ^e	38.86 ±0.05 ^e
	Fermented	1.358 ±0.02 ^g	104.40 ±0.12 ^p	42.71 ±1.99 ^a
	Cooked	0.840 ±0.05 ^l	176.91 ±1.56 ^g	15.83 ±0.01 ^m
	Fermented cooked	0.444 ±0.01 ^p	131.11 ±0.92 ^l	22.08 ±0.08 ⁱ

Values are means of three replicates ± SD. Means not sharing a common superscript(s) in a column are significantly different at $p \leq 0.05$ as assessed by Duncan's Multiple Range Test.

Table 2. Total (mg/100gm) and extractable (%) sodium (Na), potassium (K) and magnesium (Mg) of Karamaka cultivar as influenced by gamma radiation followed by traditional treatments

Radiation dose (kGy)	Treatment	Minerals					
		Na		K		Mg	
		Total	extractable	Total	extractable	Total	extractable
0	Raw	22.60 ±0.01 ^d	67.40 ±0.02 ^j	470.10 ±0.05 ^{fg}	50.00 ±0.01 ^m	50.00 ±1.98 ^{ef}	52.50 ±0.03 ^j
	Fermented	23.90 ±0.11 ^c	73.40 ±0.02 ^h	495.32 ±1.35 ^c	73.30 ±0.01 ^d	62.20 ±1.97 ^{ab}	75.00 ±0.19 ^e
	Cooked	19.87 ±2.04 ^h	69.90 ±0.11 ⁱ	458.00 ±0.08 ^k	46.20 ±0.78 ⁿ	47.40 ±0.01 ⁱ	50.60 ±0.25 ^k
	Fermented /cooked	20.30 ±0.03 ^h	72.30 ±2.01 ^h	469.80 ±1.06 ^g	70.00 ±0.03 ^e	56.80 ±2.03 ^b	72.30 ±0.14 ^g
5	Raw	22.10 ±1.09 ^{def}	76.60 ±2.97 ^g	470.00 ±1.76 ^{fg}	57.40 ±0.43 ⁱ	49.80 ±0.05 ^{fg}	75.00 ±2.42 ^e
	Fermented	24.50 ±0.43 ^b	88.60 ±0.09 ^b	479.40 ±0.53 ^d	75.10 ±0.52 ^c	64.40 ±0.98 ^a	76.10 ±0.07 ^d
	Cooked	21.33 ±0.75 ^g	70.50 ±0.52 ⁱ	450.00 ±0.87 ^l	53.30 ±0.01 ^l	43.80 ±0.08 ^j	50.00 ±0.09 ^k
	Fermented/cooked	22.31 ±0.98 ^{def}	79.90 ±0.52 ^f	466.00 ±0.09 ^h	67.10 ±0.06 ^f	53.80 ±1.23 ^c	73.30 ±0.01 ^f
10	Raw	22.52 ±1.18 ^{de}	81.40 ±3.03 ^e	473.00 ±0.38 ^e	60.00 ±0.42 ^h	50.40 ±1.23 ^e	67.30 ±0.78 ^h
	Fermented	25.3 ±0.87 ^a	97.70 ±0.52 ^a	496.40 ±0.42 ^b	81.20 ±0.64 ^b	65.50 ±0.42 ^a	100.0 ±0.07 ^a
	Cooked	21.91 ±0.76 ^f	86.60 ±0.64 ^c	460.60 ±0.98 ^j	56.20 ±0.92 ^j	47.20 ±0.76 ⁱ	55.90 ±0.09 ⁱ
	Fermented/cooked	21.93 ±0.23 ^f	86.60 ±0.85 ^c	465.00 ±0.65 ⁱ	80.40 ±0.21 ^b	51.20 ±0.01 ^d	92.50 ±0.05 ^b
15	Raw	22.5 ±1.97 ^{de}	84.00 ±2.43 ^d	470.50 ±0.12 ^f	63.40 ±0.91 ^g	49.60 ±1.13 ^g	75.10 ±0.23 ^{de}
	Fermented	25.00±0.76 ^{ab}	88.00 ±0.53 ^b	500.00 ±0.75 ^a	86.00 ±0.23 ^a	66.40 ±0.18 ^a	83.50 ±0.55 ^c
	Cooked	21.90 ±1.87 ^f	81.40 ±0.09 ^e	450.00 ±0.12 ^l	54.80 ±0.43 ^k	48.20 ±0.98 ^h	73.90 ±0.91 ^f
	Fermented /cooked	22.00 ±0.23 ^{ef}	86.63 ±0.65 ^c	473.20 ±0.76 ^e	85.80 ±0.98 ^a	53.80 ±0.65 ^c	91.00 ±0.01 ^b

Values are means of three replicates (± SD). Means not sharing a common superscript(s) in a column are significantly different at $p \leq 0.05$ as assessed by Duncan's Multiple Range Test.

Table 3. Total (mg/100gm) and extractable (%) trace minerals of Karamaka cultivar as affected by gamma irradiation followed by traditional treatments

Radiation dose (KGy)	Treatment	Minerals							
		Zn		Cu		Mn		Co	
		Total	extractable	Total	extractable	Total	extractable	Total	extractable
0	Raw	2.54 ±0.01 ^a	49.29 ±0.05 ^o	0.42 ±0.17 ^a	34.17 ±0.09 ^l	3.35 ±1.24 ^a	39.42 ±0.01 ^m	0.15 ±0.88 ^a	62.07 ±1.07 ^l
	Fermented	2.57 ±0.11 ^a	71.02 ±0.23 ^g	0.48 ±0.02 ^a	42.01 ±0.45 ⁱ	3.42 ±0.17 ^a	64.08 ±0.26 ^g	0.17 ±0.04 ^a	81.89 ±0.71 ^e
	Cooked	2.52 ±0.12 ^a	49.29 ±0.01 ^o	0.42 ±0.01 ^a	33.43 ±0.56 ^m	3.29 ±0.13 ^a	33.47 ±0.11 ⁿ	0.14 ±0.11 ^a	62.44 ±0.52 ^l
	Fermented /cooked	2.55 ±0.46 ^a	67.43 ±0.98 ⁱ	0.44 ±0.32 ^a	41.60 ±0.09 ⁱ	3.37 ±0.98 ^a	63.70 ±0.07 ^g	0.18 ±0.13 ^a	78.80 ±0.08 ^f
5	Raw	2.51 ±0.01 ^a	52.58 ±0.32 ^m	0.41 ±0.02 ^a	40.90 ±0.01 ^j	3.30 ±0.12 ^a	42.90 ±0.16 ^l	0.19 ±0.08 ^a	65.70 ±0.13 ^j
	Fermented	2.57 ±0.75 ^a	84.51 ±0.63 ^e	0.42 ±0.07 ^a	54.90 ±0.31 ^e	3.37 ±0.13 ^a	65.90 ±0.75 ^f	0.21 ±0.54 ^a	86.00 ±0.01 ^d
	Cooked	2.55 ±0.09 ^a	51.12 ±0.51 ⁿ	0.49 ±0.23 ^a	36.30 ±0.97 ^k	3.31 ±0.98 ^a	48.10 ±0.35 ^k	0.14 ±0.09 ^a	64.30 ±0.06 ^k
	Fermented/cooked	2.55 ±0.16 ^a	82.86 ±0.09 ^f	0.40 ±0.95 ^a	49.10 ±0.94 ^f	3.35 ±0.09 ^a	68.10 ±0.24 ^e	0.15 ±0.87 ^a	82.60 ±0.15 ^e
10	Raw	2.51 ±0.14 ^a	64.14 ±0.81 ^k	0.43 ±0.23 ^a	44.00 ±0.34 ^h	3.32 ±0.04 ^a	50.00 ±1.05 ^j	0.17 ±1.92 ^a	70.00 ±0.05 ^h
	Fermented	2.58 ±0.32 ^a	95.20 ±0.91 ^b	0.46 ±0.81 ^a	63.90 ±0.12 ^b	3.34 ±0.65 ^a	82.30 ±0.86 ^b	0.26 ±0.54 ^a	94.60 ±0.75 ^a
	Cooked	2.51 ±0.43 ^a	60.54 ±0.37 ^l	0.47 ±0.76 ^a	46.90 ±0.98 ^g	3.32 ±0.01 ^a	51.10 ±0.43 ⁱ	0.15 ±0.98 ^a	69.70 ±0.43 ^h
	Fermented /cooked	2.55 ±0.75 ^a	89.65 ±0.83 ^d	0.41 ±0.01 ^a	61.90 ±0.42 ^c	3.36 ±0.06 ^a	71.90 ±0.09 ^d	0.17 ±0.81 ^a	87.40 ±0.01 ^c
15	Raw	2.54 ±0.23 ^a	69.67 ±0.65 ^h	0.45 ±0.05 ^a	44.20 ±0.16 ^h	3.31 ±0.43 ^a	53.40 ±0.07 ^h	0.15 ±0.06 ^a	73.80 ±0.05 ^g
	Fermented	2.54 ±0.65 ^a	97.62 ±0.71 ^a	0.42 ±0.54 ^a	68.60 ±0.43 ^a	3.32 ±0.01 ^a	88.40 ±0.75 ^a	0.18 ±0.98 ^a	93.10 ±0.43 ^b
	Cooked	2.51 ±0.94 ^a	66.43 ±0.93 ^j	0.42 ±0.91 ^a	44.00 ±0.53 ^h	3.30 ±0.01 ^a	51.40 ±0.45 ⁱ	0.19 ±0.61 ^a	68.40 ±0.22 ⁱ
	Fermented /cooked	2.55 ±0.23 ^a	91.48 ±1.09 ^c	0.44 ±0.38 ^a	60.20 ±0.05 ^d	3.32 ±0.02 ^a	77.70 ±0.11 ^c	0.20 ±0.01 ^a	85.40 ±0.16 ^d

Values are means of three replicates (± SD). Means not sharing a common superscript(s) in a column are significantly different at $p \leq 0.05$ as assessed by Duncan's Multiple Range Test.

Table 4. Amino acids composition (mg/100gm protein) and score (AAS) of Karamaka cultivar as affected by gamma radiation followed by traditional treatments

Radiation dose (KGy)	Treatment	Amino acid													
		Isoleucine		Leucine		Lysine		Methionine +Cysteine		Phenyl alanine + Tyrosine		Threonine		Valine	
		Total	AAS	Total	AAS	Total	AAS	Total	AAS	Total	AAS	Total	AAS	Total	AAS
0	Raw	3.50 ^a	125.00	12.98 ^c	196.67	1.91 ^b	32.93	1.79 ^a	71.60	6.94 ^p	110.10	1.970 ^a	57.00	4.62 ^a	132.00
	Fermented	3.17 ^d	113.20	12.52 ^h	189.62	1.98 ^a	34.13	1.77 ^b	70.84	8.55 ^e	135.70	1.42 ^c	41.88	4.24 ^e	121.10
	Cooked	3.05 ⁱ	108.90	13.00 ^b	196.91	0.98 ^k	16.90	1.66 ⁱ	66.52	8.34 ^g	132.40	1.38 ^g	40.60	4.21 ^g	120.20
	Fermented/cooked	3.09 ^g	110.30	12.87 ^d	195.05	1.22 ^g	21.03	1.76 ^c	70.50	8.06 ^j	127.90	1.31 ⁱ	38.56	4.15 ⁱ	118.50
5	Raw	2.72 ^l	97.00	12.16 ^j	184.18	1.33 ^f	22.90	1.71 ^h	68.40	7.33 ⁿ	116.30	1.11 ^m	32.50	3.77 ⁿ	107.60
	Fermented	3.30 ^b	117.90	8.91 ^k	134.90	1.88 ^c	32.46	1.72 ^f	68.70	8.68 ^b	137.80	1.41 ^d	41.50	4.47 ^b	127.80
	Cooked	3.08 ^h	110.10	8.130 ^m	123.20	0.79 ⁿ	13.62	1.64 ^k	65.70	8.42 ^f	133.70	1.40 ^f	41.10	4.24 ^d	121.20
	Fermented/cooked	3.01 ^j	107.30	12.34 ⁱ	187.00	1.13 ^h	19.55	1.76 ^d	70.30	8.15 ⁱ	129.40	1.10 ⁿ	32.30	4.10 ^j	117.00
10	Raw	2.50 ^o	89.40	12.74 ^f	193.00	0.93 ^l	16.03	1.65 ^j	66.04	7.07 ^o	112.20	0.83 ^o	24.40	3.81 ^l	99.50
	Fermented	2.84 ^k	101.40	6.62 ^o	100.30	1.81 ^d	31.27	1.54 ⁿ	61.60	7.74 ^k	122.80	1.28 ^j	37.60	3.82 ^k	109.20
	Cooked	3.12 ^f	111.30	8.37 ^l	126.80	0.62 ^o	10.71	1.73 ^e	69.00	8.61 ^d	136.70	1.57 ^b	46.30	4.24 ^f	121.10
	Fermented/cooked	3.19 ^c	113.80	12.59 ^g	190.80	1.09 ⁱ	18.72	1.62 ^l	65.00	8.25 ^h	130.90	1.40 ^e	41.20	4.38 ^c	125.20
15	Raw	2.50 ^p	89.36	13.00 ^a	196.95	0.92 ^m	15.86	1.58 ^m	62.70	7.36 ^m	116.80	0.83 ^o	24.50	3.38 ^p	96.49
	Fermented	2.70 ^m	96.40	6.38 ^p	96.60	1.81 ^e	31.12	1.51 ^p	60.30	7.73 ^l	122.70	1.19 ^k	34.90	3.78 ^m	107.90
	Cooked	2.62 ⁿ	93.70	7.16 ⁿ	108.40	0.54 ^p	9.33	1.51 ^o	60.40	8.67 ^c	137.60	1.34 ^b	39.90	3.51 ^o	100.30
	Fermented/cooked	3.12 ^e	111.50	12.76 ^e	193.30	1.00 ^j	17.28	1.71 ^g	68.60	8.69 ^a	137.90	1.18 ^l	34.70	4.19 ^h	119.60

Values are means of duplicate samples. Means not sharing a common superscript(s) in column are significantly different at $p \leq 0.05$ as assessed by Duncan's Multiple Range Test.