

1 Sensory & nutritional quality characteristics of
2 **Quality of Kunun zaki enriched with Moringa**
3 **(Moringa oleifera) seed flour**
4

5 **Olosunde O.O., Abiodun O.A., Amanyunose, A.A. and Adepeju, A.B**

6
7
8 *Department of Food Science and Technology, Osun State Polytechnic, P.M.B.301, Iree,*
9 *Osun State, Nigeria*

10
11
12
13
14
15 **ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)**
16

Aims:The aim of this work was to determine the ^{nutritional} quality of kunun zaki enriched with moringa seed flour

Study design:Analyses of variance (ANOVA).

Place and Duration of Study:Department of Food Science and Technology, Osun State Polytechnic, Iree, Nigeria, between July 2013 to January 2014

Methodology:Sorghum grains were soaked, washed and then mixed with the spices and wet milled.Moringa seed flour (5, 10, 15%) was added to the kunun slurry. The slurry was divided into two; one portion was cooked and allowed to cool to 45°C and then the uncooked portion was added and mixed thoroughly. The mixture was allowed to ferment for 12 h and sieved. Proximate, mineral, physicochemical, anti-nutritional and sensory evaluation was determined on the enriched kunun zaki

Results:There were reductions in the moisture and carbohydrate contents of kunun zaki with increase in moringa seed flour incorporation while the protein, fat, ash and crude fibre contents increased. Kunun zaki with 15 % moringa seed flour had higher values in all the mineral contents determined. pH and total soluble solid increased as the percentage of moringa seed flour increased. Kunun zaki with 15 % moringa seed flour had higher anti-nutritional contents than other samples. Sensory analysis conducted on the kunun zaki revealed no significant difference ($p=0.05$) in the taste, appearance and overall acceptability of control and 5 to 10 % level of moringa seed substitution.

Conclusion:The proximate and mineral contents of the substituted kunu zaki were increased with the incorporation of moringa seed flour. The anti-nutrients in the beverage were low and within the range permissible for food. Addition of up to 10 % moringa seed flour to kunun zaki is desirable as higher concentrations (15%) of moringa seed flour impact undesirable effect on the taste of the product.

17
18 **Keywords:** [Kunun zaki, Moringa oleifera, Proximate composition, Anti-nutrient, Sorghum,
19 Substitution]

20
21
22 **1. INTRODUCTION**

23 Kunun zaki is a traditionally fermented non-alcoholic beverages mostly consumed in
24 Northern Nigeria [1-2]. Among the several types of kunun, kunun zaki is the most preferred

* Tel.: +2348030701354

E-mail address: funmiabiodun2003@yahoo.com

according to Gaffa et al.[3]. Preparation of kunun zaki varies amongst people and can generally be produced from either the following substrates; millet (*Pennisetum typhoideum*), maize (*Zea mays*) or sorghum (*Sorghum bicolor*), but millet is the most common substrates [4]. Spices such as ginger, black pepper, red pepper, cloves and sugar are commonly added as flavor and taste improver [5]. The method of production of kunun zaki is crude involving only household utensils and varies from locality to locality [6]. Kunun zaki is consumed as a beverage with or without addition of sugar as a sweetener[7]. Its consumption is high among all age groups and is fast spreading in Nigeria as it is cheaper than the carbonated drinks [8]. Kunun-zaki was reported to have low nutritional status providing substantial carbohydrate value [9, 10]. The beverage is deficient in protein since it is majorly from cereal crops [3]. Therefore, protein-energy malnutrition is prevalent in the areas where kunun serves as their major food. There is need to improve the protein contents of kunun zaki to solve the problem of malnutrition in our society. Due to this, Gaffa et al [11] enriched kunun zaki with soybean and the kunun zaki was acceptable at 4 % level. Substitution with *Moringa oleifera* seed flour could also improve the protein contents of kunun zaki as it has been found to contain essential nutrients needed by the body.

Moringa oleifera (Syn. *M. pterygosperma* Gaertn.) is of the family Moringaceae. It is a small graceful tree with sparse foliage, often planted in compounds or used as hedge in northern Nigeria [12]. *Moringa oleifera* is a fast growing, aesthetically pleasing small tree. The specie is characterized by its long, drumstick shaped pods that contain its seeds within the first year of growth [13]. It is well known for its nutritional and medicinal values by many communities in northern Nigeria [12]. *Moringa oleifera* was reported to be a good source of vitamins and amino acids [14]. In developing countries, moringa has potential to improve nutrition, boost food security, foster rural development and support sustainable land care [15]. Therefore, this paper presents the quality of kunun zaki substituted with moringa seed flour

2. MATERIAL AND METHODS

2.1 Materials

Sorghum and spices were obtained at Igbona market, Osogbo, Osun State, Nigeria, while *Moringa* seeds were obtained at IAR&T research Institute, Ibadan, Oyo State, Nigeria.

2.2 Methods

2.2.1 Production of kunun zaki

The grains (sorghum) were cleaned and steeped in tap water for 24 h. The grains were washed and then mixed with the spices (ginger, cloves, black pepper and cinnamon) and wet milled. The paste was divided into two portions; one portion was cooked and allowed to cool to 45°C and then the uncooked portion was added and mixed as described by Efiuvwevwere and Akoma [16] and Evans et al. [17]. The mixture was allowed to ferment for 12 h after which it was sieved.

2.2.2 Production of moringa seed flour and substitution

Moringa seeds were removed from the pods, dried in the oven and milled. *Moringa* seed flour (5, 10, 15%) was added to the kunun slurry. The slurry was divided into two; one portion was cooked and allowed to cool to 45°C and then the uncooked portion was added and mixed thoroughly. The mixture was allowed to ferment for 12 h and sieved.

mixed thoroughly

2.2.3 Proximate composition of kunun zaki substituted with moringa seed flour

* Tel.: +2348030701354

E-mail address: funmiabiodun2003@yahoo.com

delete footnote

Moisture content, crude protein, crude fibre, crude fat and ash were determined as described by AOAC [18]. Carbohydrate was calculated by difference while the metabolizable energy was calculated using the method of Adubiaro et al. [19].

2.2.4 Mineral composition of kunun zaki substituted with moringa seed flour

Kunun zaki (0.5 g) was weighed into a clean ceramic crucible. A blank was prepared with empty crucible. The crucible was placed in a muffle furnace at 500 °C for 4 hr. The sample was allowed to cool down in the oven after which it was removed carefully. The ashed sample was poured into already labeled 50 ml centrifuge tube. The crucible was rinsed with 5 ml of distilled water into the centrifuge tube. The crucible was rinsed again with 5 ml of aqua regia. This was repeated to make a total volume of 20 ml. The sample was mixed properly and centrifuged (IEC Centra GP8) for 10 min at 301.86 g. The supernatant was decanted into clean vials for mineral determination. The absorbance was read on atomic absorption spectrophotometer (Buck Scientific Model 200A) at different wavelength for each mineral element (Ca-422.7nm, Fe-248.3 nm, Mg-285.2 nm, Mn-279.5 nm) [20]

2.2.5 Physico-chemical analysis of kunun zaki substituted with moringa seed flour

pH of the kunun zaki was measured using digital pH meter (ELICO L1 614 pH analyser) and expressed as pH units. Total soluble solids (TSS) as % Brix was determined using digital ATAGO refractometer (ATAGO, PAL-Maple Pocket type). Total titratable acidity as % lactic acid was determined by titration using 0.1N sodium hydroxide and phenolphthalein indicator solution [21-22]. The relative amount of lactic acid was calculated as follows:

$$\% \text{Lactic acid} = \frac{\text{Volume of 0.1N NaOH} \times \text{Normality of Alkali} \times 9}{\text{Volume of sample}}$$

This is not the acid factor for lactic acid
= 0.09003 [0.09]

2.2.6 Antinutritional analysis of kunun zaki substituted with moringa seed flour

Method of Dairo [23] was used for phytate determination while oxalate content was determined using the method of Nwinuka et al. [24]. Total phenol and alkaloid determination were done using method of Obadoni and Ochuko [25] and Sahoré et al. [26] respectively.

* methods

2.2.7 Sensory evaluation of kunun zaki substituted with moringa seed flour

Sensory evaluation was done on each of the kunun-zaki samples produced by a panel of 20 judges comprising of experienced lecturers at Osun State Polytechnic, Iree, Nigeria, as described by Evans et al. [17]. The quality characteristics including appearance, aroma, taste and overall acceptability of the samples were evaluated based on a seven point hedonic scale (where 1 = like extremely, 2 = like very much, 3 = like slightly, 4 = neither like nor dislike, 5 = dislike slightly, 6 = dislike very much, 7 = dislike extremely).

experienced in what? Are they familiar with kunun-zaki?

2.3 Statistical analysis

The analyses were carried out in triplicate. The mean and standard deviation of the data obtained were calculated using SPSS (17.0).

The mean scores were computed and significant differences among the mean were determined using SPSS version 17 (SPSS, 2011).

3. RESULTS AND DISCUSSION

The proximate composition of kunun zaki substituted with moringa seed flours are shown in Table 1. Moisture contents ranged from 81.37 to 85.33 %. Moisture contents obtained were slightly lower than the values reported by Gaffa and Ayo [7], Ogbonna et al. [27] and Gaffa et al. [11] for kunun zaki enriched with soybean. Difference in the moisture contents may be due to the methods used in preparing the kunun zaki. Kunun zaki consists of majorly water to quench our thirst. It had low shelf stability due to high moisture content observed and they readily undergo microbial induced spoilage within 2 to 3 days of production [10]. Moisture content of the kunun zaki decreased with increase in moringa seed flour substitution.

* Tel.: +2348030701354

E-mail address: funmiabiodun2003@yahoo.com

Kunun zaki substituted with 15 % moringa seed flour was significantly different ($p=0.05$) from other kunun zaki in ash content. There was no significant difference ($p=0.05$) in the control and kunun zaki substituted with 5 % moringa seed flour in ash contents. Gaffa and Ayo [7] reported 1.22 % ash content for traditionally processed kunun zaki. The highest ash content (1.96 %) was observed in kunun zaki substituted with 15 % moringa seed flour. Ash contents increased with increase in moringa seed flour substitution.

Fat content ranged from 1.67 to 3.33 %. There were significant differences ($p=0.05$) in the fat contents of the kunun zaki with higher value in kunun with 15 % moringa seed flour. Lower fat contents were observed in the control but there was increase in the fat contents of the substituted kunun zaki which could be due to incorporation of moringa seed flour. Fat content of moringa seed flour was reported to be high (45.84 %) [28]. The storage life of the substituted kunun zaki may however be reduced owing to increase in fat content and high fat content in kunun zaki may lead to susceptibility of the kunun to oxidative rancidity.

Crude fibre contents of the kunun zaki increased with moringa seed flour substitution. The highest value (1.52 %) was in kunun with 15 % moringa seed flour. There were significant differences ($p=0.05$) in crude fibre contents of all the kunun zaki produced. This was due to addition of moringa seed flour. Moringa seed flour had 7.73 % crude fibre as reported by Abiodun et al. [28] showing that the crude fibre of moringa seed flour was high. The seed contain appreciable amount of crude fibre which aid in digestion process.

Protein contents increased from 2.96 to 6.73 % with moringa seed flour substitution. The least protein content (2.96 %) was in the control sample. Increase in protein content in the kunun zaki was recorded with higher value at 15 % moringa seed flour substitution. The high protein content of moringa seed flour (28.04 %) gives an indication of their usefulness in human diet and as animal feed [28-29]. Therefore, moringa seed flour had tendency to improve the protein contents of kunun zaki.

Carbohydrate contents decreased with increase in moringa seed flour substitution. It ranged from 5.09 to 9.63 %. The control was significantly different ($p=0.05$) from the substituted kunun zaki in carbohydrate content. Addition of moringa seed flour caused reduction in the carbohydrate contents and increase in protein content of the kunun zaki. The beverage serves as a good source of energy.

Table 1: Proximate composition of Kunun zaki substituted with moringa seed flour

Parameter	Level of moringa seed flour substitution (%)			
	0	5	10	15
Moisture content (%)	85.33±0.61a	83.03±0.70b	82.38±0.40c	81.37±0.10d
Ash content (%)	1.03±0.11 c	1.08±0.10c	1.73±0.10b	1.96±0.18a
Fat (%)	1.67±0.10d	2.20±0.05c	2.51±0.05b	3.33±0.13a
Crude fibre (%)	0.88±0.41d	1.01±0.11c	1.20±0.40b	1.52±0.40a

Crude protein (%)	2.96±0.20d	4.63±0.26c	5.57±0.32b	6.73±0.30a
Carbohydrate (%)	9.63±0.09a	8.05±0.07b	6.61±0.10c	5.09±0.13d
Energy (KJ/100g)	275.82±0.62c	296.96±0.77b	299.93±0.90b	324.15±0.50a

Values with the same subscript along the row are not significantly different (p=0.05)

The mineral compositions of kunun zaki are shown in Table 2. The major mineral contents in the control and substituted kunun zaki were calcium and magnesium. There were no significant differences (p=0.05) in the calcium content of 10 and 15 % moringa seed flour substituted kunun zaki. Likewise, no significant differences in the manganese and nickel compositions of kunun zaki substituted with 5 and 10 % moringa seed flour. All the mineral contents increased with increase in moringa seed flour substitution. There were significant differences (p=0.05) in magnesium and iron contents of the kunu zaki. Magnesium and iron contents increased rapidly with increase in moringa seed flour. This showed that moringa seed flour serves as a source of essential minerals needed by the body.

Table 2: Mineral composition (ppm) of Kunun zaki substituted with moringa seed flour

Parameter	Level of moringa seed flour substitution (%)			
	0	5	10	15
Ca	30.02±0.06c	35.42±0.35b	40.22±0.07a	40.65±0.10a
Mg	15.27±0.19d	25.10±0.37c	30.11±0.16b	55.78±0.08a
Fe	0.80±0.06d	1.38±0.05c	1.65±0.05b	2.58±0.02a
Mn	0.08±0.02c	0.14±0.11b	0.15±0.10b	0.23±0.02a
Ni	0.09±0.10c	0.15±0.02b	0.16±0.02b	0.21±0.04a

Values with the same subscript along the row are not significantly different (p=0.05)

Physico-chemical properties of kunun zaki substituted with moringa seed flour are shown in Fig. 1. pH, total soluble solid and titratable acidity ranged from 3.20 to 3.90, 3.20 to 5.20 °Brix and 0.34 to 1.25 % respectively. pH and total soluble solid increased with increase in moringa seed flour addition while the titratable acidity decreased with moringa addition. pH obtained were lower than the observed value (4.10-4.49) for instant kunun zaki substituted with mango mesocarp flour [1] and 3.80-4.08 for kunun zaki reported by Ogbonna et al. [27]. Increase in level of moringa seed flour incorporation reduced the acidity of the products therefore could affect the storability of the kunun zaki.




Fig. 1: Physico-chemical properties of kunun zaki substituted with moringa seed flour

Anti-nutritional compositions of kunun are presented in Table 3. Phytate, oxalate, tannin, phenol and alkaloid contents of kunun zaki substituted with moringa seed flour ranged from 0.12 to 0.33, 0.09 to 0.36, 0.10 to 0.64, 0.08 to 0.13 and 0.10 to 1.02 mg/100g respectively. Kunun zaki without moringa had the least values in all the anti-nutrients determined. There were no significant differences ($p=0.05$) in the kunun zaki with 5 and 10 % moringa seed flour in phytate and phenol contents. There were significant differences ($p=0.05$) in oxalate, tannin and alkaloid levels of the substituted kunun zaki. All the anti-nutrients increased with increase in moringa seed flour substitution. The levels of antinutrients in the kunun zaki were low and within the permissible level for food [30].

Table 3: Antinutritional composition of kunun zaki substituted with moringa seed flour

Parameter	Level of moringa seed flour substitution (%)			
	0	5	10	15
Phytate (mg/100g)	0.12±0.21c	0.19±0.15b	0.21±0.11b	0.33±0.19a
Oxalate (mg/100g)	0.09±0.16d	0.18±0.14c	0.23±0.09b	0.36±0.20a
Tannin (mg/100g)	0.10±0.11d	0.23±0.05c	0.41±0.13b	0.64±0.10a
Phenol (mg/100g)	0.08±0.10c	0.10±0.10b	0.11±0.11b	0.13±0.09a
Alkaloid(mg/100g)	0.10±0.10d	0.52±0.10c	0.76±0.10b	1.02±0.12a

Values with the same subscript along the row are not significantly different ($p=0.05$)

Table 4 showed the sensory properties of kunun zaki. Kunun zaki substituted with 15 % moringa seed flour was significantly different ($p = 0.05$) in appearance than other samples. There was no significant difference ($p = 0.05$) in the appearance of control and kunun zaki substituted with 5 and 10 % moringa seed flour. At 15 % level, settling and coagulation of the components of kunun zaki was noticed. This may be due to the natural coagulating properties exhibited by the seeds of moringa reported by Santos et al. [31] and Ogbe and Affiku[32]. Likewise the taste of kunun zaki substituted with 15 % moringa seed flour was significantly different ($p=0.05$) from other samples. The taste of up to 10 % moringa seed flour substitution was acceptable while above 10 % objectionable taste was perceived. The aroma of the substituted kunun zaki were not significantly different ($p=0.05$) from the control sample. Overall acceptability showed that kunun zaki substituted with up to 10 % moringa seed flour were acceptable

Table 4: Sensory properties of kunun zaki substituted with moringa seed flour

Parameter	Level of moringa seed flour substitution (%) ^{1, 2}			
	0	5	10	15
Appearance	1.43b	1.40b	1.47b	1.58a
Taste	1.95b	1.99b	2.09b	4.23a
Aroma	2.17a	2.14a	2.12a	2.15a
Overall acceptability	1.58b	1.60b	1.62b	3.84a

¹ Values with the same subscript along the row are not significantly different ($p=0.05$)

² 7 point hedonic scale (1 = like extremely, 3 = neither like nor dislike, 7 = dislike extremely)

* Tel.: +2348030701354

E-mail address: funmiabiodun2003@yahoo.com

4. CONCLUSION

The proximate and mineral contents of the substituted kunu zaki were increased with the incorporation of moringa seed flour. The anti-nutrients in the beverage were low and within the range permissible for food. Addition of up to 10 % moringa seed flour to kunun zaki is desirable as higher concentrations (15%) of moringa seed flour impact undesirable effect on the taste and appearance of the product.

"Authors have declared that no competing interests exist."

AUTHORS' CONTRIBUTIONS

'Author AOA' designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. 'Author OOO and 'Author AAA managed the analyses of the study. 'Author ABA managed the literature searches. All authors read and approved the final manuscript.'

REFERENCES

- 1 Sengey IA, Akpapunam MA and Ingbian EK. Physicochemical and sensory properties of instant kunun zaki flour blends from sorghum and mango mesocarp flours. *Nigerian Food Journal* 2012; 30 (2): 8-16
- 2 Udensi EA, Odom TC and Dike CO. Comparative studies of ginger (*Zingibr officinale*) and West African black pepper (*Piper guineense*) extracts at different concentrations on the microbial quality of soymilk and kunun zaki. *Nigerian Food Journal* 2012; 30 (2): 38-43
- 3 Gaffa T, Jideani IA and Nkama I. Traditional production, consumption and storage of kunun-zaki: a non-alcoholic cereal beverage. *Plant Food for Human Nutrition* 2002a; 57: 73-81
- 4 Akoma O, Jiya EA, Akumka DD and Mshilia E. Influence of malting on the nutritional characteristics of kunu-zaki. *African Journal of Biotechnology* 2006; 5 (10): 996 – 1000.
- 5 Ahmed EU, Musa N, and Ngoddy PO. 2003. Sensory attributes of extruded Cereal-Legume Blends for instant "kunu-zaki" Beverage analogue. *Proceedings of the 27th Annual NIFIST Conference*. Pp. 84.
- 6 Adeyemi IA and Umar S. Effect of method of manufacture on quality characteristics of kunun zaki, a millet based beverage. *Nigerian Food Journal* 1994; 12: 34-41.
- 7 Gaffa T and Ayo JA. Innovations in the Traditional Kunun Zaki Production Process. (2002). *Pakistan Journal of Nutrition* 2002; 1 (5): 202-205
- 8 Akpapunam MA, Badifu GO and Maduagwu PE. Assesment of production practices and evaluation of product characteristics of kunun zaki produced in Makurdi metropolis. Paper presented at NIFST Annual Conference held at Uyo, Nov 21-26
- 9 Edward-Inatimi B, Abasiokong SF, Chiemeka I. Kunun-zaki and Tsamiya non-alcoholic beverages prepared from sorghum grains. Chemical analysis for nutrient

* Tel.: +2348030701354

E-mail address: funmiabiodun2003@yahoo.com

- 276 content of fresh and ageing samples. Nigerian Journal of Biotechnology 1988; 5:
277 21-22.
- 278 10 Egbere OJ, Pam KV, Adesheyan KD, A'Kadir T and Oyero SK. Effects of
279 pasteurisation on survival patterns of microorganisms and vitamin C retention in
280 kunun-zaki. African Journal of Biotechnology 2009; 8 (23): 6603-6607
- 281 11 Gaffa T, Jideani IA and Nkama I. Soybean seed in kunun zaki beverage production.
282 Pakistan Journal of Biological Sciences 2002b; 5 (9): 970-973
- 283 12 Kawo, A.H., Abdullahi, B.A., Gaiya, Z.A., Halilu, A., Dabai, M. and Dakare, M.A.
284 (2009). Preliminary phytochemical screening, proximate and elemental composition
285 of moringa oleifera lam seed powder. Bayero Journal of Pure and Applied Sciences,
286 2(1): 96 - 100
- 287 13 Ajar, P.M., Ibiam, U.A., Uraku, A.J., Orji, O.U., Offor C.E. and Nwali, B.U. (2013).
288 Comparative Proximate and Mineral Composition of *Moringa oleifera* Leaf and Seed.
289 Global Advanced Research Journal of Agricultural Science, 2(5): 137-141
- 290 14 Olugbemi TS, Mutayoba SK and Lekule FP. Effect of Moringa (*Moringa oleifera*)
291 Inclusion in Cassava based diets to broiler chickens. International Journal of
292 Poultry Science 2010; 9 (4): 363-367.
- 293 15 National Research Council (2006). Moringa. Lost crops of Africa; Volume II:
294 Vegetables. National Academies Press. P 247.
- 295 16 Efiuvwevwere BJO and Akoma O. The microbiology of Kunu-Zaki, a cereal
296 beverage from northern Nigeria, during the fermentation (production) process. World
297 Journal of Microbiology & Biotechnology 1995; 11: 491-493.
- 298 17 Evans CE, Omoaruefike EO and Mohammed AV. Effect of Fermentation
299 Variables on Nutritional and Organoleptic Scores of Kunu-Zaki Produced From Rice
300 and Acha Blends. Advance Journal of Food Science and Technology 2013; 5(9):
301 1117-1119
- 302 18 AOAC. Association of Official Analytical Chemists Official methods of Analysis Vol. II
303 17th edition Washington, DC, USA, Official methods 925.09, 923.03, 979.09,
304 962.09, 4.5.1, and 923.05. 2000.
- 305 19 Adubiaro HO, Olaofo O, Akintayo ET and Babalola OO. Chemical composition,
306 calcium, zinc and phytate interrelationships in *Adansonia digitata* seed flour.
307 Advance Journal of Food Science and Technology 2011; 3(4): 228-232
- 308 20 Novozamsky I, Houba VJG, Van ECK and Van VW. Plant nitrogen and
309 phosphorus in plant tissue, novel digestion technique for multi-element. Plant
310 analysis communication in soil science and plant analysis 1983; 14: 239-248.
- 311 21 Lawal AK, Oyedoyin OB and Olatunji OO. Fate of pathogenic bacteria during
312 fermentation of cereal porridge (ogi)-A weaning food formula. Nigerian Food Journal
313 2009; 27 (10): 19-26
- 314 22 Talasila U, Vechalapu RR and Shaik KB. Preservation and Shelf life Extension of
315 Cashew Apple Juice. Internet Journal of Food Safety 2011;13: 275-280
- 316 23 Dairo FAS. Performance and haematological evaluation of weaner rabbits fed
317 loofahgourd seed meal (*Luffa cylindrica* m.j.roem). African Journal of Food
318 Agriculture and Nutritional Development 2008; 8(4): 451-463.
- 319 24 Nwinnuka NM, Ibeh GO and Ekeke GI. Proximate composition and levels of
320 some toxicants in four commonly consumed spices. Journal of Applied Sciences
321 and Environmental Management 2005; 9(1):150-155.
- 322 25 Obadoni BO and Ochuko PO. Phytochemical studies and comparative efficacy of
323 the crude extract of some homostatic plants in Edo and Delta States of Nigeria.
324 Global Journal of Pure Applied Sciences 2001;8: 203-208.
- 325 26 Sahoré DA and Amani NG. Classification of Some Wild Yam Species Tubers of
326 Ivory Coast Forest Zone. International Journal of Biochemistry Research & Review
327 2012; 2(4): 137-151

World J of
Microbiology & Biotech
Biotechnology

- 328 27 Ogbonna IO, Opobiya MY, Katuka B and Waba JT. Microbial Evaluation and
329 Proximate Composition of Kunu zaki, an Indigenous Fermented Food Drink
330 Consumed Predominantly in Northern Nigeria. Internet Journal of Food Safety
331 2011; 13: 93-97
- 332 28 Abiodun OA, Adegbite JA and Omolola AO. Chemical and Physico-chemical
333 properties of moringa flours and oil. Global Journal of Science Frontier Research:
334 Biological Sciences 2012; 12 (5): 12-17.
- 335 29 Anwar F, Zafar SN and Rashid U. Characterization of moringa seed oil from drought
336 and irrigated regions of Punjab. Grasasy Aceites 2006; 57 (2):160-168
337
- 338 30 Asuquo JE and Etim EE. Phytochemical and Antinutrients Evaluation of Oxyporus
339 Populinus. Journal of Emerging Trends in Engineering and Applied Sciences 2011;
340 2 (5): 817-820
- 341 31 Santos AFS, Luciana A, Adriana CCA, Teixeira JA, Paiva PMG, Coelho LCBB.
342 Isolation of a seed coagulant *Moringa oleifera* lectin. Process Biochemistry 2009; 44:
343 504-508
- 344 32 Ogbe AO and Affiku JP. Proximate study, mineral and anti-nutrient
345 composition of *Moringa oleifera* leaves harvested from Lafia, Nigeria: potential
346 benefits in poultry nutrition and health. Journal of Microbiology, Biotechnology and
347 Food Sciences, 2011/12; (3) 296-308
348
349
350
351
352
353
354
355
356
357
358
359

360
361
362
363
364
365
366
367
368
369
370