

PERFORMANCE AND BLOOD PROFILE OF WEST AFRICAN DWARF GOAT FED VARYING LEVELS OF CORN COBS IN COMPOUNDED RATION

Abstract

This study was conducted to investigate the effects of replacing brewer's dry grain (BDG) with corn cobs at varying proportions of 0% (control), 15%, 30% and 45% respectively, using guinea grass as basal feed on the performance and blood profile of West African Dwarf (WAD) goats. The study lasted for 84 days during which the performance and blood parameters were monitored in twelve WAD goats using a completely randomized design. Results obtained showed that bucks on diet 3 (30% CC inclusion); had the highest dry matter intake (559g/d), feed conversion ratio (7.89) and weight gain (7.08g/d). There was no significant difference ($P > 0.05$) in the serum biochemical parameters of WAD goats across all the dietary treatments except in globulin and creatinine where goats on diet 3 had the highest values ($P < 0.05$). No significant differences ($P > 0.05$) were also observed in the haematological parameters of WAD goats across all the dietary treatments. The study revealed that inclusion of corn cobs in the diets of WAD goats had no deleterious effects on the haematological and serum biochemical parameters of West African Dwarf goats and could therefore be included in ruminant diets up to 30%.

Keywords: Corncob, goats, heamatology, serum biochemical indices, performance

INTRODUCTION

Feed Scarcity especially during the dry season is one of the major constraints hindering the development of ruminant livestock production in Nigeria (Oladotun *et al.*, 2003). According to FAO, (2005) livestock production is below 5% while human population is increasing at rate above 10%. The major constraint to livestock production has been identified as nutrition, which is mainly occasioned by seasonal fluctuations in quantity and quality of available forage which is the major source of feed for ruminants. However, one possible way to mitigate this incessant challenge ravaging ruminant livestock production is to explore alternative feed resource that are rarely consumed by man but which can meet the nutritional requirement of goats with little or no processing. This will reduce the cost of animal production without a decrease in productivity.

In Nigeria, there is paucity of information on utilization of crops residues, agro-industrial by-products and non-conventional feed resources for livestock nutrition. In the classification of feed resources; soybean milk residue, cowpea seed waste and corn cob have been regarded as non-conventional feed resources (NCFR). These residues or wastes have considerable potential as feed materials and their value can be increased if they are economically justifiable (Devendra, 1991).

Corncocks are by-products of maize production and may be used as alternative feed resources for ruminants especially during dry season. Corncocks are important crop residue used by livestock farmers to supplement fodder during the dry season (Olorunju *et al.*, 1996). They are relatively available and abundant in the dry season; a major limiting factor in the utilization of this agricultural waste is its low digestibility and relatively poor nutrient composition (Kategile, 1981). However, ruminants generally had been known to be the best in the utilization of coarse materials such as corn cobs for the production of meat and other products, thereby preferring feedstuffs relatively rich in crude fibre (Williamson and Payne 1987).

Haematological and biochemical indices of animals may give some insight into the production performance potentials of West African Dwarf goats (Orheruata and Akhuomobhogbe 2006). Nutrition, Breed, Sex, Age, Reproductive status, Environmental factors, stress and Transportation are known to affect haematological and biochemical parameters (Balikel *et al.*, 2007). The present study was designed to determine the performance and blood profile of West African Dwarf Goats fed compounded ration consisting of varying levels of corncob as replacement for brewers' dry grain.

Materials and methods

Location and climate of the study area

The experiment was conducted at the Ruminant Unit of the Institute of Agricultural Research and Training, Moor plantation Ibadan, Nigeria. The experimental site was located in the derived savannah vegetation zone of south-western Nigeria. The area lies within the rainforest ecological zone, the temperature and relative humidity ranges from 30-35°C and 76-84% respectively.

Experimental animals and their management



Twelve West African Dwarf (WAD) Goats weighing between 6.5 – 8.5kg were used for this study. The goats were housed intensively in well-ventilated individual pens disinfected with morigad[®] solution two weeks prior to the experiment. On arrival, the animals were kept in the pens for proper routine management. All the goats were given antibiotic injection (Oxytetracycline L/A) while Ivomec injection was administered to control both the endo and ecto-parasites. The goats were vaccinated against *Peste des Petit Ruminant* (PPR). Concentrate feed and Guinea grass was fed to the goats during the 14 days adaptation period. Feed and fresh clean water were supplied *ad-libitum*.

Experimental diets

Wilted and chopped Guinea grass (*Panicum maximum*) was fed as basal diet to the goats. Four experimental concentrate diets were compounded to include corn cob at 0%, 15%, 30%, and 45%. Other ingredients in the diets were brewers dried grain, wheat offal, palm kernel cake, groundnut cake, limestone, salt and premix (Table 1). After two weeks of adaptation period, the animals were divided into four treatment groups of three animals each and thereafter assigned to one of the four different experimental diets formulated in a completely randomized design. Each animal was kept and fed separately during the entire period of the experiment. The concentrate supplements were given to the animals at morning while wilted and chopped panicum maximum that had been harvested the previous day was given to the animals in the evening. During the 84-day experimental period, quantities of feeds offered and that refused were measured daily to compute feed intake while weight changes of the goats were recorded weekly.

79

Table 1: Gross composition of experimental diet containing varying levels of corncobs

Ingredient (Kg)	Diet 1 (control)	Diet 2	Diet 3	Diet 4
Brewers dry grain	60	45	30	15
Corn cob	-	15	30	45
Wheat offal	10	10	10	10
Palm kernel cake	20	20	20	20
Groundnut cake	5	5	5	5
Limestone	4.5	4.5	4.5	4.5
Salt	0.25	0.25	0.25	0.25
Grower premix	0.25	0.25	0.25	0.25
Calculated Analysis				
Crude protein (%)	18.35	16.18	14.00	11.83
Metabolizable Energy (Kcal/Kg DM)	1942	1990	2038	2086

Collection of Blood Samples

Blood samples (approximately 10 ml) were collected from each goat via jugular vein puncture using hypodermic syringes before feeding. Blood collection was at the end of the feeding trial. Blood, 5 ml was drawn into a heparinized bottle to prevent coagulation while the remaining 5 ml was left in the syringe to coagulate. Blood samples were then analysed for packed cell volume (PCV), Hb, white blood cells (WBC), red blood cells (RBC), serum protein, ~~serum~~ glucose, ~~serum~~ albumin,  globulin, ~~serum~~ creatinine, ~~serum~~ urea, mean corpuscular haemoglobin (MCH), and mean corpuscular haemoglobin concentration (MCHC). The packed cell volume was measured for each animal in fresh ethyl diamine tetra acetic acid (EDTA) anticoagulant samples within 24 hours of collection using the micro- haematocrit method. Haemoglobin concentration was measured in fresh EDTA anticoagulant samples using sahl's (acid haematin) method (Benjamin 1978). RBC was measured in fresh EDTA with the aid of Neubauer counting chamber (haemocytometer). Blood smear were used for total thrombocytes, total WBC counts (Tavares-Dias *et al*, 2008) and WBC differential relative and absolute counts. Differential relative and absolute counts were classified as Lymphocyte, Neutrophils, Eosinophils, Basophils

and Monocytes. Plasma glucose was measured in fluoride oxalate anticoagulant blood samples using the enzymatic glucose oxidase method (Bauer *et al.*, 1974). MCH and MCHC values were calculated from PCV, Hb and RBC values (Jain, 1993). Total serum protein was measured in serum for individual animal using the biuret method. Serum Alanine aminotransferase and Serum Aspartate aminotransferase were analyzed spectrophotometrically by using commercially available diagnostic kits (Randox Test Kits). Serum albumin and globulin were measured using bromocresol purple method of Varly *et al.* (1980). Serum creatinine was determined using the principle of Jaffe reaction as described by Bousnes and Taussky (1945).

Chemical Analysis

Aliquot of daily feed samples (concentration and forage) was collected, oven-dried, ground and sieved through a 2-mm sieve and stored in airtight containers for proximate (AOAC 2000, reference ID Number DM 930.15, CP 984.13) and fibre (Van Soest *et al.*, 1991) analyses.



Statistical Analysis

Data obtained were subjected to one way analysis of variance (completely randomized design) and significant means were separated using Duncan Multiple Range test (SAS, 1998).

Results and Discussion

Table 2 shows the chemical composition of the diet, crude protein, ether extract and ash concentration decreased as the proportion of CC increased in the diet compounded whereas the NDF, ADF and ADL increased. The crude protein of grass was 8.4% with the lowest value of ether extract. Corn cob has the lowest crude protein (2.8%) and the highest EE and NDF which were 4.2% and 63.3% respectively. The performance characteristics of the goats fed the experimental diets differed significantly ($P < 0.05$) among the treatments (Table 3). The highest dry matter intake was observed in goats fed diet 3 with value of 559g/day and was followed by that of goats on diet 1 (540g/day); values observed for goats diet 2 and 4 were similar ($P > 0.05$). Similarly, goats on diet 3 had the highest crude protein intake (16.4g/KgW^{0.75}/day) while goats on diet 4 had the least CPI (11.2 g/KgW^{0.75}/day), the best feed conversion ratio (7.89) was also

observed in goats on diet 3 and this culminated to the highest weight gain (70.83g/day) observed when compared to goats on diet 1, 2, and 4. The grinding of the corn cob to reduce the particle size increased the surface area for rumen microbial action. Factors affecting feed intake include dietary crude protein, palatability, gut fill, rumen outflow rate/retention time in the rumen (Ahamefule and Elendu (2010). Highest weight gain of 70.83g/day is higher than reported value for goats fed cassava leaf meal in corn bran based diet (Yousuf *et al.*, (2007). Table 4 shows the Serum Biochemical parameters of goats fed diets containing varying levels of corn-cob. Similar ($P>0.05$) Total protein, globulin, urea and glucose values were obtained for all the goats in the four treatment groups. However, highest ($P<0.05$) and least creatinine levels of 16.00 and 7.00mg/dl were obtained in goats fed diets containing 30% corn-cob and the control diet respectively. Goats fed the control diet contained the highest ($P<0.05$) cholesterol level (85.86mg/dl). However, goats fed diet containing 45% corn-cob had the least ($P<0.05$) cholesterol level which did not differ significantly ($P>0.05$) from the cholesterol content in goats fed diets containing 15% and 30% corn-cob based diet. The lower values for the total protein, albumin and globulin in this study indicated that there were no traces of antinutritional factors that could diminish nutrient permeability in the gut walls. Serum urea levels reported in this study were high compared to values reported for apparently healthy Marwari goats (Tanwar *et al.*, 2000). This may probably have been due to persistent hypoglycemia and catabolic activity is increased for gluconeogenesis thus resulting in high serum urea levels.



The Haematological parameters of West African Dwarf goats fed diets containing varying levels of corn-cobs showed that there was no significant ($P>0.05$) differences in all the parameters measured Table 5. The PCV values obtained in this study were within the range of 25-30% reported by Opara *et al.*(2010) for clinically healthy WAD goats and sheep. The Hb range in this study fell within the range of 7-15g/dl reported by Daramola *et al.*(2005) but higher than the values of 5-6g/dl obtained by Balewu and Ogunsola (2010) for goats fed fungi treated *Jatropha curcas* kernel cake rations. Due to relatively higher Hb concentration obtained in this study, the dietary treatments seemed to have the wherewithal of supporting high oxygen carrying capacity blood in the goats. The RBC counts obtained in this study were within the range of 9.2-13.5 $\times 10^6/\mu\text{l}$ reported for West African dwarf goats by Tambuwal *et al.* (2002), 9.9-18.7 $\times 10^6/\mu\text{l}$ by Taiwo and Ogunsanmi (2003). The WBC count obtained in this study fell within the range of 6.8-20.1 $\times 10^6/\mu\text{l}$ reported by Daramola *et al.*(2005) The experimental animals seem to possess a

protective system capable of preventing them from infection and this is probably the physiological basis for the adaptation of this species in there ecological zone. Daramola *et al.* (2005).

The lymphocytes (%) values obtained in this study fell within the range of 47- 82% and 51.6% reported by Daramola *et al.* (2005) and Tambuwal *et al.* (2002) respectively. The results indicated there was fluctuation in the values of lymphocyte obtained across the dietary treatment and this may be attributed to physiological stress which is accompanied by increase lymphopenia or lymphocytopenia.

Conclusion

It is concluded from this study that corncobs can be included in the diets of goats to replace brewers dried grain up to 30% without deleterious effect on the performance, haematology and serum biochemical indices of the goat. The nutrient intake, feed conversion ration and weight gain observed in goats on 30% inclusion showed that corncob in compounded ration can support optimal performance of goats.

References

- AOAC, 2000. Official methods of analysis. Association of official Analytical chemists (17th edition). Arlington.
- Ahamefule, F.O. and Elendu, C. 2010: Intake and digestibility of West African Dwarf bucks fed cassava leaf maize offal based diets, *J. Anim, Vet.* 9: 535-539.
-  Balikei, E., Yildiz, A. and Gurdogan, F., 2007. Blood metabolite concentrations during pregnancy and post-partum in Akkaraman ewes. *Small Ruminant Research*, 67, 247-251
- Bauer, T.D., Ackermann, P.G and Toro, G., 1974. Methods in clinical chemistry. Clinical laboratory methods. The C.V. Mosley company, Saint Louis, Pp. 946.

189 Belewu, M. A and Ogunsola, F.O., 2010. Haematological and serum indice of goat fed fungi
190 treated *Jatropha curcas* kerner cake in a mixed ration. *Journal of Agriculture*
191 *Biotechnology and Sustainability Development*, 2(3), 035-038

192 Benjamin, M.M., 1978. *Outline of Veterinary clinical pathology*, 2nd edition. IOWA State
193 University Press. IOWA, U.S.A. Pp. 35-105

194
195 Bonsnes, R.W. and H.H.J. Taussky. 1945. Determination of creatinine in plasma and urine. *J.*
196 *Biochem.* 58: 581–589.

197 Daramola, J.O., Adeloye, A.A., Fatoba, T. A., Soladoye, A.O., 2005. Haematological and
198 biochemical parameters of West African Dwarf goats. *Livestock Res. Rural Dev*: 17.
199 90-114.

200 Devendra, C. 1991. *Feed Resources in Goat Husbandry and Breeding in the Tropics*.
201 Compiled paper presented in an International seminar carried out by German
202 Foundation for international development (DSF) at the Institution for Advanced
203 Studies, University of Malayan. Kwala Lumpur) Pp 121-135

204 **FAO**, 2005. FAOSTAT data 2005, <http://faostat.fao.org/faostat>.

205 Jain N.C., 1993. *Essentials of Veterinary Haematology*. Lea and ferbeiger, Pennsylvania, U.S.A.
206 Pp. 7.

207 Kategile, J.A. 1981. Digestibility of low quality roughages supplemented with concentrate.
208 *Proc. Of Workshop, held in Arusha, Tanzania, AUN Agricultural Development Report*
209 *Norway, AUS. NLH. 18 – 22 January, 1981. Pp 181 - 184*

210 Oladotun, O.A., Aina, A.B.J. and Oguntona, E. B. 2003. Evaluation of formulated agro-
211 industrial waste as dry season feed for sheep. *Nig. Journal. Animal Production.*, 30:71-80.

212 Olorunju, P. E., Ntare, B. R., Babalola, O.A and Alabi, O. 1996. In proceeding of a
213 workshop on nationally co-ordinated Groundnut Research Project. 25-29
214 September, 1995, Zaria, Nigeria, IAR/ABU, Zaria and ICRISAT, Kano, Nigeria, pp:47.

215 *Proc. Of Workshop, held in Arusha, Tanzania, AUN Agricultural Development*
216 *Report Norway, AUS. NLH. 18 – 22 January, 1981. Pp 181 - 184*

217 Opara, M.N., Udevi, N. and Okoli, I.C., 2010. Haematological Parameters and Blood Chemistry of
218 Apparent Healthy West African Dwarf (WAD) goat in Owerri, South Eastern Nigeria.
219 *New York Science Journal*, 3 (8), 68-72.

220 SAS. 1998. **Statistical Analytical Systems**. SAS/STAT User's Guide Statistical Analysis
 221 Institute Inc. Version 6, 3rd ed. Cary, North Carolina, USA. 943 pp.
 222

223 Taiwo, V.O. and Ogunsanmi, A.O. 2003. Haematology, plasma, whole blood and erythrocyte
 224 biochemical values of clinically healthy captive-reared grey duiker (*Sylvicapra grimmia*)
 225 and West African dwarf sheep and goats in Ibadan, Nigeria. Isreal J. Vet. Med. Retrieved
 226 from http://www.isrvma.Org/article/58_2_3.htm

227 Tambuwal, F.M., Agale,B.M. and Bangana, A., 2002. Haematological and biochemical values of
 228 apparently healthy Red Sokoto goats. In: Proceedings of the 27th Annual conference of
 229 the Nigeria Society for Animal Production (NSAP), held between 17-21 March 2002,
 230 Federal University of Technology, Akure , Nigeria. 50-53

231 Tanwar , R.K.,Tinna, N.K,Gahlot, A.K. and Sharma, S.N.,2000. Biochemical profile of clinical
 232 ketosis in goats . In : Proceedings of the 7th international Goat Association
 233 (IGA,)Toulouse, France, 306-307

234 Tavares Dias, M., Oliveira-Junior, A., Marcon, J.L., 2008.Methodological Limitations of
 235 counting total Leukocytes and thrombocytes in reptiles (Amazon turtle, *Podocnemis*
 236 *expansa*): an analysis and discussion. Acta Amazon, 38, 159-163.

237 Varly, H.,Van,E.andKass,I.,1980. Practical clinical Chemistry, New York, Inter Science
 238 Publishers Inc. 197-240
 239

240 Van Soest, P.J., J.B. Robertson and B.A. Lewis.1991. Methods for dietary fiber, neutral
 241 detergent fi ber, and nonstarch polysaccharides in relation to animal Nutrition. J. Dairy
 242 Sci. 74: 3583–3597.

243 Williamson, G And Payne, W.J.A.,1987. An introduction to animal husbandry in tropics, 3rd
 244 Edn., ELBS series. Longman, London, pp: 463 – 484.

245 Yousuf, M.B., Belewu, M.A., Daramola, J.O. and Ogundun, N.I. 2007. Replacing Cotton Seed
 246 Cake with Cassava Leaf Meal in Corn Bran Based Diet Fed to the Goat. Bulgarian.
 247 *Journal of Agricultural Science*, 13, 231 – 236.
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257

258
259
260

Table 2: Chemical composition of the diets containing varying levels of corn cobs

Parameters (%)	Diet 1	Diet 2	Diet 3	Diet 4	Guinea	Corn cob
	control	(15%)	(30%)	(45%)	Grass	
Dry matter	90.1	90.2	90.3	90.1	36.1	88.4
Crude protein	17.2	16.7	16.3	15.8	8.4	2.8
Ether extract	3.66	3.62	3.58	3.54	2.44	4.2
Ash	9.9	8.8	9.6	9.0	9.9	3.8
NDF	51.4	53.7	52.8	55.2	61.4	63.3
ADF	42.4	44.2	43.2	46.3	46.8	30.2
ADL	6.8	7.8	6.9	7.0	9.7	9.5

261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276

277

278

279

280

281

282

283

284

Table 3: Performance characteristics of goats fed diet containing corncob in varying proportions

Parameters	Diet 1	Diet 2	Diet 3	Diet 4	SEM
	(0%)	(15%)	(30%)	(45%)	
Dry Matter (g/day)					
Grass	350	351	362	349	
Concentrate	190	182	197	180	
Total	540 ^b	533 ^c	559 ^a	529 ^c	5.44
Nutrient intake (g/KgW ^{0.75} /day)					
Dry matter	99.1	96.0	97.4	97.1	
Crude protein	15.1	15.7	16.4	11.2	
NDF intake	61.5	57.7	55.5	56.0	
ADF intake	43.4	45.2	44.4	47.4	
ADL intake	6.6	8.7	8.4	9.2	
Feed conversion ratio	10.19 ^a	9.18 ^a	7.89 ^b	10.79 ^a	1.24
Initial weight (Kg)	7.43	7.40	7.37	7.53	
Final weight (Kg)	11.88	12.28	13.32	11.65	
Live weight (g/Kg W ^{0.75} /day)	5.45	5.56	5.77	5.45	0.25
Weight gain (g/day)	52.98 ^b	58.09 ^b	70.83 ^a	49.04 ^b	5.12

abc=Mean in the in the row with similar superscript are not significantly different (P>0.05)

NDF= Neutral detergent Fiber; ADF= Acid detergent fiber; ADL= Acid detergent lignin

Table 4: Serum Biochemical Parameters of West African Dwarf Goats Fed Diets Containing Varying Levels of Corncobs

Parameters	Diet 1 (0%)	Diet 2 (15%)	Diet 3 (30%)	Diet 4 (45%)	SEM
Total protein (g/dl)	5.08	5.40	5.33	5.11	0.13
Albumin (g/dl)	3.87 ^{ab}	3.80 ^{ab}	4.52 ^a	3.40 ^b	0.17
Globulin (g/dl)	1.21	1.61	0.81	1.72	0.20
Urea (mg/dl)	28.20	31.29	33.07	27.31	1.04
Creatinine (mg/dl)	7.00 ^b	11.13 ^{ab}	16.00 ^a	11.00 ^{ab}	1.38
Glucose (mg/dl)	71.45	57.63	71.29	65.87	3.48
Cholesterol (mg/dl)	85.86 ^a	49.50 ^{ab}	68.94 ^{ab}	36.61 ^b	7.93

^{ab} Means on the same row with different superscripts differ significantly(P<0.05).

SEM - Standard Error of Mean

Table 5: Haematological Parameters of West African Dwarf Goats Fed Diet Containing Varying Levels of Corncobs

Parameters	D ₁	D ₂	D ₃	D ₄	±SEM
	(0%)	(15%)	(30%)	(45%)	
Packed cell volume (%)	24.00	22.50	24.00	20.00	0.80
Haemoglobin (g/dl)	10.10	9.10	9.20	9.85	0.42
Red blood cell (x10 ⁶ /μl)	11.25	10.58	10.59	10.92	0.35
White blood cell (x10 ³ /μl)	17.03	13.33	17.20	12.55	19.76
Lymphocyte (%)	50.50	55.00	50.50	54.00	1.69
Heterophils (%)	45.50	42.50	46.00	43.00	1.76
Monocytes (%)	2.50	2.50	2.50	2.00	0.18
Eosinophils (%)	1.50	1.00	1.00	1.00	0.35
Platelets (x10 ⁴ mm ³)	20.40	19.50	16.10	15.30	1.89
MCV (fl)	21.54	21.37	22.71	18.28	0.81
MCH (pg)	0.90	0.86	0.87	0.90	0.01
MCHC (g/dl)	4.22	4.04	3.85	4.95	0.22

MCV= Mean Corpuscular Volume; MCH= Mean Corpuscular Haemoglobin
MCHC = Mean corpuscular Haemoglobin Concentration; SEM = Standard Error of Mean