

Original Research Article**Direct Seeded and Transplanted Maize: Effects of Planting Date and Age of Seedling on the Yield and Yield Attributes****ABSTRACT**

An experiment was conducted at the Bangladesh Agricultural Research Institute, Regional Station, Jamalpur during rabi 2003-04 to study the performance of maize varieties at different planting dates and ages of seedling. The experimental site is located at 24°56'11" N latitude and 89°55'54" E longitude and at an altitude of 16.46 m. The duration of the study period was November 2003 to May 2004. Design of the experiment was split-split plot having 3 (three) replications assigning planting date in the main plot, variety in the sub plot and age of seedling in the sub-sub plot. Five planting dates (20 November 30 November, 10 December 20 December 30 December), two maize varieties (BARI maize-6 and Pacific-11) and three ages of seedlings (Direct planting of seed, 14-day old and 21-day old seedlings) were included as treatments in the experiment. Seedlings were raised in the dry bed underneath polythene sheet nursery. The results revealed the variety Pacific-11 produced significantly higher grain yield in the earlier November 20 and 30 planting irrespective of direct seeding and age of seedling. There were no significant difference in respect of grain yield in the earlier November 20 and November 30 plantings with all seedling ages but planting 14-day old seedling the crop may be harvested about 7 days earlier while planting 21-day old seedling crop may be harvested about 12 days-earlier than direct seeding. Transplanting of 14-day old seedling at 10, 20 and 30 December may produce higher grain yield compared to both direct seeding and 21-day old seedling along with 5-6 days reduced field duration than direct seeding. This early harvest may save the crop from natural vulnerability of hailstorm and rains, and may help in early establishment of the next crop like sesame, aus rice, mungbean blackgram, jute etc.

Key words: Planting date, age of seedling, yield and yield attributes, maize

1. INTRODUCTION

Maize (*Zea mays* L.) is the top most ranking cereal in terms of higher grain yield and holds third position in respect of total production following wheat and rice in the world. Maize is nutritionally superior to most other cereals as it contains 9.0% protein, 3.4% fat, 1.1% ash, 1.0% starch fibre, 0.30% thiamine, 0.08% riboflavin and 1.9% niacin (Paliwal, 2000). In Bangladesh, maize is primarily

used as feed for livestock and poultry, and a considerable proportion of grains and green cobs is used for human consumption. The demand of maize is thus rapidly increasing. It is expected to be increased further with the establishment of maize-based food industries, and poultry, dairy and fish farms. This may lead to an enormous increase in maize import, which might result in depleting the hard-earned foreign currency. Hence, special emphasis should be given for increasing the indigenous maize production in order to fulfill the present demand as well as of the future. The agro-climatic conditions of Bangladesh are favourable for maize cultivation round the year. However, *rabi* season is reported to be the best time for maize cultivation with the optimum time of planting ranging from mid November to end November (BARI, 1997). Maize is generally grown after harvest of transplant aman rice and consequently, planting gets delayed. In the *char* lands, the potential maize growing areas of Bangladesh, the lands attain 'joe' condition in late after recession of flood water and thus, maize planting goes beyond the optimum time. Mohamed and Shams (1991) reported that grain yield of hybrid maize was affected less by late sowing than the synthetic and open pollinated varieties. Porwal and Jain (1999) stated that grain yield of maize reduced due to early and delayed plantings to mid November by 3.9 to 66.4%. Hence, planting time and choice of variety should be important considerations for maize cultivation. However, in addition to reduced grain yield the late sown crops may experience natural hazards at the later stage of growth due to nor'wester. Again the sowing of following crops like jute, aus rice, mungbean, blackgram etc. may be affected by the late maize harvest. So, the development of an appropriate early plant establishment technique through agronomic management is essential for maize cultivation under these situations. Basu *et al.* (2003) opined that transplanting of 21-day old seedlings gave identical grain yields with direct sown crop and matured 8-10 days earlier. Maize crop could also be established much earlier through transplanting technique which may produce higher yield (Dale and Drennan, 1997b). Badran (2001) stated that under late planting conditions, transplanting of maize may be a possible alternative to direct sowing. Therefore, transplanting of seedling may be an important area of study for maize cultivation considering the field duration and early plant establishment under Bangladesh conditions. On the basis of the discussion made above, the present research work was undertaken to evaluate the performance of transplanting techniques including varying seedling ages of maize against direct planting of seed under different dates.

60 2. Materials and Methods

61 The experiment was carried out to find out suitable planting material either seed or seedling in
 62 different planting time for producing higher grain yield of maize. The treatments of the experiments
 63 were five planting dates (20 November, 30 November, 10 December, 20 December, 30 December),
 64 two varieties (BARI maize-6 and Pacific-11) and three ages of seedlings (Direct planting of seed, 14-
 65 day old and 21-day old seedlings). Seedlings were raised in the dry bed underneath polythene sheet
 66 nursery. Seeds were planted earlier in the bed to attain 14- and 21-day old seedlings at different
 67 dates. For example, to attain 14- and 21-day old seedlings seeds were planted in the nursery on 6
 68 November and 31 October 2003, respectively for 20 November planting. A split-split plot design
 69 arranging planting date (Factor A) in the main plot, variety (factor B) in the sub plot and age of
 70 seedling (factor C) in the sub-sub plot was used to carry out the experiment. Twenty plants were
 71 randomly selected to collect data on length of cob, cob diameter, number of grain cob⁻¹, 1000 grain
 72 weight, grain yield and harvest index. Harvest area 10.5 m² was used for calculation of grain yield in t
 73 ha⁻¹. Seedling raising method, fertilizer application rate and methods, and all intercultural operations
 74 were described by Biswas *et al.* (2009). Data were analysed with the help of a computer package
 75 program MSTATC and means were separated as per DMRT at 5% level of significance.

76 3. Results and Discussion

77 3.1 Effects of planting date

78 3.1.1 Yield and yield attributes

79 Planting date did not exhibit significant effect on number of grain cob⁻¹ but on plant height at harvest,
 80 length of cob, cob diameter, number of cobs plant⁻¹, number of grain rows cob⁻¹, number of grains
 81 row⁻¹ and 1000 grain weight (Table 1). The tallest plant was found in 10 December planting while the
 82 shortest plant was found in 20 November planting irrespective of variety and age of seedling. The
 83 results did not agree with Prodhan (2001) who found the tallest plant in November than December
 84 sowing. This might be due to variation in the soil and climatic factors as the location was different.
 85 Length of cob was recorded the highest in 20 November planting. November 30 and 10 December
 86 also gave statistically similar length of cob to 20 November planting while the lowest length of cob was
 87 found in 30 December planting. There were no significant differences up to 20 December in terms of
 88 cob diameter having the in 20 November planting. December 30 planting had the lowest cob
 89 diameter. Number of cobs plant⁻¹ and number of grain rows cob⁻¹ were also significantly highest in 20

November planting. December 10 planting produced statistically similar number of grain row cob⁻¹ to the 20 November. Number of cob plant⁻¹ decreased progressively with delayed plantings while number of grain row⁻¹ increased in the delayed plantings. But 1000-grain weight was found to reduce as the plantings were delayed. Grain yield was mainly ascribed to higher number of cobs plant⁻¹, number of grains cob⁻¹ and 1000 grain weight. Hence, the earlier two plantings viz. 20 November produced the highest grain yield that was closely followed by 30 November planting. Grain yield reduced significantly beyond 30 November planting and the lowest was obtained from the last planting of 30 December due to its lowest number of cobs plant⁻¹, and 1000 grain weight mainly. Sandhu and Hundal (1991) stated that grain yields decreased in later sowing were mainly associated with lower 100 grain weight. The highest stover yield was obtained in 30 November planting which was statistically similar to 20 November. Stover yield reduced significantly beyond 30 November planting and the lowest stover yield was obtained from 30 December planting. The early planting also gave the highest harvest index and beyond 10 December planting, harvest index reduced significantly compared to 20 November planting.

Table 1. Yield and yield attributes of maize as influenced by planting date during *rabi* 2003-04

Planting date	Plant height at harvest (cm)	Length of cob (cm)	Cob diameter (cm)	Number of cobs m ⁻²	Number of grains cob ⁻¹	1000-grain weight (g)	Grain yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)	Harvest index (%)	Field duration (days)
20 November	191.1 b	17.5 a	5.2 a	7.04 a	407.2	370.2 a	10.49 a	8.88 ab	54.26 a	139
30 November	205.9 ab	17.0 a	5.0 a	6.71 b	412.9	373.68 a	10.36 a	9.06 a	53.18 ab	139
10 December	214.7 a	16.7 ab	5.0 a	6.39 c	426.2	352.5 b	9.57 b	8.62 b	52.59 bc	136
20 December	203.8 ab	16.0 bc	5.0 a	6.06 d	423.9	340.7 c	8.76 c	7.93 c	52.33 bc	132
30 December	194.3 b	15.8 c	4.9 b	5.66 e	418.7	339.4 c	8.00 d	7.42 d	51.83 c	129
Level of significance	*	**	*	**	NS	**	**	**	**	-
CV(%)	10.03	6.29	2.62	5.46	5.24	2.66	2.20	5.93	2.69	-

Figures in a column having no or similar letter(s) do not differ significantly while those with different letter(s) differ significantly at 5% level of probability; * = Significant at p≤0.5; ** = Significant at p≤0.01; NS = Not significant at 5% level.

108 3.2 Effects of variety

109 3.2.1 Yield and yield attributes

110 Yield and all the yield components except cob diameter and number of grain row⁻¹ differed
111 significantly between the variety (Table 2). The results revealed that Pacific-11 had the higher values
112 than BARI maize-6 in respect of all the yield components which in turn produced the highest grain
113 yield (Table 2). Grain yield of Pacific-11 was about 24.11% higher than BARI maize-6. Pacific-11 also
114 produced the highest stover yield and harvest index. Anil and Sezer (1995) observed that there were
115 significant differences between the cultivars in terms of plant height, number of grain, ear weight etc.
116 of sweet corn transplant.

117
118 **Table 2. Yield and yield attributes maize as influenced by the variety during *rabi* 2003-04**

Treatment	Plant height at harvest (cm)	Length of cob (cm)	Cob diameter (cm)	Number of cobs m ⁻²	Number of grains cob ⁻¹	1000-grain weight (g)	Grain yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)	Harvest index (%)	Field duration (days)
Variety										
BARI maize-6	193.2 b	16.0 b	5.0	5.92 b	408.2 b	341.1 b	8.15 b	7.49 b	52.10 b	134
Pacific-11	210.8 a	17.3 a	5.0	6.83 a	427.4 a	369.4 a	10.72 a	9.27 a	53.57 a	135
Level of significance	**	**	NS	**	**	**	**	**	**	-
CV(%)	7.06	4.50	2.37	5.41	6.62	2.82	2.44	3.85	2.20	-

119 120 3.3 Effects of age of seedling

121 3.3.1 Yield and yield attributes

122 All the characters except number of grain row⁻¹ showed significant variation among the seedling ages
123 (Table 3). Direct planting had the highest values in respect of all yield contributing characters except
124 1000 grain weight. Direct planting produced the longest plant and it reduced gradually and the
125 shortest plant was found in 21-day old seedling. Thousand grain weight increased significantly in the
126 older age of seedling. The lowest 1000 grain weight was obtained from direct seeded while the
127 highest was obtained from the older age of seedling (21-day). The highest grain yield was found in
128 14-day old seedling while lowest was found in 21-day old seedling. Grain yield in 14-day seedling was
129 possibly compensated by the heavier grain size in spite of its lower number of cobs plant⁻¹ and
130 number of grains cob⁻¹ than direct planting. Direct planting produced the second highest grain yield

131 which was statistically similar to 14-day old seedling. Stover yield was decreased progressively with
132 the increase of seedling age while HI increased with the increase of seedling age.

133 **Table 3. Yield and yield attributes of maize as influenced by age of seedling during *rabi* 2003-**

134 **04**

Age of seedling (days)	Plant height at harvest (cm)	Length of cob (cm)	Cob diameter (cm)	Number of cobs m ⁻²	Number of grains cob ⁻¹	1000-grain weight (g)	Grain yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)	Harvest index (%)	Field duration (days)
0	224.1 a	17.1 a	5.0 a	6.58 a	430.4 a	337.5 c	9.47 a	9.23 a	50.43 c	141
14	204.2 b	16.7 a	5.0 a	6.42 b	423.3 a	355.1 b	9.59 a	8.35 b	53.28 b	135
21	177.6 c	16.1 b	4.9 b	6.11 c	399.8 b	373.2 a	9.25 b	7.56 c	54.79 a	129
Level of significance	**	**	**	**	**	**	*	**	**	-
CV(%)	6.10	4.37	2.05	4.72	6.09	2.51	2.54	5.19	2.34	-

135 Note: '0' = Direct seeded

136 There were no significant differences for the interaction of variety and age of seedling, and interaction
137 of planting date, variety and age of seedling in terms of yield and yield attributes. Hence, the results of
138 these interactions were not presented here. Interaction of variety and planting date, and planting date
139 and age of seedling were found significant and the results have been described as follows.

140

141 3.4 Interaction effects of variety and planting date

142 3.4.1 Yield and yield attributes

143 There were significant differences in respect of number of cobs plant⁻¹, grain yield, stover yield and
144 harvest index but other parameters did not differ significantly (Table 4) due to interaction effect of
145 planting date and variety. The results exhibited that the highest number of cobs was produced by the
146 combination V₂ x D₁ and the lowest number was recorded from V₁ x D₅ combination. However, the
147 highest grain yield was obtained from the combination V₂ x D₁ which was closely followed by the
148 combination V₂ x D₂. The lowest grain yield was recorded from the combination V₁ x D₅. Grain yield
149 was decreased significantly in the later plantings beyond November 30 in both the varieties. Hybrid
150 maize was affected less and yielded higher than the synthetic and open pollinated varieties by late
151 sowing (Mohamed and Shams, 1991). The highest stover yield was obtained from the combination V₂
152 x D₁ combination and it was significantly different from the others. The combination V₂ x D₂ gave the
153 highest harvest index which was statistically similar to the combinations of V₁ x D₁, V₂ x D₁, and V₂ x

D₄ (Table 4). The results revealed that Pacific-11 yielded significantly higher than BARI maize-6. As the interaction effect of variety and planting date was significant it was found that grain yield also reduced beyond 30 November planting and the lowest grain yield was obtained from 30 December in both the varieties irrespective of age of seedling. It seems to be number of cobs m⁻² was the main determinant for grain yield. Although no significant difference was observed for number of grains cob⁻¹ but an increasing trend was observed with late planting while a reverse result was observed for 1000 grain weight in both the varieties. Grain yield decreased with delayed plantings at or after 10 December and 30 December produced the lowest grain yield for both the varieties Pacific-11 and BARI maize-6.

163

Table 4. Yield and yield attributes of maize as influenced by the effects of interaction of variety and planting date during *rabi* 2003-04

Interaction	Plant height at harvest (cm)	Length of cob (cm)	Cob diameter (cm)	Number of cobs m ⁻²	Number of grains cob ⁻¹	1000-grain weight (g)	Grain yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)	Harvest index (%)	Field duration (days)
V ₁ × D ₁	181.1	17.1	5.1	6.40 c	398.0	353.8	8.92 d	7.48 d	54.48 a	139
V ₁ × D ₂	196.3	16.2	5.1	6.33 c	396.2	357.7	8.90 d	8.48 c	51.21 cd	138
V ₁ × D ₃	206.0	16.0	5.0	5.82 d	426.6	335.4	8.22 e	7.38 d	52.59 b	135
V ₁ × D ₄	196.4	15.5	5.0	5.60 de	419.9	329.3	7.69 f	7.51d	50.65 d	130
V ₁ × D ₅	186.1	15.1	4.8	5.43 e	400.5	329.3	7.01 g	6.58 e	51.56 b-d	130
V ₂ × D ₁	201.1	17.8	5.1	7.68 a	416.4	386.5	12.06 a	10.27 a	54.03 a	139
V ₂ × D ₂	215.5	17.8	5.0	7.09 b	429.7	389.5	11.83 a	9.64 b	55.14 a	139
V ₂ × D ₃	223.4	17.5	5.0	6.95 b	425.9	369.6	10.92 b	9.87 b	52.58 b	136
V ₂ × D ₄	211.3	16.6	5.0	6.52 c	427.9	352.0	9.83 c	8.35 c	54.00 a	133
V ₂ × D ₅	202.6	16.6	4.9	5.90 d	437.0	349.5	8.99 d	8.26 c	52.11 bc	129
Level of significance	NS	NS	NS	*	NS	NS	**	**	**	-
CV(%)	7.06	4.50	2.37	5.41	6.62	2.82	2.44	3.85	2.20	-

Figures in a column having no or similar letter(s) do not differ significantly while those with different letter(s) differ significantly at 5% level of probability; * = Significant at p≤0.05; ** = Significant at p≤0.01; NS = Not significant at 5% level; V₁=BARI maize-6, V₂= Pacific-11, D₁=20 November, D₂= 30 November, D₃= 10 December, D₄= 20 December, D₅= 30 December.

3.5 Interaction effects of planting date and age of seedling

3.5.1 Yield and yield attributes

None of the characters except 1000 grain weight, grain yield, stover yield and harvest index differed significantly due to interaction effects of planting date and age of seedling (Table 5). The results revealed that in the early plantings of 20 and 30 November, direct planting had the highest grain yield but thereafter 14-day old seedling produced the highest. Twenty-one-day old seedling produced statistically similar grain yield to the direct planting at all planting dates. The results clearly indicated that under late planting conditions 14-day old seedling may be transplanted to obtain higher grain yield. The results also indicated that 21-day old seedling may be used at all planting dates sacrificing grain yield to some extent than direct planting with the advantage of about 2 week early harvest. This could save the crop from natural hazards and the grower could avail high market price of grain. The following crops like aus rice, jute, pulses etc. may also be grown in optimum time after harvest of maize. The combination $D_2 \times D_p$ gave the highest stover yield. Stover yield reduced significantly in the 14-d and 21-d under any planting dates. The combination $D_1 \times 21\text{-d}$ had the highest harvest index. The combination $D_2 \times 21\text{-d}$ was also produced statistically similar harvest index to the combination $D_1 \times 21\text{-d}$. However, in general, increased harvest indices were found in the 21-d than 14-d and D_p in all planting dates. Grain yield differed significantly due to interaction effects of planting date and age of seedling. The results showed that direct planting of seed produced higher grain yield but not significantly different from 14- and 21-day old seedlings in November 20 and 30 plantings. At 10 December, 14-day old seedling gave the highest grain yield statistically at par with direct planting but significantly different from 21-day old seedling. However, onward 10 December, 14-day old seedling gave significantly higher grain yields than the others while direct planting of seed and 21-day old seedling yielded statistically similar to each other. But field duration variation was much wider among the planting dates. Late planted crops matured very early and there was short period of grain filling (silking to maturity) compared to early planted crops. December 10 planting produced the longest plant irrespective of variety and seedling age. Planting of 14-day old seedling was clearly found producing higher grain yield than direct planting of seed in most cases and particularly at late condition. The crops planted at the optimum time silked by the end of February and matured by the middle of April. Apparently, this time might be suitable for flowering and grain filling. The photoperiod and temperature of this period are conducive to grain development. The major effect of cool

temperature is the extending duration of grain filling period, increasing total light interception, and enhancing yield. Transplanted crops gave higher grain yields than direct planted crops and might be a possible alternative to direct planting specially under late planting condition (Lazim, 1985 and Badran, 2001). Dale and Drennan (1990a) also found that planting date and seedling age interacted significantly. They reported that 12-day old seedling performed best in early planting, whereas older seedling (42-day old) gave higher grain yield in later plantings. Khehra *et al.* (1990) described that differences due to age of seedling, and date of planting and direct planting were significant for plant height, days to silk and days to maturity. Increased seedling age or delayed transplanting decreased days to silk, days to maturity and plant height. They also recorded the taller plants in direct planted crops.

Table 5. Yield and yield attributes of maize as influenced by the combined effect of planting date and age of seedling during *rabi* 2003-04

Interaction	Plant height at harvest (cm)	Length of cob (cm)	Cob diameter (cm)	Number of cobs m ⁻²	Number of grains cob ⁻¹	TGW (g)	Grain yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)	Harvest index (%)	Field duration (days)
Planting date (D) × Age of seedling (days)										
D ₁ × 0	216.90	18.11	5.10	7.275	414.00	360.15 de	10.61 a	9.59 b	52.37 d	145.5
D ₁ × 14	193.17	17.55	5.03	7.074	412.23	367.68 cd	10.48 ab	9.06 b-d	53.71 cd	138.7
D ₁ × 21	163.30	16.80	5.02	6.763	395.38	382.72 b	10.38 ab	7.98 e	56.69 a	132.2
D ₂ × 0	228.48	17.59	5.09	6.947	428.03	353.43 e-g	10.50 ab	10.48 a	49.71 e	144.7
D ₂ × 14	204.82	17.05	5.04	6.805	414.55	369.10 cd	10.33 ab	8.69 d	54.00 cd	138.2
D ₂ × 21	184.38	16.37	4.98	6.380	396.22	398.20 a	10.26 b	7.99 e	55.81 ab	132.7
D ₃ × 0	231.93	17.28	5.06	6.654	439.57	332.60 hi	9.61 cd	9.26 bc	50.70 e	143.0
D ₃ × 14	222.87	16.65	4.97	6.489	429.93	351.67 e-g	9.77c	8.88 cd	52.37 d	136.5
D ₃ × 21	189.33	16.30	4.92	6.016	409.13	373.23 bc	9.33 de	7.73 ef	54.69 bc	128.8
D ₄ × 0	231.30	16.45	5.01	6.230	436.92	322.57 ij	8.67 f	8.66 d	49.82 e	137.3
D ₄ × 14	202.57	15.98	4.98	6.108	427.60	345.08 fg	9.14 e	7.80 ef	53.73 cd	131.2
D ₄ × 21	177.68	15.67	4.91	5.840	407.27	354.35 ef	8.47 fg	7.33 f	53.43 cd	126.3
D ₅ × 0	211.87	15.90	4.91	5.807	433.33	318.62 j	7.95 h	8.13 e	49.57 e	134.3
D ₅ × 14	197.80	16.38	4.92	5.615	432.03	342.10 gh	8.24 g	7.34 f	52.61d	129.3
D ₅ × 21	173.37	15.23	4.75	5.571	390.87	357.53 de	7.82 h	6.78 g	53.32 cd	124.2
Level of significance	NS	NS	NS	NS	NS	*	*	**	**	

CV %	6.10	4.37	2.05	4.72	6.14	2.51	2.54	5.19	2.34
------	------	------	------	------	------	------	------	------	------

212

213 Figures in a column having no or similar letter(s) do not differ significantly while those with different
 214 letter(s) differ significantly at 5% level of probability; * = Significant at $p \leq 0.05$; ** = Significant at
 215 $p \leq 0.01$; NS = Not significant at 5% level; D₁=November 20, D₂= November 30, D₃= December 10, D₄=
 216 December 20, D₅ =December 30, '0' = Direct seeded.

217 Combined effect of variety and age of seedling, and planting date, variety and age of seedling were
 218 found non-significant for all the characters.

219 4. Conclusion

220 The results revealed that farmers could transplant maize seedling on the mud just after recession of
 221 flood water and thus the crop may be established earlier in the season. There were no significant
 222 difference in respect of grain yield in the earlier November 20 and November 30 planting but the crop
 223 may be harvested about 7 days earlier with 14-day old seedling while 12 days-earlier with 21-day old
 224 seedling. Under late planting condition beyond November 30, transplanting of 14-day old seedling
 225 may produce higher grain yield compared to both direct planting and 21-day old seedling along with 5-
 226 6 days reduced field duration than direct planting of seed. The early harvest may save the crop from
 227 natural vulnerability of hailstorm and rains, and may facilitate in early establishment of the
 228 subsequently crop like sesame, aus rice, mungbean, blackgram, jute etc.

229 References

230 Ahmed F. Maize Production Technology (in Bengali). Intl. Fertilizer Dev. Centre, Bangladesh. 1994;
 231 pp.13-15.
 232 Badran, MSS. Effect of transplanting and seedling age on grain yield and its components of some
 233 maize cultivars. Alexandria J. Agril. Res. 2001;46(2):47-56.
 234 BARI (Bangladesh Agricultural Research Institute). Effects of sowing date on the growth and yield of
 235 maize. Annual Research Report 1996-97. Bangladesh Agril. Res. Inst., Regional Agril. Res.
 236 Inst., Jamalpur, Bangladesh. 1997.
 237 Basu S, Sharma SP, Basu S. Effect of transplanting on vegetative, floral and seed characters of
 238 maize (*Zea mays*) parental lines in spring-summer season. Indian J. Agril. Sci. 2003;73(1):44-
 239 48.

- 240 Bhinder SS, Sawhney JS. Studies on the time and method of sowing winter maize. Agril. Sci. Digest,
241 India. 1985;5(3):145-147.
- 242 Cirilo AG, Andrade FH.. Sowing date and maize productivity: II. Kernel number determination. Crop
243 Sci. 1994b;34(4):1044-1046.
- 244 Dahiya DS, Narwal SS. Effect of sowing date on phenology and growing degree day requirement of
245 winter maize (*Zea mays*). Indian J. Agril. Sci. 1989;59(9):589-592.
- 246 Dale AE, Drennan DSH. Transplanted maize for grain production in Southern England. I. Effects of
247 planting date, transplant age at planting and cultivar on grain yield. J. Agril. Sci., Cambridge.
248 1997a;128(1):27-35.
- 249 Dale AE, Drennan DSH. Transplanted maize (*Zea mays*) for grain production in southern England. II.
250 Effects of planting date, transplant age at planting and cultivar on growth, development and
251 harvest index. J. Agril. Sci., Cambridge. 1997b;128(1):37-44.
- 252 Gomez KA, Gomez AA. Statistical Procedures of Agricultural Research (2nd ed.). A Wiley-Interscience
253 Publication, John Wiley and Sons, New York. 1984.
- 254 Ibrahim SM, Gopalasamy N. Effect of age of seedlings on growth and yield of transplanted maize.
255 Madras Agril. J. 1989;76(4):181-183.
- 256 Narwal SS, Poonia S, Singh G, Malik DS. Influence of sowing dates on the growing degree days and
257 phenology of winter maize (*Zea mays* L.). Agril. Forest Met. 1986;38(1-3):47-57.
- 258 Oswald A, Ransom JK, Kroschel J, Saureborn J. Transplanting maize and Sorghum reduces *Striga*
259 *hermonthica* damage. Weed Sci. 2001;49(3):346-353.
- 260 Paliwal RL. Introduction to maize and its importance. In: Tropical Maize Improvement and Production.
261 Food and Agriculture Organization of the United Nations, Rome, Italy, 2000.
- 262 Prodhan HS. Influence of appropriate cultivar and sowing dates on rabi maize productivity. Indian J
263 Agril. Sci. 2001;71(1): 55-56.
- 264 Rai B, Jha MM, Kumar B. Effect of direct seeded and transplanted maize in relation to the
265 development of Turcicum leaf blight disease. Ann. Agri-Bio-Res. 2002;7(1):55-58.
- 266 Razzaque MA, Satter MA, Amin S, Quayyum A, Alom MS. Edited. *Krishi Projukti Hatboi* (In Bengali).
267 2nd Edn. Bangladesh Agril. Res. Inst., Gazipur 1701, Bangladesh; 2000.
- 268 Sandhu BS, Hundal SS. Effect of method and date of sowing on phenophases and productivity of
269 winter maize (*Zea mays*). Indian J. Agril. Sci. 1991;61(3):178-181.

- 270 Sharma RK, Brar HS, Khehra AS, Dhillon BS. Gap filling through transplanting in maize during winter.
271 J. Agron. Crop Sci. 1989;162(3):145-148.
- 272 Singh VK, Bajai RP, Sisodia RJ. Effect of sowing dates and plant population on yield of maize.
273 Current Res., Univ. Agril. Sci., Bangalore. 1992;21(2):24-25.
- 274 Sudipta B, Sharma SP, Basu S. Effect of transplanting on vegetative, floral and seed characters of
275 maize (*Zea mays*) parental lines in spring-summer season. Indian J. Agril. Sci. 2003;73(1):44-
276 48.
- 277 Thompson LM. Climatic change, weather variability and corn production. Agron. J. 1986;78:649-653.
- 278 UNDP, FAO. Land Resources Appraisal of Bangladesh for Agricultural Development. Report No. 2.
279 Agro-Ecological Regions of Bangladesh. United Nations Development Programme and Food
280 and Agriculture Organization. 1988;pp. 212-221.
- 281 Uy T H. 1996. Transplanting maize on wetland. A technical manual based on a successful case-study
282 in Vietnam. Food and Agriculture Organization of the United Nations, Rome, Italy; 1996.
- 283 Vasa, SK, McLean S. The Lowland Tropical Maize Subprogram. Maize Program Special Report
284 Mexico, CIMMYT, Mexico; 1994.
- 285 Yousafzai HK, Muhammad A, Rozina G, Naseer A, Khan IA. Effect of sowing dates on maize
286 cultivars. Sarhad J. Agric. 2002;8(1):11-15.
- 287 Zheng HJ, Dong ST, Wang KJ, Guo Y, CHHu. Effects of ecological factors on growth of different
288 maize varieties and corresponding regulative measures. J. Shandong Agril. Univ.
289 2001;32(2):117-123.