

Research Paper

**RESPONSE OF MAIZE TO ORGANIC AND INORGANIC SOURCES OF NUTRIENTS
IN ACID SOILS OF KERICHO COUNTY, KENYA**

ABSTRACT

Maize yields in Kericho County are limited mainly by infertile acidic soils. The effect of inorganic sources of nutrients and amendments; triple superphosphate (TSP), calcium ammonium nitrate (CAN) and lime, were compared to a range of manures; Farmyard manure (FYM) of low and high quality, dried cow dung, goat manure, tithonia applied as green manure or dried, in a greenhouse and field experiment. Two soils collected from two farmers fields (Joseph and Samuel) were used in the greenhouse where maize was grown for six weeks and its biomass yield determined. The treatments that showed promise were used in a subsequent field experiment where maize was grown to maturity and grain yields determined. In the greenhouse, maize responded to application of all the sources of nutrients and amendments, except lime when applied without TSP, on Joseph's soil. On Samuel's soil, maize did not respond to application of lime alone or with TSP, TSP and dried tithonia. High quality FYM gave the highest increase (136%) in dry matter yields on Samuel's soil. In the field experiment, goat manure gave the highest grain yields. Maize failed to significantly respond to either CAN or TSP when applied alone but the application of the two in combination (TSP + CAN) effected a significant response indicating that both N and P were deficient in this soil. All the manures, except low quality FYM, gave yields that were higher or comparable to the standard recommended fertilizer practice (TSP + CAN) and could be economically attractive substitutes as they are locally

23 available. There was a  low correlation between dry matter biomass yield in the greenhouse and
24 grain yield in the field likely due to different durations in growth hence differences in nutrient
25 demands. Extrapolation of greenhouse findings to different fields should therefore be treated
26 with caution.

27 Key words: *Acid soils; inorganic fertilizers; maize yield, organic materials*

28 1.0. INTRODUCTION

29 The densely populated humid regions of western Kenya have experienced high rates of soil
30 fertility depletion [1]. In addition to the widespread P-deficiencies, nitrogen and potassium
31 limitations are common, especially when P limitations have been overcome [2]. The situation is
32 exacerbated by the high acidity of these soils with possibility of aluminum toxicity.  Such
33 conditions  obtain in Kericho County. Thus although most parts of Kericho receive favorable
34 rainfall, crop productivity is seriously constrained, with yields of the staple food maize rarely
35 exceeding 1 ton ha⁻¹ on smallholder farms. Consequently, food insecurity is common in the
36 region.

37 Management of nutrient deficiencies requires addition of organic or inorganic fertilizers or
38 their combinations. Present management practices for amelioration of P-deficiencies and soil
39 acidity include the utilization of the low input strategy consisting of spot or band fertilizer
40 placement and use of soil amendments such as lime [3]. Use of the low input strategy, however,
41 remains negligible among the smallholder farmers in Kenya due to the ever increasing costs of
42 purchased fertilizer (particularly) and lime inputs. In addition, fertilizers alone cannot guarantee
43 long-term productivity on many soils, hence inputs of organic materials are needed to maintain
44 soil organic matter (SOM) levels [4]. Thus in Kenya, there is growing interest in finding

45 alternative, affordable, viable and sustainable technologies to replenish soil fertility with focus
46 shifting to solutions that utilize more local resources and less external inputs [5].

47 The use of organic amendments, in particular could be both agronomically and economically
48 attractive because they can serve the dual purpose of providing a range of plant nutrients while
49 ameliorating soil acidity [6]. Several studies have demonstrated the positive effects of OM s in
50 greenhouse studies and assumed the same would hold under field conditions. There are not many
51 studies, however, comparing the results of greenhouse and field experiments. Greenhouse
52 experiments have the advantage of a good control of environmental variables but with a lack of
53 realism, while field experiments are the opposite [7]. The objective of this study was therefore to
54 i) explore the effectiveness of a range of organic amendments, commonly available on
55 smallholder farms in Kericho County, in terms of ameliorating soil acidity, increasing available
56 phosphorus and nutrient uptake by maize in a greenhouse pot experiment and field experiment
57 and ii) to correlate the results obtained under greenhouse conditions to those of field conditions.

58

59 2.0. MATERIALS AND METHODS

60 2.1. Site description

61 The study was conducted at the Kabianga University College Farm, in Kericho County,
62 Kenya. The site lies at an altitude of 2163 m above sea level and is within latitude 0° 49' 0 N and
63 longitude 35° 49' 60 E. The average rainfall is 900 - 1200 mm per annum distributed mainly
64 between the months of March and December with two distinct peaks in May and October. The
65 soils at the site are Nitisols which are acidic, moderately deep and well drained [8].

2.2. Preparation of organic materials and analyses

Organic materials which represent common alternatives to inorganic fertilizers on smallholder farms, but varying in their chemical characteristics, were used. These were farmyard manures of varying quality, *Tithonia diversifolia* (tithonia) and goat manure. They were obtained in the vicinity of the experimental site and air-dried. A sub sample of each OM was then oven-dried at 70°C and ground to pass through a 0.5 mm sieve. The nutrient content of these organic materials (total N, P, K, Ca, Mg) were determined as described by [9].

2.3. Experimental layout and management

2.3.1. The greenhouse pot experiment

Soil (0-20 cm) was collected randomly from various spots at two farmers (Joseph and Samuel) fields that were reportedly not responding to fertilizers. The collected soil was air-dried, sieved through a 4 mm mesh and applied in all the pots at a rate 4 kg of soil pot⁻¹.

The proposed treatments are given below.

1. Control

2. Lime

3. Lime + TSP

4. TSP

5. Dried cattle dung

6. FYM high quality

7. FYM low quality

8. Tithonia green manure

9. Tithonia dry

10. Goat manure

A randomised complete block design with three replicates was installed. All the treatments with no organic materials application received muriate of potash (60 kg K ha^{-1}) and CAN (100 kg N ha^{-1}) to ensure that all the major nutrients (except the control) were non-limiting [10]. The organic amendment were analysed at the Mumias Sugar Company laboratory and their characteristics are given in Table 1. All organic materials were applied at a rate of 5 t ha^{-1} and TSP at 60 kg P ha^{-1} . Lime was applied at 4 t ha^{-1} to the appropriate pots receiving lime treatment and incubated for 30 days at approximately field water holding capacity to allow reaction before planting. The other treatments were applied on the day of planting (15th March 2011) after which 4 maize seeds were planted per pot at a depth of approximately 2 cm. The maize plants were thinned to 2 per pot one week after emergence from the soil. Soil water was maintained at almost equal levels for all treatments by regular watering and weeds were regularly removed by hand. Each treatment was destructively harvested at 6 weeks after planting by cutting the maize tops at soil level. The harvested plants were oven dried at 70°C to a constant weight.

102

2.3.2. The Field Experiment

A Field trial was conducted at the Kabianga University College Farm for one season (May to November 2011) to test whether the observed responses in the green house study could be replicated under field conditions. However, lime was not included in the field experiment because of its poor performance in the greenhouse while tithonia was excluded because of its unavailability in adequate quantities in the study area. A randomized complete block design with 3 replicates was used with treatment plot sizes of $4 \text{ m} \times 3 \text{ m}$. The imposed treatments were as follows;

1. Control
2. Calcium ammonium nitrate (CAN)
3. Triple superphosphate (TSP)
4. TSP + CAN
5. Dried cow dung
6. FYM low quality
7. FYM high quality
8. Goat manure

All the manures were applied at 5 t ha⁻¹ while TSP was applied to supply 60 kg P ha⁻¹. All the nutrient inputs were uniformly broadcast on the plots and then incorporated into the top soil (0-15 cm) by a hand hoe prior to planting of maize. CAN at 100 kg ha⁻¹ was split applied in plots without organic amendments, one third at planting and two-thirds at 5 weeks after planting. The rates of fertilizers and manures were based on recommendations [10] and previous studies in western Kenya [11]. The experiment was managed using recommended agronomic practices for the study area. The grain and stover yields were determined when maize matured.

2.4. Data analyses

Analysis of variance (ANOVA) was conducted to determine the effects of treatments on the measured parameters. The Least Significant difference between means (LSD) was used to compare the treatment means. Mention of statistical significance refers to $p = 0.05$ unless otherwise stated. Quantitative relationships between dry matter yields obtained in the greenhouse study and stover and grain yields, obtained in the field study were developed by regression analyses. Statistical analysis was conducted using the Genstat statistical package [12].

136 3.0. RESULTS

137 3.1. Nutrient content of the organic materials used in the study

138 Tithonia had the highest levels of N, K and Mg while FYM of high quality had the highest
 139 amounts of P (Table 1). The nutrient content of the high quality FYM was almost double that of
 140 the poor quality FYM. Apart from the low P content, the nutrient content of goat manure was
 141 comparable to that of high quality FYM.

142

143 Table 1. Nutrient content of the experimental organic materials

Organic material	Parameter				
	N%	P%	K%	Ca	Mg
FYM high quality	2.27	1.50	0.88	0.41	0.24
FYM low quality	0.98	0.62	0.56	0.23	0.16
Dried cattle dung	1.18	1.30	0.64	0.32	0.21
Tithonia	4.23	0.41	2.64	1.81	0.38
Goat manure	2.21	0.36	0.52	0.53	0.31

144 3.2. Greenhouse study

145 The dry matter yields for maize grown in the soils collected from the two sites in the
 146 greenhouse are presented in Table 2. There were highly significant ($p < 0.001$) differences
 147 among the treatments at in both soils. However, the response to the application of the nutrient
 148 sources and amendments varied with the soil. Maize responded to application of all the sources
 149 of nutrients and amendments, except lime when applied without TSP, on Joseph's soil.
 150 Application of goat manure gave the highest yield increase (135%) above the control for this soil
 151 (Figure 1). On Samuel's soil, maize did not also respond to application of lime when applied

152 alone. In addition, maize did not respond to lime applied with TSP, TSP and dried tithonia on
 153 this soil. High quality FYM gave the highest increase (136%) in dry matter yields on Samuel's
 154 soil (Figure 1). Among the cattle manures, high quality FYM was significantly better than dried
 155 cattle dung but not low quality manure on both soils. Low quality FYM had slightly higher but
 156 none significant yields compared to the dried cattle dung. Tithonia green manure gave
 157 significantly higher yields than dried tithonia on Joseph's soil but not Samuel's. Generally,
 158 organic sources of nutrients performed better than inorganic fertilizer TSP on both soils.


159

160 Table 2. Effect of nutrient sources on biomass dry matter yields

Treatment	Dry matter yield (g/pot)	
	Joseph	Samuel
1. Control	11.5	11.9
2. Lime	15.0	9.2
3. Lime + TSP	19.4	11.4
4. TSP	21.6	15.2
5. Dried cattle dung	20.1	21.6
6. FYM high quality	26.5	27.9
7. FYM low quality	23.3	25.1
8. Tithonia green	26.6	17.6
9. Tithonia dry	20.5	15.0
10. Goat manure	27.9	25.9
LSD	5.2	5.0

CV%	20	16
-----	----	----

161



162

163 **Figure 1. % Increase in drymatter yield over the control**

164

165 3.3. Field experiment

166 There were significant treatment effects ($p < 0.01$) on stover yields which ranged from 11.43
167 t ha^{-1} (control) to 16.76 t ha^{-1} (high quality FYM) (Table 3). Maize did not respond to application
168 of CAN alone or FYM of low quality but gave significantly higher stover yields relative to the
169 control for all the other treatments. Unlike in the greenhouse study where **dried cowdung** was
170 inferior to FYM of high quality, in the field experiment the two were not significantly different.
171 Among the manures, only FYM of high quality gave higher stover yields than TSP. The rest had
172 comparable yields with those of TSP. The maize grain yields ranged from 0.76 t ha^{-1} (CAN) to
173 3.25 t ha^{-1} (goat manure). Maize failed to respond to either CAN or TSP when applied alone but

the application of the two in combination (TSP + CAN) effected a significant **resonse**. All the manures, except low quality FYM gave grain yields that were higher or comparable to the standard recommended fertilizer **practice** (TSP + CAN). The increase in grain yield was dependent on the type of the manure and followed the order goat manure > high quality FYM > dried cow dung > low quality FYM. The increases, over the control, for grain yields were much higher than those of stover. For example, goat manure increased grain yield by 242% but increased stover by only 33% (Figure 2). The trend was observed in all the other treatments.

181

182 Table 3. Maize stover and grain yields as affected by nutrient sources

Treatment	Yield t/ha	
	Maize stover	Grain
1. Control (no nutrient input)	11.43	0.95
2. Calcium ammonium nitrate (CAN)	12.44	0.76
3. Triple superphosphate (TSP)	13.71	1.47
4. TSP + CAN	14.72	2.70
5. Dried cow dung	14.22	2.57
6. FYM low quality	13.20	1.43
7. FYM high quality	16.76	2.93
8. Goat manure	15.23	3.25
LSD	2.14	0.67
CV%	10	19

183




Figure 2. % increase in stover and grain yield over the control

There was a poor correlation between the biomass dry matter yields obtained in the greenhouse and the grain yield in the field study for both Joseph ($r^2 = 0.47$) and Samuel's soils ($r^2 = 0.29$). (Figures 3 and 4). The correlation with stover was also low for Samuel's soil. However, there was a high correlation ($r^2 = 0.74$) of the greenhouse biomass to stover yield for Joseph's soil.




Figure 3. Relationship between biomass dry weight yield (Joseph's soil) in greenhouse with stover and grain yields in the field.




Figure 4. Relationship between biomass dry weight yield (Samuel's soil) in greenhouse with stover and grain yields in the field.

4.0. DISCUSSIONS

The responses to the nutrient sources in the greenhouse were soil specific likely due to differences in the soil properties of the two soils. Such soil specific responses have been reported by others [13]. The failure of maize to respond to TSP on Samuel's soil and the low response on Joseph's soil confirmed the farmers' observations on the ineffectiveness inorganic fertilizers on their farms. It had been hypothesized that high acidity of these soils could be causing Al toxicity, but the failure of lime application to increase yields suggest other limitations. The synergism that was observed when lime was combined with TSP compared to application of lime alone or TSP alone on Joseph's soil (Table 2) suggests a simultaneous P deficiency and acidity problem. However, on Samuel's soil where no response to lime and TSP, whether applied alone or in combination, was observed, other factors other than acidity and P deficiency could be at play. The soil in the field study did not respond to N when applied alone (CAN) but responded to P applied alone implying that P was more limiting than N although both N and P were deficient as confirmed by the higher yields when TSP was combined with CAN. This confirms reports by [14] that most soils in western Kenya do not respond to N unless P limitations are overcome.

Application of organic materials generally had positive effects on yields but the magnitude varied with the type and quality of organic input. FYM of high quality out-performed that of low quality both in the greenhouse and field while goat manure was better than the other manures likely due to their high nutrient content. All the manures were applied at 5 t ha^{-1} and therefore the amount of nutrients supplied by them varied with their quality. For example, for the most limiting nutrients N and P, FYM of high quality provide 114 kg of N and 75 kg P ha^{-1} compared to 49 and $31 \text{ kg of N and P ha}^{-1}$ respectively for the low quality FYM. This does not however explain the slightly higher or comparable yields obtained by the high quality OM_s in the field compared to the standard to inorganic fertilizers; TSP combined with CAN, whose application provided 60 kg P ha^{-1} and 100 kg N ha^{-1} respectively. However, there are other factors other than nutrient content that could also have contributed to the responses observed among the manures. For example, low biomass yields were observed for dried tithonia compared to the green tithonia although both provided the same quantity of nutrients. This is likely due to increased polyphenol content and reduced nutrient release following drying of plant biomass [15]. In addition, organic materials positively affect other soil properties such as cation exchange capacity, improvement of soil structure and moisture holding capacity [16, 17]. They also provide other nutrients such as micronutrients which were not provided by TSP and CAN. This could also have contributed to their general superiority over inorganic fertilizers.

There low correlations between dry matter biomass yield in the greenhouse and grain yield in the field could be attributed to the fact that in the greenhouse, the maize was harvested before the period of high nutrient demand unlike in the field where the crop grew to maturity. Therefore some of the treatments that performed well in the greenhouse e.g., FYM of low quality may have failed to supply enough nutrients demanded for grain production in the field. This is

corroborated by the generally better correlation with stover yields, especially for Joseph's soil, suggesting that some of the inputs were able to meet the nutrient requirements of the vegetative stages of growth (biomass) but were not adequate for grain production. Furthermore, some of the manures e.g. the goat manure and dried cattle dung may not have had not had enough time to mineralize in the short period in the greenhouse (6 weeks) and the longer period in the field, coupled with more favorable conditions for mineralization compared to the greenhouse, allowed them to perform better in the field. Moreover, the use of different soils in the greenhouse and field experiment and differences in their environments may have contributed to the poor correlation between greenhouse and field data.

5.0. CONCLUSIONS

The responses to the nutrient sources in the greenhouse were soil specific. Joseph's soil was more responsive to most of the organic and inorganic inputs tested while Samuel's soil was none responsive to inorganic inputs and low quality manures. Among the manures, quality in terms of nutrient content and ability to mineralize seemed to have largely affected their effectiveness with goat manure and FYM performing well both in the greenhouse and field. In the field experiment, maize failed to significantly respond to either CAN or TSP when applied alone but the application of the two in combination (TSP + CAN) effected a significant response suggesting a simultaneous deficiency of N and P at this site. All the manures tested, except low quality FYM, gave grain yields that were higher or comparable to the standard recommended fertilizer practice (TSP + CAN) in the study area and could be economically attractive substitutes as they are locally available. Differences in response by maize to the different organic and inorganic inputs on the two soils in the greenhouse and lack of correlation between biomass dry matter yields in the greenhouse with the maize grain yields in the field indicate that

259 recommendations of nutrient sources for maize cannot easily be transposed among diverse soils
260 of Kericho County or from greenhouse to field.

261

262 REFERENCES

263 1. Smaling EMA, Stoorvogel JJ, Windmeijer PN. Calculating soil nutrient balances in Africa in
264 different scales: II. District scale. *Fert. Res.* 1993; 35:237-250.

265 2. Jama B, van Straaten. Potential of East African phosphate rock deposits in integrated nutrient
266 management strategies. *Annals of Brazilian Academy of Sciences.* 2006; 78:781-790.

267 3. Sanchez PA, Salinas JG. Low-input technologies for managing Oxisols and Ultisols in Tropical
268 America. *Adv. Agron.* 1981; 34:279-406.

269 4. Syers JK. Managing soils for long-term productivity. *Philosophical Transactions of the Royal*
270 *Society of London. Series B, In: Biological Sciences.* 1997; 352:1011-1021.

271 5. Pretty J and Hine R. Reducing Food Poverty with Sustainable Agriculture. A summary of new
272 evidence. Final report from SAFE-World Research Project. Colchester, U.K. University of
273 Essex. 2001.

274 6. Opala PA, Othieno CO, Okalebo JR, Kisinyo P. Effects of combining organic materials with
275 inorganic phosphorus sources on maize yield and financial benefits in western Kenya.
276 *Experimental Agriculture.* 2010; 46:23-34.

277 7. Marañón T, Zamora R, Villar R, Zavala MA, Quero JL, Pérez-Ramos I, Mendoza I, Castro J.
278 Regeneration of tree species and restoration under contrasted Mediterranean habitats: field and
279 glasshouse experiments. *International Journal of Ecology and Environmental Sciences* 2004; 30:
280 187-196.

- 281 8. Jaetzold R, Schmidt H. Farm management handbook of Kenya Vol. 2. Natural conditions and
 282 Farm management information. Part A: Western Kenya. Ministry of Agriculture. Nairobi,
 283 Kenya; 1982.
- 284 9. Anderson JM, Ingram JSI. Tropical soil Biology and Fertility: A handbook of methods.
 285 Wallingford: CAB International; 1993.
- 286 10. FURP. Fertilizer Use Recommendation Project. Compilation of results from former fertilizer
 287 trials in Kenya. Ministry of Agriculture, Nairobi, Kenya; 1987.
- 288 11. Nziguheba G, Merckx R, Palm CA, Mutuo P. Combining *Tithonia diversifolia* and fertilizers
 289 for maize production in phosphorus deficient soil in Kenya. *Agroforestry Systems*. 2002;55:
 290 165-174.
- 291 12. GENSTAT. GenStat Procedure Library Release PL20.2. GenStat Release 12.2 (PC/Windows
 292 XP). Copyright 2010, VSN International Ltd; 2010.
- 293 13. Sakala WD, Kumwenda JDT, Saka AR. The Potential of Green Manures to Increase Soil
 294 Fertility and Maize Yields in Malawi pp, 378-384. In: A. Bationo (ed). *Managing nutrient cycles*
 295 *to sustain soil fertility in sub-Saharan Africa*. Nairobi: Academy Sciences Publishers; 2004.
- 296 14. Jama B, Swinkels RA, Buresh RJ. Agronomic and economic evaluation of organic and
 297 inorganic sources of phosphorus in western Kenya. *Agron. J.* 1997; 89:597-604.
- 298 15. Mafongoya PL, Nair PKR. Multipurpose tree prunings as a source of nitrogen to maize under
 299 semiarid conditions in Zimbabwe. 1. Nitrogen recovery in relation to pruning quality and method
 300 of application. *Agroforestry Systems*. 1997; 35: 31–46.
- 301 16. Palm CA, Gachengo CN, Delve C, Cadisch G, Giller KE. Organic inputs for soil fertility
 302 management in tropical agroecosystems: Application of an organic resource database. *Agric.*
 303 *Ecosyst. Environ.* 2001; 83:27-42.

- 304 17. Opala PA, Okalebo JR, Othieno C. Comparison of effects of phosphorus sources on soil
305 acidity, available phosphorus and maize yields at two sites in western Kenya, Archives of
306 Agronomy and Soil Science, 2012: DOI:10.1080/03650340.2011.627681.