

EFFECT OF DIFFERENT RATES OF PHOSPHORUS FERTILIZER ON SOYBEAN {GLYCINE MAX (L.) MERRIL} INOCULATED WITH RHIZOBIUM

Abstract

The experiment was conducted during the 2012 farming season at the University for Development Studies, Nyankpala in the Northern Region of Ghana. The objective of the study was to determine the influence of rhizobium inoculants and phosphorus (P) at different application rates on yield and yield components of soybean. Two levels of inoculation regimes (un-inoculated (-In) and inoculated (+In)) were combined with four application rates of phosphorus (0 kg P/ha, 15 kg P/ha, 30 kg P/ha and 45 kg P/ha). The experiment was laid in a Randomized Complete Block design with three replications. Parameters measured were crop emergence, plant height, leaf area, canopy spread, number and weight of nodules, number of pods and total grain yield. The results obtained indicated significance differences in all the parameters measured. Phosphorus application at 45 kg P/ha plus Rhizobium inoculants recorded significantly higher growth and yield parameters than the rest of the treatments. The economic analysis of the treatments also showed that inoculation of soybean seeds or a combination of inoculation with 45 kg P/ha was more profitable than the application of 45 kg P/ha without inoculation. The study recommends inoculation of soybean seeds with Rhizobium inoculants in association with application of phosphorus fertilizer at 45 kg P/ha.

Keywords: Inoculation, phosphorus rates, Yaralegume, Grain yield

1.0 Introduction

Soybean {*Glycine max* L. (Merrill)} is one of the important oil grain legume crops in the world. In the international world trade market, soybean is ranked number one in the world among the major oil crops such as rapeseed, groundnut, cotton seed, sunflower, linseed, sesame and safflower [1]. Soybean is the cheapest protein source as compared to other protein sources such as egg, fish and meat. With an average protein content of 20% on dry matter basis, soybean has the highest protein content of all field crops and is second only to groundnut in terms of oil content among the food legumes [2] and [3]. Besides, its rich nutrition source for human and animals, it fixes atmospheric nitrogen which is a major boost

37 to African farmers where the soils are low in nitrogen and commercial fertilizers are
38 expensive to afford. It is also an important source of income to farmers [4].

39 In spite of its great potential, its production is still limited by low yields resulting in a wide
40 gap between current production levels and the crop's potential. In order to improve
41 production levels, areas to address include development of high yielding varieties, disease
42 and pest control and development of improved cultural practices [5]. In northern Ghana,
43 where the largest production occurs in the country, the average yield is about 2.5 tonnes/ha
44 [6] as compared to that of USA which is 4.6 tonnes/ha [7]. The low yields can be attributed to
45 soils which are inherently poor and deficient in organic matter and other vital nutrients such
46 as phosphorus and nitrogen. Studies in parts of Northern, Upper West and Upper East
47 Regions showed that available phosphorus is very low of about 6.0 mg/kg soil [8].
48 Phosphorus deficiency can limit nodulation by legumes and phosphorus fertilizer application
49 can overcome the deficiency [9]. Availability of phosphorus in the soil influences the
50 efficiency of *Rhizobium* that fixes atmospheric nitrogen in association with nodulating
51 legumes as it is directly involved in biological nitrogen fixation via legume-*Rhizobium*
52 symbiosis. This is attracting considerable research attention world-wide due to its economic
53 viability for resource poor farmers and environmental friendliness [10], [11] and [12]. The
54 objective of this study was to assess the impact of phosphorus at different application rates
55 and *Rhizobium* inoculants on the growth and yield of soybean.



56 2.0 Methodology

57 2.1 Site description

58 The experiment was  ducted on the experimental field of the Faculty of Agriculture of the
59 University for Development Studies, Nyankpala in the Northern Guinea savannah ecological
60 zone during the 2012 cropping season. The area experiences a unimodal annual rainfall of
61 1000 mm-1200 mm from April to November. The temperature distribution is uniform with
62 mean monthly minimum and maximum values of 21°C and 34.1°C, respectively and a
63 relative humidity of 53%-80% [13].

64

65 2.2 Experimental design

66 A 2 x 4 factorial experiment consisting of two inoculation regimes and four application rates
67 of phosphorus fertilizer were laid out in a Randomized Complete Block design with four

replications. The P treatments were 0 kg P/ha, 15 kg P/ha, 30 kg P/ha and 45 kg P/ha from Yaralegume fertilizer with or without inoculants. The plot size was 2.5 m x 2.5 m. Yaralegume fertilizer is composed of 0%N-18% P_2O_5 13% K_2O +31%Ca+4%S+2%Mg+0.6%Zn. This was applied two weeks after sowing by burying it in a trench dug about 5 cm away from the plant hill.

2.3 Planting materials and sowing

The improved soybean variety, **Jenguma** which was used was obtained from the CSIR-Savanna Agricultural Research Institute (SARI) located at Nyankpal. For the inoculated treatments, the seeds of Jenguma were inoculated with commercial Rhizobium inoculants, *Legumefix* (containing 1×10^9 cells/g) at the rate of 1 kg seed per 5 g of inoculants [14] before sowing. The sowing was done early in the morning to avoid exposing the inoculants to the direct rays of the sun which might affect the quality of the inoculants. Sowing was done at three seeds per hill at a distance of 50 cm between ridges and 10 cm within ridges. In order to avoid contamination of the un-inoculated treatments with the inoculants that might get stuck to the hand during sowing, the un-inoculated seeds were sown before the inoculated ones. The plants were allowed to grow to maturity under rain-fed conditions while the necessary management practices like weeding were observed.

2.4 Data collection

Data were collected on percent crop establishment, plant height, leaf area, canopy spread, shoot biomass, number of nodules, nodule biomass, number and dry weight of pods at maturity and grain yield. Partial budget analysis for the treatments was also carried out.

Percent crop establishment was determined three weeks after sowing by counting the emerged plants in the four inner ridges and dividing the number by the expected number of plants (i.e. number of seeds sown per the four hills) multiplied by 100. Plant height, leaf area and canopy spread were measured in the third, sixth and ninth weeks after sowing (3 WAP, 6 WAP and 9 WAP). A meter rule was used to take the height of each of the eight plants randomly selected from each of the four inner ridges (two plants per ridge), respectively and tagged by measuring from the base of the plant (ground level) to the top of the main stem.

For leaf area, one leaf each was selected from the lower, middle and upper parts of each plant and the breadth and length were measured with a meter rule. The product of these two parameters constitutes the area of the leaf. The means of the areas of the three leaves on each

plant were taken as the leaf area of that plant while the mean leaf area of the eight sampled plants on each plot (treatment) was calculated and recorded as the leaf area of that plot. Canopy spread was measured from the last leaf on one side of each of the eight plants to the last leaf on the other side of the same plant using a measuring tape.

For the assessment of shoot biomass and nodulation, the eight plants which had been tagged for plant height measurements were sampled at full pod stage (R4 stage) by gently pulling them out of the ground after carefully loosening the soil around the plants with a cutlass before pulling them out gently with the hand making sure to collect the few nodules that got detached from the roots in the process. The shoots from each plot were then severed from the root system, bulked and weighed on a digital scale after which they were air-dried for one day followed by oven-drying at 80°C for 48 hours (to a constant weight) and the weights recorded. The nodules were detached from the roots, counted, weighed and then oven-dried at 70°C for 48 hours and the weights recorded.

At harvest, all the pods from the plants of the two innermost ridges of each plot were harvested manually and counted after which they were sufficiently sun-dried (to about 11-13% moisture content) on a concrete platform and weighed on a digital scale. The dried pods were threshed and winnowed to separate the grains from the husk and the former weighed to obtain the grain yield.

2.5 Data analysis

Data collected were subjected to a one-way analyses of variance (ANOVA) using the computer statistical package Genstat (2008 Edition) and treatment means compared using the least significance difference (LSD) at 5% probability level.

3.0 Results and Discussion

3.1 Percent crop establishment

Crop establishment was seen to be significantly better in inoculated plots than in uninoculated plots and this was observed irrespective of inoculation regime though the effects due to higher P application rates were more pronounced in inoculated treatments (Figure 1). There was therefore an interaction effect between P fertilizer application rate and Rhizobium inoculation resulting in the highest percent crop establishment being observed in treatments with joint applications of 45 kg P/ha and Rhizobium inoculants. This observation confirms findings by [15] who intimated that increased plant establishment from seeds appears to be a

result of improved soil productivity due to bacterial activity and available nutrients. Increases in whole plant growth in response to phosphorus supply have been reported in soybean [16], cowpea [17] and groundnut [18]. With Rhizobium inoculation showing a better crop establishment than without inoculation in the 0 kg P/ha treatment, the improved establishment could be attributed to rhizobial activity and perhaps also to a vigorous germination and growth of the seeds used.




Figure 1: Influence of Rhizobium inoculation and phosphorus fertilizer application rates on establishment of soybean

3.2 Plant height

The application of phosphorus fertilizer and Rhizobium inoculants impacted on the growth of soybean when compared to the non-phosphorus and non-Rhizobium treatments (Figure 2). As expected, the plants grew taller with time within a fertilizer application regime, especially when inoculated (Figure 2). Irrespective of growth stage, plants fertilized with 45 kg P/ha and inoculated grew tallest (51.8 cm). The current study confirms other research findings by [19] and [20] that increased levels of phosphorus significantly increased plant height. Also, [21] observed that Rhizobium inoculation increased plant height. [22] reported incidences of increased plant height due to interactions between phosphorus fertilizer application and

Rhizobium inoculants whereas [23] observed that over-supply of phosphorus resulted in decreases in plant growth, nodulation and acetylene reduction in soybean.





Figure 2: Influence of Rhizobium inoculation and phosphorus fertilizer application rate on plant height of soybean at different growth stages.

3.3 Leaf area

The response of leaf area of soybean to Rhizobium inoculation and rate of P fertilizer (Figure 3) showed a similar trend as that of plant height (Figure 2). Inoculation increased leaf area at all the three stages of sampling irrespective of the rate of P applied (Figure 3) and whether plants were inoculated or not, leaf area increased with plant age becoming more significant at 9 WAP in uninoculated plants. However, leaf area was highest in all inoculated plants fertilized with 45 kg P/ha compared with other treatments. The results of the current study confirm results of a similar study conducted by [24] who reported that primary whole plant response to phosphorus addition was an enhancement in shoot growth caused by increases in leaf area expansion. In a similar study on common bean, the observed increase in cell division due to phosphorus application resulted in increasing amounts of ATP at growth centers [25] which affected many plant characteristics such as leaf area.




172

173 Figure 3: Influence of Rhizobium inoculation and phosphorus fertilizer application rate on
 174 leaf area of soybean at different growth stages.

175 3.4 Canopy spread

176 Phosphorus application rate, Rhizobium inoculation and their interaction significantly
 177 affected canopy spread of soybean plants (Figure 4). Phosphorus application rate of 45 kg
 178 P/ha with Rhizobium inoculation recorded the highest canopy spread at 3, 6 and 9 WAP
 179 compared to the spreads at 15 kg P/ha and 30 kg P/ha 9 WAP. The above results are in line
 180 with the findings of [26] that application of phosphorus significantly increased most of the
 181 soybean growth parameters.




182

183 Figure 4: Influence of Rhizobium inoculation and phosphorus fertilizer application rate on
184 canopy spread in soybean at different growth stages.

185 3.5 Shoot biomass

186 Shoot biomass per plot was significantly influenced by Rhizobium inoculations and
187 increasing fertilizer rates (Figure 5). The highest shoot biomass was obtained in inoculated
188 plants fertilized with 45 kg P/ha whilst plants that received no P application with or without
189 inoculation produced the lowest shoots biomass. Across the P application rates, inoculation
190 increased soybean shoot dry weight by a significant average of 30% over the uninoculated
191 plants varied phosphorus rates and their interactions (Figure 5). Shoot biomass increased with
192 increasing application of phosphorus + Rhizobium inoculants. Phosphorus at 45 kg P/ha +
193 Rhizobium inoculant recorded the highest shoot biomass per plot as compared to 15 kg P/ha
194 + Rhizobium inoculants which recorded the least with regards to phosphorus and Rhizobium
195 inoculants. With the sole fertilizer treatments, shoot biomass increased with increasing
196 phosphorus rates with 45 kg P/ha recording the highest. At low phosphorus application, the
197 straws yield was lower than treatments that received the highest amount. In an earlier work,
198 the authors reported of similar increases in shoot biomass of soybean as a result of increased
199 rates of P from different fertilizer sources [27]. The significant increase in shoot biomass due
200 to increased P application rates in the current study also agrees with [28] who reported of a
201 20.7% increase in biomass yield due to phosphorus application.



202

Figure 5: Influence of Rhizobium inoculation and phosphorus fertilizer application rate on shoot dry biomass of soybean at full pod stage.

3.6 Nodulation

Nodule number was significantly higher in inoculated plants and increased with P application rate whether plants were inoculated or not (Figure 6). Nodule formation in inoculated plants increased by 28%, 39% and 56% by applying 15, 30 and 45 kg P/ha, respectively compared to nodulation in the unfertilized plants whereas inoculation effect was greatest (80%) in the unfertilized treatments compared to 64%, 32% and 30% increases in the 15, 30 and 45 kg P/ha treatments, respectively (Figure 6). This suggests that Rhizobium inoculation influenced nodulation more positively than P fertilizer. [29] demonstrated from pot studies that nodulation and N₂ fixation of promiscuous soybean may be increased by inoculation with effective *Bradyrhizobia*. [30] also attributed the increase in nodulation to the competitive ability of *Bradyrhizobium* used, a result which was confirmed by [31].




Figure 6: Influence of Rhizobium inoculation and phosphorus fertilizer application rate on number of nodules per plant at full pod stage of soybean.

3.7 Nodule biomass

Nodule biomass production was enhanced significantly by inoculating soybean with Rhizobium inoculants whether the crop was fertilized with phosphorus or not (Figure 7). The effect of inoculation was, however, greater in the presence of P and also when higher rates of

P were applied but the highest percent increase (105%) in biomass due to inoculation occurred with P rate of 30 kg/ha. The other increases were 43%, 78% and 88% for 0 kg P, 15 kg P and 45 kg P per hectare, respectively. Figure 7 reveals that in un-inoculated plants the highest biomass of 1.6 g/plant was obtained at 45 kg P/ha application rate but a co-application of inoculants and only 15 kg P/ha was able to produce the same nodulation effect. This means that with respect to nodule biomass (which is an index of biological N fixation if all nodules are active), application of 250 g of inoculants (inoculation rate used in this study was equivalent to 250 g inoculants/ha) could make savings equivalent to 30 kg P (i. e. 381.7 kg Yaralegume) in this study. The positive effect of Rhizobium inoculation on nodulation has been confirmed by [32] who recorded a significant increase in nodulation by native soil *Rhizobium* population in single inoculations conducted on a legume. Phosphorus is also known to initiate nodules formation, increase the number of nodule primordial and is essential for the development and functioning of formed nodules [33] and [34]. Several researches have reported that the supply of phosphorus plays important roles in the establishment, growth and function of nodules [26], [35] and [36]. Both figures 6 and 7 show that the interactive application of phosphorus and Rhizobium inoculants had a greater impact on nodulation than either treatment alone.




Figure 7: Influence of Rhizobium inoculation and phosphorus fertilizer application rate on nodule biomass of soybean at full pod stage.

3.8 Pod production

The number and weight of pods produced by soybean in response to applications with P fertilizer and Rhizobium inoculants are shown in figures 8 and 9, respectively. Rhizobium inoculation remarkably increased both pod number and pod weight but whereas the increases due to inoculation were observed at all the P application rates in the former (Figure 8), there was no effect of inoculation on pod weight at 15 kg P/ha rate (Figure 9). Also, whether plants were inoculated or not, pod production generally increased with increasing rates of P application (Fig. 8 & 9) which agrees with [37] and [20] who reported high numbers of pods per plant in response to high doses of phosphorus. [38] also reported that a mixed inoculation improved the productivity of soybean when compared to the control treatment. Increased numbers of pods per plant in response to inoculation were also reported by [39] and [21]. The increases in pod weight at the different increasing rates of P plus Rhizobium inoculation may have been due to enhanced stimulation of pod filling in such treatments as confirmed by [21] when they reported of significant high numbers of seeds per pod when 100 kg P was applied in contrast to minimum numbers of seeds where no P was applied




Figure 8: Influence of Rhizobium inoculation and phosphorus fertilizer application rate on number of pods of soybean at full pod stage.



3.9 Pod weight

There were significant difference ($p<0.05$) among varied phosphorus rate, *Rhizobium* inoculants and the interaction between phosphorus rates and *Rhizobium* inoculants on pod weight (figure 9). Different rates of phosphorus fertilizer plus *Rhizobium* inoculant stimulated pod filling and thereby improved the pod weight as the phosphorus level progressively increased. Phosphorus at 45 kg P ha⁻¹+ *Rhizobium* inoculant recorded the highest pod weight as compared to phosphorus at 15 kg P/ha which recorded the least. The current study agrees with the findings of [40] and [21] who also significantly recorded the highest number of seeds per pod produced when 100 kg P ha⁻¹ was applied and minimum number was recorded where no phosphorus was applied. Similar results were obtained by [41] who reported of significant effects of phosphorus rates on harvest index of soybeans.




Figure 9: Influence of *Rhizobium* inoculation and phosphorus fertilizer application rate on pod weight of soybean at full pod stage.

3.10 Total grain yield

Similar to pod production, grain yield of soybean was significantly influenced by varied phosphorus rates, *Rhizobium* inoculation and their interaction (Figure 10). The grain yield increased significantly at every higher rate of P application from 0 to 45 kg/ha with or without inoculation. Phosphorus application at 45 kg P/ha plus *Rhizobium* inoculant recorded the highest grain yield of 1955 kg/ha compared to 897 kg grain per hectare in the unfertilized inoculated treatment which constitutes a yield increase of 118% (Figure 10). However, though inoculation enhanced grain yield at every P application rate, within the P application

regimes Rhizobium inoculation caused the highest percent grain yield increase (25%) at 15 kg P/ha. Also, these results are in agreement with [42] who concluded that phosphorus and inoculation induced a pronounced effect on grain yield. [39] also reported significant increases in grain yield with Rhizobium inoculation.




Figure 10: Influence of Rhizobium inoculation and phosphorus fertilizer application rates on grain yield of soybean.

Economic analysis

Information on costs and benefits of treatments is a prerequisite for adoption of technical innovation by farmers [43]. The studies assess the economic benefits of the treatments to help develop recommendation from the agronomic data. This enhances selection of the right combination of resources by farmers in the study area. The results in this study indicated that the inoculated treatments resulted in higher net benefits than the uninoculated treatments for all P fertilizer treatments (Table 1). The implications are that farmers will be better off inoculating soybeans seeds before planting as it will improve incomes of farmers through increased grain yields. The partial budget analysis showed that the application of 45 kg P/ha for both inoculated and uninoculated treatments yielded higher net benefits than all the treatments. The control (no fertilizer) treatment also produced the lowest net benefit for both the inoculated and un-inoculated treatments. This implies that farmers would be better off inoculating their soybean in combination with application of 45 kg P/ha as these increase soybean yields and thus increase farmers' income.

306

307

308

309

310 **Table 1:** Partial budget for inoculated and un-inoculated soybean seeds under four different
311 treatments

ITEM	TREATMENT							
	Inoculated				Un-inoculated			
	15 kg P/ha	30 kg P/ha	45kg P/ha	0 kg P/ha	15 kg P/ha	30 kg P/ha	45kg P/ha	0 kg P/ha
Average yield (kg/ha)	1230	1640	1955	897	984	1350	1580	738
Adjusted yield (kg/ha) ¹	1107	1476	1759	807	885	1215	1422	664
Gross field benefit (C) ²	2656	3542	4222	1937	2125	2916	3412	1594
Cost of Yaralegume (C/ha)	16	33	48	0	16	33	48	0
Cost of inoculation (C/ha)	15	15	15	15	0	0	0	0
Total cost that vary (C)	31	48	63	15	16	33	48	0
Net benefit (C)	2625	3494	4159	1922	2109	2883	3364	1594

312 ¹ Average yield adjusted 10% downwards; Farm gate price of soybean as at December 2012= GH

313 C2.4 per kg; Price of 50 kg of Yaralegume as at December, 2012 = GH C50.00 and price of inoculants

314 = GH C15.00/ha. 

315 **Conclusion and Recommendation**

316 Growth and development parameters of the soybean measured increased with increasing rates
317 of phosphorus fertilizer coupled with Rhizobium inoculation. Generally, plants that received
318 phosphorus fertilizer performed better than plants that were not fertilized as well as than
319 plants that received inoculation only. However, the positive effects of P fertilizer application
320 on growth and grain yield of soybean in this study were significantly boosted in the presence
321 of Rhizobium inoculants. The growth and grain yield enhancements observed in this
322 investigation may be attributed to an increased symbiotic relationship of rhizobia (bacteria)

with the roots of leguminous crops which reflected in increased nodulation resulting in the fixation of atmospheric nitrogen into the roots of soybean which was favored by increased P nutrition.

From the results of the economic analysis conducted, this study therefore recommends that farmers inoculate their soybean seeds with *Rhizobium* and apply phosphorus fertilizer at 45 kg P/ha to maximize their grain yield and their farm income.

References



1. Chung G, Singh RJ. Broadening the Genetic Base of Soybean: A Multidisciplinary Approach Critical Rev. in Pl. Sci. 2008; 27: 295–341pp.
2. Frankel FN. Soybean oil flavour stability. In: Erikson, d.r., pryde, E. H., Brekke, O. L., Mont, T. L., Falk, R. A. (Eds), Handbook of soy oil processing and Utilization American Soybean Association, St. Louis. 1980; 229-244 pp.
3. Singh RR, Rachie KO. Introduction to soybeans. In: soybean for the tropics: Research, production and Utilization. Sing, S. R., Rachie, K. O. and Dashiell, K. (eds), Chichester, John Wiley and Sons Publishing Ltd: (1987). 15-20pp.
4. Langyintuo AS, Lowenberg-D Boer JK, Faye M, Lambart D, IbroMoussa B. Cowpea supply and demand in West and Central Africa I Field Crops Research. 1996; 82: 215-231pp.
5. Mahamood J., Abayomi YA., Aduloju MO. Comparative growth and grain yield responses of soybean genotypes to phosphorous fertilizer application. Department of Agronomy, University of Ilorin, P. M. B. 1515, Ilorin, Nigeria. African Journal of Biotechnology. 2008; Vol. 8 (6), pp. 1030-1036, 20 March, 2009. Available online at <http://www.academicjournals.org/AJB>

- 348 6. Awuku KA. Agriculture and Environmental Studies. Rebl. Evans Brothers Ltd.
349 London. 1991; 344pp.
- 350 7. Richard JD, Gerub LJ, Anlgren HL. Crop protection, 5th end. Prentice-Hall Inc. Engle
351 wood cliffs New Jersey. 1984; pp.252-279
- 352 8. Issaka RN, Senayah JK, Adjei EO. Soil Resources of Saboba- Cheriponi District. An
353 assessment for Agricultural production West Africa journal of Applied Ecology.
354 2004;5: 114pp.
- 355 9. Carsky RJ, Singh BB, Oyewole R. Contribution of early-season cowpea to late-season
356 maize in the Savanna zone of West Africa. Biol. Agric. Hort. 2001;18: 303-315pp.
- 357 10. Beijiga G. Current status of food legume production and use of biological nitrogen in
358 Ethiopia. In: symbiotic nitrogen fixation prospects for enhanced application in
359 Tropical Agriculture, serraj, R. (Ed), oxford and IBH. Publishing CO. Pvt. Ltd. New
360 Delhi, India. 2004;263-26pp.
- 361 11. Wolde-Meskel E. Genetic diversity of *Rhizobia* in enhances Biological Nitrogen
362 Fixation (BNF) and soil fertility for sustainable agriculture. Ethiopian J. Biol. Sci.
363 2007;6: 77-95
- 364 12. Ellafi A, Gadalla MA, Galla YGM. Biofertilizers inaction; contribution of BNF in
365 sustainable agricultural ecosystems. Int. Sci. Res. J. 2011;3:108-116pp.
- 366 13. Savannah Agricultural Research Institute. Annual report 2008.
- 367 14. Lamprey S, Ahiabor BDK, Yeboah S and Osei D. Effect of Rhizobium Inoculants
368 and Reproductive Growth Stages on Shoot Biomass and Grain Yield of Soybean
369 (*Glycine max* (L.) Merril) (*In press*).

- 370 15. Gunarto L. *Rhizosphere* microbes: their roles and potential. J. Penelitian Dan
371 Pengembangan Pertanian. 2000;19: 39–48pp.
- 372 16. Kwari JD. Soil fertility status in some communities of southern Borno. Final report to
373 Prosab Project, Maiduguri, Nigeria. 2005; 21pp.
- 374 17. Uzoma AO, Osunde AO, Bala A Effect of phosphorus and *Rhizobium* inoculation on
375 the yield and yield components of cowpea breeding lines in Minna. 2006; In
376 Proceedings of 31st Annual Conference of Soil Science Society of Nigeria Zaria.
- 377 18. Rebafka FP, Ndunguru BJ, Marchner H. Single superphosphate depresses
378 molybdenum uptake and limits yield response to phosphorus in groundnut (*Arachis*
379 *hypogaeal*) grown on an acid. 2003;4: 55p.
- 380 19. Tomar SS, Singh R, Singh PS. Response of phosphorus, sulphur and *Rhizobium*
381 inoculation on growth, yield and quality of soybean. Prog. Agric. 2004;4 (1): 72-
382 73pp.
- 383 20. Rani BP. Response of soybean to nitrogen and phosphorus application in the black
384 soils of Krishna-Godavari zone of Andhra Pradesh. Annuals Agric. Res. 1999;20(3):
385 367-368pp
- 386 21. Hernandez M, Cuevas F. The effect of inoculating with *Arbuscular mycorrhiza* and
387 *Radyrhizobium* strains on soybean (*Glycine max (l.) Merrill*) crop development
388 cultivos-tropicales. 2003; 24 (2): 19-21pp.
- 389 22. Menaria BL, Singh RK, Nagar P, Singh P. Effect of nutrients and microbial
390 inoculants on growth and yield of soybean (*Glycine max L.*) .J. Soils and Crops.
391 2003;13(1): 14-17pp.
- 392 23. Tsveitkova GE, Georgiev GI. Effect of phosphorus nutrition on the nodulation,
393 nitrogen fixation and nutrient use efficiency of *Bradyrhizobium japonicum* soybean

- 394 (Glycine max L. merrill) symbiosis. Bulgarian Journal of Plant Physiology. 2003;331-
395 335pp.
- 396 24. Zafar M, Abbasi M, Rahim N, Khaliq A, Shaheen A, Jamil M, Shahid M. Influence of
397 integrated phosphorus supply and plant growth promoting rhizobacteria on growth,
398 nodulation, yield and nutrient uptake in Phaseolus vulgaris. Afr. J. Biotechnol. (2011);
399 10:16793-16807.
- 400 25. Chiera JM, Thomas JF, Rufty TW. Growth and localized energy status in phosphorus-
401 stressed soybean. J. Plant Nutr. 2004; 27: 1875-1890pp.
- 402 26. Abbasi MK, Majeed A, Sadiq A, Khan SR. Application of *Bradyrhizobium japonicum*
403 and phosphorus fertilization improved growth, yield and nodulation of soybean in the
404 sub-humid hilly region of Azad Jammu and Kashmir, Pakistan. Plant Prod. Sci. 2008;
405 58: 368–376pp.
- 406 27. Lamptey S, Ahiabor BDK, Yeboah S, Asamoah C. Response of soybean (Glycine
407 max) to Rhizobial inoculation and phosphorus application. Journal of Experimental
408 Biology and Agricultural Sciences. 2014;2(1):72-77.
- 409 28. Asia M, Safder A, Shahzad SI. Effect of Phosphorus and potassium on yield and
410 nitrogen fixation by Mash bean. Sarhad J. Agric. 2005; 21(4): 667-670pp
- 411 29. Kumaga FK, Etu-Bonde K. Response of two promiscuous soybean genotypes to
412 Bradyrhizobial inoculation in two Ghanaian soils. J. Ghana Sci. Assoc. 2002;vol.2:
413 99–104pp
- 414 30. Okereke GU, Onochie CC, Onukwo, AU , Onyeagba E, Ekejindu, GO. Response of
415 introduced Bradyrhizobium strains infecting a promiscuous soybean Cultivar. World
416 J. Microbiol. Biotech. 2000;16: 43–8

- 417 31. Fening JO, Danso SKA. Variation in symbiotic effectiveness of cowpea
418 *Bradyrhizobia* indigenous to Ghanaian soils. Appl. Soil Ecol. 2002); 21: 23–9pp.
- 419 32. Elkoca E, Turan M, Donmez MF. Effects of Single, dual and triple Inoculations with
420 *Bacillus subtilis*, *Bacillus megaterium* and *Rhizobium leguminosarum* Bv. *Phaseolus*
421 nodulation, nutrient uptake, yield and yield parameters of common bean (*Phaseolus*
422 *vulgaris* L. cv. ‘Elkoca-05’). J. Plant Nutr. 2010; 33: 2104–2119pp
- 423 33. Waluyo SH, Lie TA, Mannetje L. Effect of phosphate on nodule primordia of soybean
424 (*Glycine max*) in acid soils in rhizotron experiments. Ind. J. Agric. Sci. 2004;5: 27-
425 44pp
- 426 34. Tagoe SO, Horiuchi T, Matsui T. Effects of carbonized and dried chicken manures on
427 the growth, yield, and N content of soybean. Plant Soil. 2008; 306: 211-220pp.
- 428 35. FAO. Legume inoculants and their use. 1984; A pocket manual jointly prepared by
429 NIFTAL project, USA.
- 430 36. Shu-Jie MS, Yun-Fa, Xiao-Zeng H, An M. Nodule Formation and development in
431 soybeans (*Glycine max* L.) in response to phosphorus supply in solution culture.
432 Pedosphere. 2007;17: 36–43pp
- 433 37. Mohan KK, Rao LM. Effect of phosphorus and molybdenum on growth, nitrogen
434 fixation and yield of soybean (*Glycine max* L.). Angrau. J. Research. 1997;25(4): 77-
435 80pp.
- 436 38. Konde, VB, Deshmukh RB. Response of chickpea varieties to *Rhizobium* and VAM
437 inoculation. J. Maharashtra Agri. Univ. 1996; 21: 304–6 (Field Crop Abstracts, 51(1):
438 426; 1999)
- 439

- 440 39. Sable S, Sontakey PY, Tagade R, Deotale RD, Manapure P. Effect of *Rhizobium* and
441 molybdenum on quality aspects, yield and yield contributing characters of soybean. J.
442 Soils Crops. 1998;8 (2): 157-159pp.
- 443 40. Patel RS, Patel ZG. Effect of farmyard manure, nitrogen, phosphorus and Rhizobium
444 inoculation on the growth and yield of gram (*Cicer arietinum* L.). Ann. Agri. Res.
445 1991;12: 200–02.
- 446 41. Malik M F A, Qureshi AS, Ashraf M, Ghafoor A. Genetic variability of the main
447 yield related characters in soybean. Intl. J. Agric. and Biol. 2006; 8(6): 815-819pp.
- 448 42. Stefanescu M, Palanciuc V. Efficiency of bacterial inoculation and mineral nitrogen
449 and phosphorus fertilization in rainfed soybean. Roma. Agric. Res. 2000; 13-14: 75-
450 83pp.
- 451 43. Das A, Biswas M, Mandal N. An economic analysis of stevia (*Stevia rebaudiana*
452 Bert.) cultivation through stem cutting and tissue culture propagule in India Trends
453 Agric. Econ. 2010; 3: 216-222
454
455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498