

Potassium Nutrition and Translocation in Cassava (*Manihot esculenta* Crantz) intercropped with Soybean

Abstract

Experiment to investigate the effect of potassium nutrition and translocation in cassava (*Manihot esculenta* Crantz) intercropped with soybean (*Glycine max* (L) Merrill) was conducted at University of Nigeria, Nsukka in a derived savannah location of South Eastern Nigeria. The effect of application of potassium fertilizer on two varieties of cassava intercropped with six cultivars of soybean was studied. Cassava tuber yield was significantly ($P<0.05$) affected by their varieties, fertilizer rate and cropping system. An increase in the supply of K^+ by fertilization (N_0K_{50} and $N_{45}K_{50}$) increased K^+ accumulation in source leaves and caused increase in cassava tuber yield. The highest cassava tuber yield (42.0 t.ha^{-1}) was obtained with the fertilizer rate of N_0K_{50} x NR 8230 intercropped with TGX 1894-3E followed by the yield (40.5 t.ha^{-1}) of the same cassava cultivar (NR 8230) x $N_{45}K_{50}$ intercropped with Samsoy-2.

Key words: Cassava, soybean, potassium, nutrition, intercrop, translocation

27 INTRODUCTION

28 Potassium is an element essential to plant for the formation and transfer of
 29 carbohydrate in photosynthesis and also for protein synthesis. It is particularly important for
 30 fruits, leaves and stems and is needed to strengthen the plant's structure. Potassium promotes
 31 high crop yields, particularly in root and tuber crops. Potassium helps to direct free nutrients
 32 such as carbon, hydrogen and oxygen out of the atmosphere and into the plant. Various
 33 studies provide evidence that K^+ nutrition promotes the translocation of assimilates in plants.
 34 Without the activity of potassium, photosynthesis would be severely restricted and plant
 35 would struggle to make starches, sugars, proteins, vitamins, enzymes and cellulose. Net
 36 carbon exchange in plant increases as a result of increased K^+ fertilization (Amir and
 37 Reinhold, 1971). Potassium aids in helping the plant through adverse climate in that, when
 38 potassium is out of balance, plant stress is very high (Umeh and Mbah, 2010a). When too
 39 much potassium is used, synergistic reaction occurs between calcium and magnesium,
 40 thereby changing the soil pH rapidly. At high pH, up to pH 6, potassium mobility slows
 41 down, and as soil reaches 7.0, mobility is severely hindered. Potassium reserve in soil is
 42 largely dependent on the type of soil mineral present in the soil. Soil potassium may be
 43 classified according to its availability to plants and falls into three categories: the inert reserve
 44 or slowly available K, the dynamic reserve and the readily available reserve (Defoer *et al.*,
 45 2000). The ratio of potassium exported through crop removal: exchangeable K indicates how
 46 long a farmer can afford to continue to extract K without adding potassium fertilizer.

47 Brennan and Jayasena (2007) reported that increasing applications of potassium
 48 fertilizer to barley crops grown on deficient sandy soils increased grain yield and decreased
 49 some foliar diseases. At the international Potash Institute (IPI), Srinivasarao *et al.* (2007)
 50 categorised soils based on potassium reserve and production systems. They observed positive
 51 correlations with exchangeable and non-exchangeable K in all soil types indicating the

dynamic equilibrium between the two soil- K^+ fractions. Increasing the K^+ available to the roots of cassava caused an increase to the delivery of K^+ to the leaves of the plant and possible increase in the cassava tuber yield (Kang and Okeke, 1983). Haeder and Mengel (1972) found that increasing K^+ supplies to the roots caused increased translocation of carbon in tomato fruits. Earlier studies (Kang and Okeke, 1983) examined the responses of cassava varieties to N and K. This study was conducted to investigate the effect of potassium nutrition and translocation in cassava intercropped with soybean.

MATERIALS AND METHODS

The experiment was conducted at University of Nigeria, Nsukka farm which is located at latitude $06^{\circ} 52''N$ and longitude $07^{\circ} 24''E$ and at 447m above the sea level, between August 2008 and July 2009. Experimental design was split plot in randomised complete block design, having two factors; fertilizer rates and cropping systems. Six varieties of soybean of three maturity classes (early, medium and late maturing) were intercropped with two varieties of cassava. Four fertilizer rates were randomised in the main plots, while twenty cropping systems comprising six sole soybean, two sole cassava and twelve cassava/soybean intercrop were randomised in the sub- plots.

Cassava varieties used were TMS 30572 and NR 8230. The six varieties of soybean used were TGX 1448-2E and Samsoy-2, (early maturing varieties), TGX 1894-3E and TGX 1805-31F, (medium maturing varieties), TGX 1889-12F and TGX 1864-17F, (late maturing varieties). The four fertilizer rates were: N_0K_0 , N_0K_{50} , $N_{45}K_0$ and $N_{45}K_{50}$ kg. ha^{-1} , where N and K were nitrogen and potassium respectively, subscripts 0, 45 and 50 were levels of N and K kg. ha^{-1} respectively. The nitrogen source was Urea and potassium source was muriate of potash. A uniform application of 30 kg. ha^{-1} of P as single supper phosphate was applied to all plots.

Plot size was 4.0 m X 3.0 m., containing 4 ridges at 1.0 m spacing. Soybean was planted on both sides of the ridge at a plant distance of 10.0 cm while cassava was planted at a plant distance of 0.75 m. on the crest of the ridges. Both cassava cuttings (20 cm long) and soybean seeds were planted at the same time. Cassava plant population was 13,333.33 plants ha⁻¹ while soybean was thinned down to plant population of 200,000 plants ha⁻¹. Planting was done in early August and weed was controlled manually. Cassava tuber yield, shoot dry matter and potassium content were determined at 12 months after planting. Uptake of K was determined as accumulation of K in cassava leaf and cassava tuber using atomic absorption spectrophotometric method.

Key: N₀K₀ = 0kg N ha⁻¹ and 0kg K ha⁻¹; N₀K₅₀ = 0kg N ha⁻¹ and 50kg K ha⁻¹;
N₄₅K₀ = 45kg N ha⁻¹ and 0kg K ha⁻¹; N₄₅K₅₀ = 45kg N ha⁻¹ and 50kg K ha⁻¹

RESULTS

Effect of potassium rate and cropping system on cassava tuber yield (t.ha⁻¹) at 12 MAP was presented in table 1. Application of N₀K₅₀ fertilizer rate out-yielded all other fertilizer rates (30.7 t.ha⁻¹), which was 204% higher than the yield at N₀K₀ (12.6 t.ha⁻¹), 108% higher than at N₄₅K₀, (28.4 t.ha⁻¹) and at N₄₅K₅₀ (28.0 t.ha⁻¹). The tuber yield at N₄₅K₀ and N₄₅K₅₀ did not differ significantly. There were no significant yield differences between the two cassava cultivars as sole crop. However, at intercrop, NR 8230 intercropped with TGX 1894-3E produced significantly (P < 0.05) the highest tuber yield (30.8 t.ha⁻¹) which was 57% higher than the sole crop (19.03 t.ha⁻¹).

All intercrop cassava significantly (P < 0.05) out yielded sole cassava except NR 8230 intercropped with TGX 1448-2E (17.9 t. ha⁻¹) which significantly yielded lower than both cassava sole crops (19.0 and 18.6 t.ha⁻¹ respectively). Generally, NR 8230 significantly out

yielded TMS 30572 when intercropped with all varieties of soybean except with TGX 1448-2E, where the yield of TMS 30572 was 65.8% higher than NR 8230.

The interaction of sole cassava x all fertilizer rates had generally lower tuber yields than when intercropped except at $N_{45}K_0$ and $N_{45}K_{50}$ x NR 8230 intercropped with TGX 1448-2E, $N_{45}K_{50}$ x TMS 30572 intercropped with Samsoy-2 and TMS 30572 intercropped with TGX 1894-3E. The fertilizer rate of N_0K_{50} x NR 8230 intercropped with TGX 1894-3E gave significantly ($P < 0.05$) the highest cassava yield of 42.0 t.ha^{-1} followed by the yield (40.5 t.ha^{-1}) of the same cassava cultivar (NR 8230) x $N_{45}K_{50}$ intercropped with Samsoy-2. Medium maturing varieties, TGX 1894-3E and TGX 1805-31F, intercropped with the two cassava varieties gave a more close consistent tuber yield at N_0K_{50} (42.0, 38.9, 38.2 and 38.7 t.ha^{-1}) which were significantly ($P < 0.05$) higher than all other soybean varieties x fertilizer rates both at sole and at intercrop. Cassava tuber yield was lowest at sole NR 8230 (4.2 t. ha^{-1}).

Effects of potassium rate and cropping system on cassava leaf-K at 12 MAP were presented in table 2. Cassava leaf-K differed significantly ($P < 0.05$) with potassium rates. Fertilizer rates of N_0K_{50} , $N_{45}K_0$ and $N_{45}K_{50}$ were 30.26, 11.69 and 29.21% higher than N_0K_0 respectively. Leaf-K at N_0K_{50} (2.71%) was significantly higher than other fertilizer rates, while leaf-K at N_0K_0 (1.89%) was the lowest.

Sole crops did not differ significantly among themselves. Intercrops of TGX 1448-2E with either varieties of cassava were similar (2.33%), while the intercrops of NR 8230 (2.45%) was significantly ($P < 0.05$) higher than TMS 30572 intercropped with Samsoy-2 (2.40%), (early maturing variety) TGX 1894-3E (2.55%) and TGX 1805-31F (2.30%), (medium maturing varieties) and at late maturing varieties, intercrops of TMS 30572 with both soybean varieties out yielded intercrops of NR 8230. At early maturing variety, highest leaf-K (2.45%) was obtained at NR 8230 intercropped with Samsoy-2 while at medium

maturing varieties, NR 8230 intercropped with TGX 1894-3E gave the highest leaf-K (2.55%). At late maturing variety, the highest leaf-K was obtained with TMS 30572 intercropped with TGX 1864-17F (2.38%).

DISCUSSION

The importance of inclusion of potassium was shown where the highest parameters were obtained at N_0K_{50} and $N_{45}K_{50}$ fertilizer rates. Increasing the K^+ available to the plants (N_0K_{50} and $N_{45}K_{50}$ fertilizer rates) caused significant increase in the cassava tuber yield at more than other fertilizer rates (N_0K_0 and $N_{45}K_0$) where there was no K^+ in the fertilizer. Tarrace and Gieiger (1982) made similar observation and reported that increased K^+ nutrition on sugar beet plants promoted the translocation of products of photosynthesis, affect the metabolic conversion of sucrose in the sink tissues and resulted in net carbon increase and C/N ratio. An increase in the supply of K^+ in the plant root zone by fertilization (N_0K_{50} and $N_{45}K_{50}$) increased K^+ accumulation in source leaves and caused increase in cassava tuber yield. The amount of K^+ in cassava leaf was between 1.5 – 3.5%, while tuber yield was 2.0 – 45.0 t.ha⁻¹. The role of K^+ as a cofactor in synthetic metabolism has been well established (Rains 1976). The increased K^+ at the sink region may enhance the conversion of sucrose to synthesized metabolites and thus promote translocation into those sinks with enhanced K^+ . There was also enhancement of synthesis of protein, lipids and cell wall in the growing plant resulting from K^+ supply as reported by Mengel and Haeder (1977). This observation explains why nitrogen fertilizer or biological nitrogen fixation (BNF) alone did not achieve the highest growth and yield of cassava. Also, it has been reported that the imbalance between exchangeable K and other soil minerals, notably calcium and magnesium, may result in total crop failure (Howeler and Candavid 1983). There was significant ($P < 0.05$) interaction between potassium rate and cropping system. Highest leaf-K (2.9%) was obtained at $N_{45}K_{50}$ x

TMS 30572 intercropped with TGX 1894-3E, TMS 30572 intercropped TGX 1864-17F, N₀K₅₀ x NR 8230 intercropped with Samsoy-2 and TMS 30572 intercropped with TGX 1889-12F. Lowest leaf-K was obtained at N₀K₀ x all the varieties and cropping system. Highest leaf-K was obtained at N₀K₅₀ and N₄₅K₅₀ fertilizer rates. Without K fertilization (N₀K₀ and N₄₅K₀), residual soil potassium was very low. Howelar and Candavid (1983) reported also that even if the stems and leaves of cassava were returned to the soil, there would still be a considerable net loss of K from the soil because much K is removed by the cassava roots and fresh tuber. CIAT (1981) reported that without adequate K fertilization in cassava, there was yield decline.

CONCLUSION

The highest cassava tuber yield (42.0 t.ha⁻¹) was obtained with the fertilizer rate of N₀K₅₀ x NR 8230 intercropped with TGX 1894-3E. Cassava leaf-K at N₀K₅₀ was significantly higher than other fertilizer rates, while leaf-K at N₀K₀ was the lowest.

REFERENCES

- Amir, S. And Reinhold, L. (1971). Interaction between potassium deficiency and light 14C-sucrose translocation in bean plants. *Physiol Plant* 24:226-231
- Brennan, R.F. and Jayasena, K.W. (2007). Increasing application of Potassium fertilizer barley crops grown on deficient sandy soils increased grain yields while decreasing some foliar diseases. *American Journal of Agricultural Research* 58:680-689.
- CIAT, (Centro International de Agricultural Tropical) (1981). Annual Report-Cassava production system 53-54.
- Defoer, T., Budelman, A., Toulmin., and Carter, S.E. (2000). Building Common Knowledge. Participatory learning and action Research (part 1). In: Defoer, T. Budelman A (eds). *Managing Soil Fertility in the Tropics. A Resource Guide for participatory learning and action research*. Amsterdam. The Netherlands: Royal Tropical Institute.
- Kang, B.T. and Okeke, J.E. (1983). Nitrogen and potassium responses of two cassava

184 varieties grown on an alfisol in Southern Nigeria.
 185 [http://www.istr.org/sites/default/files/files/symposium/internal/1983_](http://www.istr.org/sites/default/files/files/symposium/internal/1983_peru/6th_symposium_proceedings_0040_section_3_2)
 186 [peru/6th_symposium_proceedings_0040_section_3_2](http://www.istr.org/sites/default/files/files/symposium/internal/1983_peru/6th_symposium_proceedings_0040_section_3_2)
 187

188 Mengel, K. and Haeder H-E (1977). Effect of potassium supply on the rates of phloem sap
 189 exudation and the composition of phloem sap of *Ricinus communis*. Plant physiology
 190 59:282-284.
 191

192 Haeder, H-E and Mengel, K. (1972). Translocation and respiration of assimilates in tomato
 193 plants as influenced by K nutrition. Z Pflanzenernahr Bodenkd 131: 139-148.
 194

195 Howeler, R.H. and Candavid, L.F. (1983). Accumulation and Distribution of Dry matter and
 196 Nutrients during a 12 month Growth cycle of Cassava. Field Crops Research.
 197 7, 123- 139.
 198

199 Srinivasarao, Ch., Vitta, K.P.R., Tiwari, K.N. Gajbliye, P.N. and Kundu, S. (2007).
 200 Categorization of Soil based on potassium reserves and production Systems.
 201 Implication in K management. Australian Journal of Soil Research 45: 436-447.
 202

203 Terrance, R. C. and Geiger, R. D. (1982). Potassium Nutrition and Translocation in sugar
 204 beet. Plant Physiology, 70: 169-172.
 205

206 Umeh, S. I and Mbah, B. N. (2010a). Soybean contribution to nitrogen balance in a cassava
 207 (*Manihot esculenta* Crantz) based cropping system. Agro-Science Journal 9(3):147-153
 208

209 Umeh, S. I and Mbah, B. N. (2010b). Measuring the benefits of biological nitrogen
 210 Fixation of soybean (*Glycine max* (L) Merrill) in cassava (*Manihot esculenta* Crantz
 211 and soybean intercrop. African Journal of Agricultural Research 5(24): 3354-3359.
 212

213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232

233 Table 1: Effects of fertilizer rate and cropping system on cassava tuber yield (t.ha⁻¹)
234 at 12 MAP

Cropping systems (C)	Fertilizer rates (F)				Mean
	N ₀ K ₀	N ₀ K ₅₀	N ₄₅ K ₀	N ₄₅ K ₅₀	
NR 8230 Sole	4.2	22.1	24.2	25.6	19.0
TMS 30572 Sole	6.8	20.0	21.4	26.1	18.6
NR 8230/TGX1448-2E	7.1	24.2	17.8	12.4	17.9
TMS30572/TGX1448-2E	13.1	23.5	36.5	35.6	27.2
NR 8230/Samsoy-2	10.5	33.8	30.1	40.5	28.7
TMS 30572/Samsoy-2	12.0	31.5	32.9	16.7	23.3
NR 8230/TGX 1894-3E	15.7	42.0	30.6	34.7	30.8
TMS30572/TGX1894-3E	11.5	38.9	32.6	22.0	26.3
NR 8230/TGX 1805-31F	9.7	38.2	30.4	32.8	27.8
TMS30572/TGX1805-31F	11.8	38.7	27.8	27.8	26.4
NR8230/TGX1889-12F	14.6	35.5	32.6	32.6	28.8
TMS30572/TGX188912F	21.8	32.4	31.6	23.5	27.3
NR 8230/TGX 1864-17F	18.2	25.8	25.8	27.4	24.3
TMS30572/TGX186417F	19.6	23.6	23.6	25.3	21.7
Mean	12.6	30.7	28.4	28.0	
LSD _{0.05} for comparing two Fertilizer (F) means =0.51					
LSD _{0.05} for comparing two Cropping system (C) means = 0.58					
LSD _{0.05} for comparing two (F x C) interaction means = 1.14					

235
236
237
238
239
240
241
242
243
244
245
246

247

248

249

Table 2: Effects of fertilizer rate and cropping system on cassava leaf-K (%) at 12MAP

Cropping systems (C)	Fertilizer rates (F)				Mean
	N ₀ K ₀	N ₀ K ₅₀	N ₄₅ K ₀	N ₄₅ K ₅₀	
NR 8230 Sole	1.9	2.6	2.1	2.8	2.35
TMS 30572 Sole	1.9	2.7	2.1	2.6	2.33
NR 8230/TGX1448-2E	1.8	2.8	2.2	2.5	2.33
TMS30572/TGX1448-2E	1.9	2.8	2.1	2.5	2.33
NR 8230/Samsoy-2	1.9	2.9	2.2	2.8	2.45
TMS 30572/Samsoy-2	2.0	2.7	2.1	2.8	2.40
NR 8230/TGX 1894-3E	2.1	3.2	2.1	2.8	2.55
TMS30572/TGX1894-3E	1.9	2.6	2.2	2.9	2.40
NR 8230/TGX 1805-31F	1.8	2.5	2.3	2.6	2.30
TMS30572/TGX1805-31F	1.8	2.5	2.2	2.4	2.23
NR8230/TGX1889-12F	1.9	2.6	2.1	2.5	2.28
TMS30572/TGX188912F	1.8	2.9	2.1	2.6	2.33
NR 8230/TGX 1864-17F	1.8	2.6	2.1	2.8	2.33
TMS30572/TGX186417F	1.9	2.6	2.1	2.9	2.38
Mean	1.89	2.71	2.14	2.68	
LSD _{0.05} for comparing two Fertilizer (F) means = 0.01					
LSD _{0.05} for comparing two Cropping system (C) means = 0.02					
LSD _{0.05} for comparing two (F x C) interaction means = 0.04					

250