

Research Paper

EFFECTS OF LOW PROTEIN DIETS
SUPPLEMENTED WITH HIGH AMINO ACIDS
(METHIONINE AND LYSINE) ON
PERFORMANCE OF BROILERS.

8 **ABSTRACT ALL CORRECTIONS HAVE BEEN INSERTED**

9

Aims: These studies were to evaluate the effects of high level methionine and lysine supplementation on the performance and carcass characteristic of broilers fed low crude protein diets.

Study design The experiments employed a complete randomized design, all data generated were subjected to analysis of variance.

Place and Duration of Study: The experiment was carried out at the Teaching and Research Farm of the University of Ibadan between April and July 2011.

Methodology: Two hundred day old unsexed broiler chicks were used in a six week feeding trials. For each study four experimental diets were formulated such that they contained methionine at four dietary levels of 0, .35,.45 and .55% while the other four set contained lysine at four dietary levels of 0, .60, .70 and.80% respectively

Results: Results indicated that broilers fed diet1 with normal protein level without methionine and lysine were higher (($P=.05$)) in body weight gain compared with others. Feed conversion also had the same statistical trend. It was indicated that level of crude protein, lysine and methionine had significant effect on body weight gains. It was concluded that CP levels in the diets of the broilers could be reduced from 24.25 to 10.60% provided their methionine and lysine levels are higher than NRC recommended levels. This supplementation increased monetary losses due to low feed intake, reduced body weight gain and suboptimal feed conversions of the experimental animals.

Conclusion: The aim of the poultry farmer is to achieve optimum production with least inputs. From these studies, it can be concluded that crude protein content of broiler diet reduced from 24.25 to 10.60% and augmented with either methionine between .35 and .45 and Lysine between .60 and .70%. This supplementation was found to increase monetary losses due to low feed intake, reduced body weight gain and suboptimal feed conversions of the experimental birds.

Keywords: [Lysine Methionine,low protein, broiler }

10

11

12

13

INTRODUCTION

Poultry in one form or the other are kept in most areas of the world. There are fewer religious or social taboos associated with them than there are with pigs, rabbit and cattle thus products from poultry provide an acceptable form of animal proteins for most people throughout the world with the exception of strict vegetarians and Vegans [1].

With the increasing human population the consumption of meat is increasing in the world and intensive animal production has many more challenges to solve including environmental pollution and animal welfare [2].

The prevailing scarcity and cost of conventional plant protein supplements such as full fat soybean meal and groundnut cake have resulted in the recent high cost of commercial feeds and poultry products in Nigeria [3]. This major constraint necessitate intensified research on the possibilities of increasing the levels of amino acids and reducing the crude protein in the diets of poultry. Schutte and Pack [4] reported that in practical poultry diets, methionine is the first limiting amino acid followed by lysine. Although the advantages of the digestible amino acids are recognized, diet formulation on the basis of total amino acid content is still widely practiced [5]. Widespread commercial production of supplemental amino acids and further redefining of swine and poultry amino acid requirements have provided nutritionists with the option of reducing dietary CP without compromising performance levels. Therefore supplementation of methionine and lysine to practical poultry diets provides a means of increasing the efficiency of protein utilization and as a result nitrogen excretion will be reduced. In the past, swine and poultry diets were formulated using a specified crude protein level that was proven to provide an acceptable level of production. However widespread commercial production of supplemental amino acids and further redefining of monogastrics amino acid requirements have provided nutritionists with the option of reducing dietary crude protein without comprising performance levels [6]. The crude protein and amino acid status of a diet influence the carcass composition of broilers with increased carcass protein and reduced carcass fat accompanying increase in dietary protein or essential amino acid content [7]. The positive benefits of using supplemental amino acids have been reported to include decreased nitrogen in manure, decreased ammonia emission into the air, decreased water consumption by the experimental birds and decreased manure volume up to 55% [6]. When the CP is lowered, the ability of the diet to supply essential amino acids in a ratio that the birds require is impaired. The use of supplemental amino acids allows nutritionists to use a wide range of alternative ingredients in the diet that otherwise may not be economically or nutritionally practicable. The aim of this study was to

investigate the effects of methionine and lysine in low crude protein diets on performance and carcass characteristics of broilers.

MATERIAL AND METHODS

The experiment was carried out at the Teaching and Research Farm of the University of Ibadan. Maize and cassava flour were the main energy source while soybean was the main protein source. For each experiment, four diets were formulated. Diet1 which served as the control contained 24.25% crude protein while diets T2,T3 and T4 had 19.7, 15.15 and 10.6% respectively. The first set of Methionine supplemented diets 2,3 and 4 had recommended level of lysine with varying levels of methionine while the other set contained recommended level of methionine with varying levels of lysine (Table 1).

One hundred and ninety two day old broilers of Anak strain, bought from a reputable farm at Ibadan were used for these feeding trials. The broilers were raised on deep litter with conventional feed during the first week to allow for acclimatization, they were weighed on the eighth day and then randomly assigned to the experimental diets in such a way that each dietary treatment had 24 birds in three replicates of eight birds each. The experimental feed and water were supplied *ad libitum*. Routine vaccination and necessary medications were administered as when necessary. Records of feed intake were taken daily while body weight gain and mortality were taken on a weekly basis.

At the last week of the experiment, an indigestible marker Titanium dioxide (TiO_2) was included in the diets to monitor digestibility of the feed. Four birds per replicate were sacrificed and ilea digesta were collected in different bowls which were freeze-dried and analyzed. The sacrificed birds were scalded in moderately hot water and defeathered. They were eviscerated by carefully removing the gastro intestinal tracts. The weight of the cut up parts precisely the drumstick and breast muscle were taken.

2.1 Chemical analysis

Proximate analysis of the experimental diets and ilea digesta were carried out using AOAC methods AOAC [8].

2.2 Statistical Analysis

The experiments employed a complete randomized design, all data generated were subjected to analysis of variance and where significant differences occurred, the means were separated using Duncan multiple range test. (Which statistical package did the author use?)

Comment [AS1]:

Comment [AS2]:

Comment [AS3]:

3. RESULTS AND DISCUSSION

The gross composition of the experimental diets are as shown in Table 1 The first four sets of diets are designated as methionine diet where the CP varied from 10.60 to 24.25% and methionine inclusion from .25 to.50 while the other four sets had lysine inclusion of .50 to .80. The feed intake of broilers on control diets were significantly higher ($P \leq .05$) compared with lysine and methionine supplemented diets. It implied low level of crude protein with either high methionine or lysine depressed feed intake of broilers. The feed intake of birds on diets T2 T3 and T4 were similar in methionine supplemented diets while T2 and T3 were similar and higher than T4 in lysine supplemented diets (Table 2). The body weight gain also had a similar statistical trend as the feed intake. Those fed control diets had significantly ($P \leq .05$) higher weight than others fed dietary supplementation of Methionine and lysine. However there were no significant differences ($P \geq .05$) among the dietary treatments considering the feed conversion ratio. From these trials, it is apparent that in the methionine supplemented diets, feed intake in treatments T2, T3 and T4 were similar (2.43kg, 2.38kg and 2.32kg respectively). This is at variance with the reports of [9], [10], [4], [11] that methionine levels above NRC recommendations improves body weight [12], feed efficiency and breast meat yield.

Table 1. Gross composition of the Experimental diet

Ingredients(%)	Methionine supplemented Diets				Lysine supplemented Diets			
	T1	T2	T3	T4	T1	T2	T3	T4
Maize	54.50	54.4	54.3	54.2	54.50	54.4	54.3	54.2
Soybean meal	40	30	20	10	40	30	20	10
Cassava flour	0	10	20	30	0	10	20	30
Methionine	0.25	0.35	0.45	0.5	0.25	0.25	0.25	0.25
Lysine	0.50	0.50	0.50	0.50	0.50	0.60	0.70	0.80
T1O ₂	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Bone meal	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50
Oyster shell	1.70	1.70	1.70	1.70	1.70	1.70	1.70	1.70
Salt	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
*Premix	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Crude protein	24.25	19.70	15.15	10.60	24.25	19.70	15.15	10.60
%								
ME kcal	2821.8	2694.77	2970.42	3044.98	2821.8	2894.77	2970.42	3044.9
Meth (%)	0.60	0.64	0.67	0.71	0.60	0.54	0.47	0.41

Lysine (%) 1.84 1.54 1.24 0.93 1.84 1.64 1.44 1.23

*2.5 kg Premix used supplied vitamin A, 12,500,000iu; vitamin D, 2500000iu; vitamin E,40,000mg; vitamin K3,2000mg;vitamin B1, 3000mg; vitamin B2 5500mg Niacin, 55000mg;calcium pantothenate, 11500mg vitamin B6, 5000mg; vitamin B12, 25mg; Folic acid,1000mg;biotin,80mg;choline chloride 500000mg;manganese, 120000mg; Iron100,000mg;zinc,80000mg;copper 8500mg;iodine 1500mg cobalt 3000mg;selenium 120mg and anti-oxidant 120,000mg.

Table 2 Performance and carcass characteristics of broiler fed lysine and methionine supplemented diets

Parameters Kg/bird	METHIONINE					LYSINE				
	T1	T2	T3	T4	SEM	T1	T2	T3	T4	SEM
Feed intake	2.75 ^a	2.43 ^b	2.38 ^b	2.32 ^b	0.03	2.67 ^a	2.41 ^a	2.32 ^b	2.01 ^c	0.02
Weight gain	1.18 ^a	1.03 ^{ab}	0.88 ^b	0.85 ^b	0.01	1.06 ^a	0.93 ^b	0.87 ^b	0.55 ^c	0.00
Feed conversion ratio	2.33	2.36	2.70	2.73		2.52	2.60	2.67	3.65	
Drumstick	0.24 ^a	0.18 ^b	0.20 ^{ab}	0.19 ^{ab}	0.003	0.28 ^a	0.20 ^{ab}	0.21 ^{ab}	0.15 ^b	0.00
Breast	0.19 ^a	0.14 ^a	0.18 ^a	0.16 ^a	0.0009	0.19 ^a	0.15 ^{ab}	0.17 ^{ab}	0.13 ^b	0.00

abc-means along the same row with different superscripts differs significantly ($P < 0.05$)

Feed intake values across the methionine and lysine diets decreased as the amino acid inclusion level increased, this was contrary to the findings of [13] who reported increased feed consumption in low CP diet supplemented with lysine in broiler. This however agrees with [14] that broiler chickens given a very high dietary concentration of a single amino acid reduce their voluntary feed intake, consequently reducing growth rate and feed conversion efficiency. The decreased feed intake of the broilers fed low crude protein supplemented with high levels of methionine and lysine could probably be due to reason adduced by [15] that high methionine and lysine diets could result in amino acid imbalance. It was further reported that amino acid imbalance precipitates ill effects by reducing feed intake. The reduced feed intake coupled with the imbalance in the ratio of methionine and lysine may have contributed to the significantly ($P \leq 0.05$) lowered body weight gain recorded by broilers across the diets. Possibly the imbalance had a deleterious effect on the broilers. This confirmed the assertion that feed intake is influenced by dietary crude protein and amino acid levels[16,17]. This is in agreement with the report of [18] who reported that broiler

chicks receiving a low protein starter diet had inferior body weights and feed conversion ratio at four weeks of age compared with those receiving the required amount. This is contrary to [19] who reported no adverse effect on feed intake of growing broilers fed low crude protein with supplemental methionine. The digestibility result of this study had been reported elsewhere. The insignificant trend of feed conversion efficiency across the diets is an indication that low crude protein diet supplemented with methionine and lysine can be fed to broilers. This could possibly be due to reason adduced by [20] who opined that greater growth efficiency is expected when protein deposition is maximized and the ratio of protein synthesis to protein deposition is minimized. In the view of Reeds [21] it could imply that increase in nutrient intake above those required for nitrogen equilibrium are moderated by increase in both synthesis and degradation. Therefore inclusion levels of lysine and methionine in these trials above those required for optimal growth increase protein synthesis.

These results indicated that reduction up to 10.6% in the CP level of the broiler diets had no effect on the feed conversion ratio during the experimental period of eight weeks because the diets were supplemented with high methionine and lysine which was similar to the reports of [22] but contrary to [23] who observed an improvement in the feed conversion ratio of birds fed diets with low CP supplemented with essential amino acids in hot summer months. This agrees with [24] and [25] who reported that lower protein content supplemented with amino acid will improve efficiency of protein utilization because equal gain is achieved from a lower protein intake and this amount to reduced cost of production. According to [26] it is reasonable to expect that maximum protein deposition and growth rate is associated with an optimum requirement of crude protein and amino acid, whereas amino acid levels above those needed for growth does not improve deposition likely due to increased rate of protein catabolism.

The breast weight in the methionine possibly (Some missing words here! The author should check his/her manuscript) implying muscle accretion reached a plateau in response to methionine at each level of CP. This was contrary to the report of [14] that increased methionine but not lysine significantly improved breast weight and feed conversion ratio at six weeks of age and not at three weeks. This result also ran contrary to [9] who opined that lysine above NRC levels did not improve breast meat yield unless methionine was also greater than the recommended level. This result further revealed that there are significant interactions between methionine and lysine in broiler diets for carcass parameters with respect to drum stick and breast. This is at variance with the conclusion of [14] that there were no significant interactions between methionine and lysine levels in broiler diets for any parameter studied whether fed amount recommended or in excess.

4. Conclusion

The aim of the poultry farmer is to achieve optimum production with least inputs. From these studies it can be concluded that supplemental methionine and lysine increased monetary losses due to low feed intake, reduced body weight gain and suboptimal feed conversions of the experimental broilers.

COMPETING INTERESTS

The Authors declare no conflict of interest and the studies were not funded by any agency.

AUTHORS' CONTRIBUTIONS

Author A' designed the study and contributed to literature search, Authors C and D managed the analyses of the studies and performed the statistical analysis, while Author B wrote the protocol and the first draft of the manuscript. All authors read and approved the final manuscript."

ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

REFERENCES

1. Smith J. Poultry production in the Tropics. Third edition pp 2000.
2. Ishibashi I. and Yonemochi U. Effects of reducing crude protein in broilers diet. Ani Sci. J 2002;73: 155-165
3. Ogbonna U. Effect of replacing groundnut cake with raw soybean residue diet on performance and nutrient digestibility of cockerel chicks. A preliminary study Nig Poultry Sci J 2000;1:24
4. Schuttle J.B. and Pack M. Effect of sulphur containing amino acids on performance and breast meat deposition of broiler chicks during the growing and finishing phases. Br Poultry sci.1995; 36: 747 -762
5. Bryden W.L., Hew L.I. and Ravindran V. Digestible amino acid values: variation and application .Proceedings of the Australian Poultry Science Symposium 2000;12:51-58
6. Moehn S.,Bertolo R.F., Pencharz P.B., and Ball R.O. Indicator amino acid oxidation responds rapidly to changes in Lysine or protein intake in growing and adult pigs. J Nutr.2004;134: 836-841

7. Carbel M.C. T.L.Godwin and P.W. Waldroup Relationship of dietary lysine level to the concentration of all essential amino acids in broiler diets. Poultry science 1987;80:1472-1479
8. A.O.A.C. Official methods of analysis Association of Official Analytical Chemists Washington D.C.2000.
9. Hickling D.M. Genter and M.E Jackson The Effect of dietary methionine and lysine on broiler chicken performance and breast meat yield. Can. Journal Anim.Sci.1190;70: 67 -68
10. Morgan E T. and S.F.Bilgili Processing losses carcass quality and meat yield of broiler chickens receiving diets marginally deficient to adequate in lysine prior to marketing .Poultry sci.1990;69: 702 -710
11. Si J., C.A.Fritts, D.J.Burnham and P.W.Waldroup Extent to which crude proteins may be reduced in corn-soybean meal broiler diets through amino acid supplementation. Int. J. Poult.Sci.,2004;3:46-50.
12. National Research Council. Nutrient requirement of swine. 9th reviewed National Acad. Press, Washington D.C.1994.
13. Sterling K.G.,G.M.Pesti and R.I Bakalli Performance of broilers fed various levels of dietary lysine and crude protein Poult.Sci 2003;.82:1939-1947.
14. Al-saffar A.A. and Rose S.P. The response of laying hens to dietary amino acids. World poultry journal 2002;58:209-234.
15. D'Mello J.P.F. . Amino acid supplementation of cereal- based diets for non ruminants.Animal Feed Science Technol 1993;45 1-18
16. Aleator V.A. Hamid I.I. Nieb E and Pfeffer E Low protein amino acid supplemented diets in broiler chickens: effects on performance, carcass characteristics, whole body composition and efficiency of nutrient utilization. Journal of Science, Food and Agriculture 2000;80:547-554.
17. Sklan D and I.Plavnik Interactions between dietary crude protein and essential amino acids intake on performance in broilers.Br. Poult.Sci 2002;.30:442-449
18. Jianlin S.I Fritts C.A Burnham D.J. and Waldroup P.W Extent to which crude protein may be reduced in corn-soybean meal broiler diets through amino acid supplementation. International journal of poultry science. 2004;3 (1):46-50
19. Razaei, M Nassiri Moghaddam,H PourReza J. and Kermanshahi H The effects of dietary protein and lysine levels on broiler performance, carcass characteristics and N excretion.Inter.J of poult.sci. 2004;3(2):148-152.

20. Birkett S, and K. de Lange Limitation of conventional models and a conceptual framework for a nutrient flow representation of energy utilization by animals.Br. J. Nutr.2001;86 :647-659.
21. Reeds P.J. Regulation of protein turnover Pages 183-210 in Animal Growth Regulation1989
22. Bregendahl K.J,L. Sell and D.R Zimmerman Effects of low protein diets on growth performance and body composition of broiler chicken. Poultry science 2002; 81: 1156-1167
23. Zarate A.J. ,E.T. Jr.Moran and D.J. Burnham Reducing crude protein and increasing limiting essential amino acid levels in summer reared, slow and fast feathering broilers. J.Appl.Poult.Res.,2003;12:160-168.
24. Jensen L.S. (1989) Relationship between protein and amino acid requirements of poultry.Proceeding of the Georgial Nutrition Conference, 1989; 8-15.
25. Schuttle J.B.Utilization of synthetic amino acids in poultry.6th European symposium on poultry Nutrition 1987;11-12 (Koning Slutte,Germany,World's poultry Science association).
- 26.** Urdaneta-Rincon M. and Leeson S Muscle (pectoralis major) protein turnover in young broiler chickens fed graded levels of lysine and crude protein.Poultry Science2004; 83:1897-1903.

PART 1:

Journal Name:	American Journal of Experimental Agriculture
Manuscript Number:	2013_AJEA_6265
Title of the Manuscript:	EFFECTS OF LOW PROTEIN DIETS SUPPLEMENTED WITH HIGH AMINO ACIDS (METHIONINE AND LYSINE) ON PERFORMANCE OF BROILERS.

280

PART 2:

FINAL EVALUATOR'S comments on revised paper (if any)	Authors' response to final evaluator's comments
1. The authors should react to the questions and comment in lines 44, 82 and 155 respectively. 2.The authors should submit this version of the revised paper for publication as I have gone through it once again to correct possible punctuation and grammatical mistakes	

281

PART 3: Objective Evaluation:

Guideline	MARKS for this REVISED manuscript
Give OVERALL MARKS you want to give to this REVISED manuscript (Highest: 10 Lowest: 0) <u>Guideline:</u> Accept (8-10) Revision required: (4-8) Rejected: (0-4)	8

282