

Species Complex and Seasonal Abundance of Panicle Pests of Sorghum, *Sorghum bicolor* (L.) (Moench) in North-Eastern Nigeria

¹D.M. Dunuwel and ^{2*}A.M. Malgwi

^{1,2} Department of Crop Protection, School of Agriculture and Agricultural Technology, Modibbo Adama University of Technology, P.M.B. 2076, Yola, Nigeria

Authors' contributions

This research was carried out by both authors DMD and AMM. The experiment was designed, supervised, proof read, sent for publication and paid by AMM. The experiment was carried out with data collection and the first draft written by DMD. Both authors contributed for the literature. The final write up was done by AMM.

ABSTRACT

Aims: The species complex and abundance of panicle pests of sorghum was studied on eight sorghum varieties and cultivars in field trial in order identify the pest spectrum complexity on sorghum and at what time of the season they are in abundance and identify which of the sorghum varieties supports fewer pests and incorporate it in integrated pest management programme (IPM) for future studies.

Study design: The experiment was laid in a randomized complete block design (RCBD) which was replicated 3 times on 12m² plots consisting of eight sorghum varieties and cultivars as the.

Place and Duration of Study: The field trial was conducted in two locations at Yola, Adamawa State and Kaltungo, Gombe State in 2012 cropping season in North-eastern Nigeria.

Methodology: Data was collected as from Complete Anthesis Stage (CAS) in September up to Hard Dough Stage (HDS) of sorghum development in November ending. Ten panicles /plot were sampled using transparent polythene bags.

Results: During this trial, 21 pests were identified, but only 18 were common at both Yola and Kaltungo. At both locations three of the pests found were different. Insects identified in order of their abundance were: *Eurystylus oldi*, *Sitophilus zeamais*, *Monolepta* sp., *Orius* sp., *Nola sorghiella*, *Agnoscelis versicola*, *Mirperus jaculus*, *Spilostethus* sp., *Nezara viridula*, *Campylomma* sp., *Achaeearanea tepidoriorum*, *Forficula senegalensis*, *Apis mellifera*, *Messor* sp., *Erythroneura variabilis*, *Poophilus costalis*, *Apocrita polistes*, *Kraussaria angulifera*, *Pachnoda chordata*, *Mantis religiosa* and *Silidius apicalis*. The variety that haboured the highest population of pests was SAMSORG-17 at Yola with a mean population of (609.68/10 panicles) and the least was on Tiksha-Mamza (Ex-Garkida) with a population of (153.33/10 panicles). At Kaltungo, it was SAMSORG-14 haboured the highest population of panicle insect pests (834.34/10 panicles) and the least was on Ex-Tula Farafara cultivar (182.34/10 panicles). The cultivars (Tiksha-mamza, Ex-Tula red and farafara) haboured fewer pests and are more promising than the Samsorgs (14, 17, 37, 39 and 41) improved varieties that haboured more pests. Peak population was in mid-october at soft dough stage.

Conclusion: The local cultivars were more promising than the Samsorg varieties which could be improved for further studies to plant breeders and planting dates could be manipulated to escape period of high pests abundance and hence incorporated in IPM programme as a recommendation for further studies.

Keywords: pests, species, sorghum, varieties, abundance, panicle, Nigeria

1. INTRODUCTION

Sorghum (*Sorghum bicolor* (L.) Moench) belongs to the family Poaceae and originated in Africa which is the fifth world most important crop after wheat, rice and barley [1, 2 and 3]. Millions of people living in semi-arid tropic especially in Africa, Asia and India depend on sorghum as a staple food [4]. Sorghum is faced with many important problems that lead to large yield losses especially caused by parasitic weeds (especially *Striga hermonthica*) anthracnose, downy mildew, grain moulds and sorghum midge and stem borers [5]. The crop is also attacked by a wide and complex range of insect pests worldwide and the total pests load in any area can be very heavy. In spite of the importance of the crop there are few reports on different aspects of sorghum insect pests; progress has been low because of the perceived low market value of the sorghum in developing countries [6, 2 and 3]. Insect pests are a major constraint to increased sorghum production in West Africa, with over 100 species recorded, of which more than 40 are panicle - feeding pests [8, 9, 10, 11 and 12]. In Nigeria, insect pests attacked sorghum at virtually all phonological Stages.

However, only a few of these are considered to be economically important pests in West Africa, among which are stem borers, Sorghum midge, shoofly, spittle bugs and a complex of pentatomid and mirid head bugs [13,14,15, 2,16 and 12]. Panicle infesting insects, feed directly on the reproductive parts of cereal, they often cause direct and irreversible damage [16]. However, [12] further stated that, in Africa, there are few isolate reports which have been documented more specifically on head bugs, but no detailed reports on the species complex and their abundance. However it is believed that improved varieties hold a key to increased and sustainable sorghum production because they combine early maturity with higher yield and have better potential for commercial exploitation and industrial use than the local cultivars. Therefore the need to develop effective management practices for sorghum panicle feeders cannot be over emphasized [11].

This paper, reports on species complex and abundance of panicle pests on different sorghum varieties and cultivars, with the aim of suggesting some peak period of abundance and when to escape high damage which could be incorporated in integrated pest management programme.

2.0 MATERIALS AND METHODS

2.1 Site Description

Field trials were established at the teaching and research farm of the Department of Crop Protection, Modibbo Adama University of Technology, Yola and school farm of Government Girls' Secondary School Kaltungo, Gombe state. Yola is located within latitude $9^{\circ}11'N$ to $9^{\circ}19'N$ and longitude $12^{\circ}31'E$ [17] and Kaltungo is located within latitude $9^{\circ}48'N$ to $9^{\circ}51'N$ and longitude $11^{\circ}18'E$ to $11^{\circ}32'E$ [18] both in Guinea savanna zone of Nigeria during 2011 cropping season.

2.2 Sorghum Varieties and Cultivars

Eight varieties and cultivars of sorghum were obtained from three sources. Five of the varieties (SAMSORG-41, SAMSORG-14, SAMSORG-17, SAMSORG-37 and SAMSORG-39) were obtained from the Institute for Agricultural Research (IAR) /Ahmadu Bello University (ABU) Samaru, Zaria, one cultivar Tiksha Mamza (Ex-Garkida) was obtained from Garkida and two cultivars (Ex-Tula Red Cultivar and Ex-Tula Farafara) were obtained from Tula. This varieties and cultivars characteristic have already been described and discussed [19 and 20].

2.3 Experimental Design

The experiment was laid out in a randomized complete block design (RCBD) replicated three times with varieties and cultivars of sorghum mentioned above as the treatments. Each replication was separated by 2m apart and inter - plot spacing was 1.5m while the plot size was $3 \times 4m^2$ ($12m^2$). The intra-plot spacing was 0.4m and inter-plot spacing was 0.75m.

2.4 Cultural practices

The fields were ploughed with a tractor mould-board plough. The fields were later marked out as earlier mentioned. Apron star 50 DS (Metal-axyl 10%, carboxin 6% +furathiocarb 34%) at the rate of 1 sachet per 10 kg of sorghum seeds was used to dress the seeds before planting. The seeds were sown on 24th June at Yola and 28th June at Kaltungo in 2011 for effective data collection because the distance is more about 246km from Yola since the sampling and data collection of pests have to be carried out in the morning hours. The dressed seeds were sown on the prepared plots in holes of depth of 2cm at the rate of 4-6 seeds per hole. The plants were later thinned to 2 plants per stand at two weeks after emergence. Split

application of fertilizer NPK 15:15:15 was applied at four and eight weeks after emergence at the rate of 64 kg Nha⁻¹. Manual hoe weeding was carried out two times depending on the weed infestation and followed by earthening around the sorghum stands in order to prevent lodging of the sorghum plants.

2.5 Sampling of Pests, Predators and Pollinators

Sampling and pests collection was carried out in the morning hours (06:00-08:30hrs) when the pests were at rest or inactive using transparent polythene bags size measuring 54cm (0.5m) length and 36cm (0.36m) width as described by [21, 22 and 12]. The date of the Sampling varied with the varieties and cultivars depending on the date of flowering, maturity and location. This was done on weekly basis from September to November ending (from CAS to HDS of sorghum development) in each location. Due to the distance in site from Yola to Kaltingo which is about 246km, time of data sampling of 48hours was spaced to allow effective data collection since this has to be carried out in the early morning hours before sun set. Ten (10) panicles were sampled randomly per plot at on a weekly basis and the pests were collected by shaking each panicle vigorously into the polythene bags. The insects were immobilized by placing a cotton wool swab with chloroform to prevent flying pests from jumping out while collecting the other panicles using same polythene bag to sample 10 panicles per plot. The contents of the polythene bags were taken to the laboratory and were sorted out, counted and recorded. Identification of pests was done at the Institute for Agricultural Research / Ahmadu Bello Zaria Insect Museum with the help of the insect reference collection boxes/cabinets and the Curator. The larvae found in each of the sampled panicles was later collected in insect rearing jars of capacity 1liter with fragments of sorghum panicle at soft dough stage reared to adult for clarity in identification.

2.6 Statistical Analysis

The data obtained on the eight sorghum varieties and cultivars were subjected to Analysis of Variance (ANOVA) using (23) 12 Software Version 8 (2) and Student-Newman-Keuls (SNK) was used to separate the means at $P = .05$ test for variables.

3.0 RESULTS AND DISCUSSION

The result for the species complex for both locations (Yola and Kaltungo) showed that 21 species of pests were identified. This include: *Monolepta* sp., *Spilostethus* sp. (Stal), *Eurystylus oldi* (Poppius), *Nola sorghiella* (Riley), *Agnocelis versicola* (Thumberg), *Compylomma* sp. (Odhanbo), *Erythroneura variabilis*, *Mirperus jaculatus* (Thumberg), *Messor* sp. (Emery), *Pachnoda chordata* (Olivier), *Sitophilus zeamais* (Motschusky), *Poophilus costalis* (Walker), *Kraussaria angulifera* (Krauss), *Nezara viridula* (Linnaeus) and *Silidius apicalis* are pests; (*Achaeearanea tepidariorum*, *Orius* sp., *Forficula senegalensis* (Serville) and *Mantis religiosa*) are predators and (*Apis mellifera*) and (*Apocrita polistes*) are pollinators (Table 1). *A. polistes* though a pollinator, but is also a parasitoid a pest, because it was found feeding on developing grains and deposit some exudates which leads to growth and development of mould on the sorghum grains.

However, it has been well documented [24] that honey bees, *A. mellifera* and Wasp, *A. polistes* are well-known pollinators and they occurred during the anthesis stage of sorghum development, these are not pests except they sting human beings working on the farm. *Orius* sp. and preying mantis, *M. religiosa*, occurred in abundance at complete anthesis and milk stages of sorghum development, while earwigs, *F. senegalensis* occurred at soft dough stage of sorghum development in low population. Spiders, *A. tepidariorum* were found at all stages of sorghum development which they are not pests, but a predator mostly on headbugs and these findings also agree with [24, 25, 8, 26, 2 and 12].

Table 1: Species Complex Identified in Sorghum Panicle

Species	Order	Family	Stage found	Pest Status
<i>Monolepta</i> sp.	Coleoptera	Chrysomelidae	Adult	Pest
<i>Spilostethus</i> sp.	Heteroptera	Lygaeidae	Adult	Pest
<i>Eurystylus oldi</i>	Hemiptera	Miridae	Adult & nymphs	Pest
<i>Nola sorghiella</i>	Lepidoptera	Nolidae	Larva & flying Adult	Pest
<i>Agnoscelis versicola</i>	Hemiptera	Pentatomidae	Adult & nymph	Pest
<i>Campylomma</i> sp.	Hemiptera	Miridae	Adult & nymph	Pest
<i>Erythroneura variabilis</i>	Hemiptera	Cicadellidae	Adult & nymph	Pest
<i>Mirperus jaculus</i>	Hemiptera	Alydidae	Adult & nymph	Pest
<i>Messor</i> sp.	Hymenoptera	Formicidae	Adult	Pest
<i>Pachnoda cordata</i>	Coleoptera	Scarabaeidae	Adult	Pest
<i>Sitophilus zeamais</i>	Coleoptera	Curculionidae	Larvae & Adult	Pest
<i>Poophilus costalis</i>	Hemiptera	Aphrophoridae	Adult & nymph	Pest
<i>Kraussaria angulifera</i>	Orthoptera	Acrididae	Adult & nymph	Pest
<i>Nezara viridula</i>	Hemiptera	Pentatomidae	Adult	Pest
<i>Silidius apicalis</i>	Coleoptera	Cantharidae	Adult	Pest
<i>Achaearanea tepidariorum</i>	Theridiidae	Araneae	Adult	Predator
<i>Orius</i> sp.	Hemiptera	Anthocoridae	Adult	Predator
<i>Forficula senegalensis</i>	Dermaptera	Forficulidae	Adult	Predator
<i>Mantis religiosa</i>	Dictyoptera	Mantoidae	Adult	Predator
<i>Apis mellifera</i>	Hymenoptera	Apidae	Adult	Pollinator
<i>Apocrita polistes</i>	Hymenoptera	Vespidae	Adult	Pollinator/pest/Parasitoid

162

163

164

165

166

167

168

169

170

171

172

173

The result showed that, weekly population trend of panicle insect pests in the two locations (Yola and Kaltungo) differs slightly. At Yola, it was the 7th week (24 October) that had the highest population of insect pests (60.50), the lowest population of insect pests was recorded in the 12th week (28 November) (0.75) while at Kaltungo the highest population was recorded in the 6th week (17 October) (100.38) and the lowest population in the 11th week (21 November) (1.46). When the two locations were combined, it was the 6th week (17 October) with the highest population of about 74.65 and the lowest in the 12th week (28 November) with 1.45 (Table 2). The finding also corroborates the earlier results of [2] and [8, 12], that panicle insect pests usually peak at mid-October which coincides with the SDS of sorghum development. This signifies the point at which control measures should be initiated before the pest situation reaches economic damage.

The chrysomelid beetle, *Monolepta* sp. occurred in abundance at Yola and mostly on Ex-Tula farafara cultivar, sap sucking bug, *Spilostethus* sp.; headbug, *E. oldi* the insect pest that has the highest population count and SAMSORG-14 harboured these pests in high populations both at Yola and Kaltungo. This corroborates the findings of [2] and [1] that among the panicle insect pests found in West Africa, most especially Nigeria, *E. oldi* remained the most abundant and important in reducing sorghum grain quality. The webworm, *N. sorghiella* were found on sorghum panicles which formed webs on developing grains that further contaminates the quality of the grains. The stink bug, *A. versicola*; head bug, *Campylomma* sp.; hoppers, *E. variabilis*; pod sucking bug *M. jaculus*; harvester ant, *Messor* sp.; pod sucking bug, *P. cordata* and maize weevil, *S. zeamais* were common in both locations and were common on all the varieties at SDS. *Campylomma* sp. and *M. jaculus* are sap suckers, while the rest were found feeding on the grains which are typical of their characteristics as earlier documented by [7, 8, 910 and 11]. The *S. zeamais*, spittle bug, *P. costalis* and grasshopper, *K. angulifera* occurred in abundance only in Kaltungo. The green stink bug, *N. viridula* and *S. apicalis* were only found in Yola. *N. viridula* occurred only on SAMSORG-17. The differences in specie and abundance of pests in Yola and Kaltungo could be due to the differences in geographical zones regarding rainfall and probably temperature. Such differences also validate the findings of [27, 2 and 28], who found that geographical locations could affect pests population negatively or positively depending on the habitat, rainfall patterns, temperature and relative humidity at a point in time when data was collected.

The mean cumulative pests on the eight sorghum varieties and cultivars in 2011 cropping season in the two locations (Yola and Kaltungo) are presented in Table 3. At Yola, the variety that harboured the highest population was SAMSORG-17 with 50.81 / 10 panicles, followed by Ex-Tula Farafara cultivar (46.19 / 10 panicles), SAMSORG-14 (44.56 / 10 panicles) and least was on Tiksha-Mamza (Ex-Garkida) (12.78 / 10 panicles). At Kaltungo, it was SAMSORG-14 had the highest population of 69.53 / 10 panicles followed by SAMSORG-37 (52.19 / 10 panicles), and the least was on Ex-Tula Farafara cultivar (15.19 / 10 panicles). In the combined analysis, SAMSORG-14 was leading in the population of insect pests with (57.04 / 10 panicles) followed by SAMSORG-17 (44.10 / 10 panicles) and the least occurred in Tiksha-Mamza (Ex-Garkida) with a population of insect pests 19.11 / 10 panicles (Table 3).

Table 2: Mean weekly population of panicle pests of sorghum 2011 cropping season in the two locations.

Time of the Year	Yola	Kaltungo	Combined analysis
12 and 14 Sept.	10.33 ^{de}	3.54 ^d	6.94 ^f
19 and 21 Sept.	22.54 ^{cd}	18.38 ^c	20.46 ^e
26 and 28 Sept.	31.38 ^c	26.67 ^c	29.02 ^{de}
3 and 5 Oct.	20.83 ^{cd}	57.25 ^b	39.04 ^c
10 and 12 Oct.	45.83 ^{ab}	52.04 ^b	48.94 ^b
17 and 19 Oct.	47.75 ^{ab}	100.38 ^a	74.06 ^a
24 and 26 Oct.	60.50 ^a	30.17 ^c	45.33 ^{bc}
31 Oct. and 2 Nov.	50.83 ^{ab}	96.46 ^a	73.65 ^a
7 and 9 Nov.	49.29 ^{ab}	24.54 ^c	36.92 ^{cd}
14 and 16 Nov.	35.13 ^{bc}	4.29 ^d	19.71 ^e
21 and 23 Nov.	1.88 ^e	1.46 ^d	1.67 ^f
28 and 30 Nov.	0.75 ^e	1.54 ^d	1.45 ^f
Mean	31.42	34.73	33.07
CV (%)	66.99	33.38	59.82
SE	21.05	18.54	19.78

Means with the same letter(s) are not significantly different at $P=.05$ using Student-Newman-Keuls (SNK) test for variables

Table 3: Mean cumulative pests on sorghum varieties in 2011 cropping season in two locations

Varieties	Yola	Kaltungo	Combined analysis
SAMSORG-41	21.94 ^{bc}	25.58 ^{de}	23.76 ^{cd}
SAMSORG-14	44.56 ^a	69.53 ^a	57.04 ^a
SAMSORG-17	50.81 ^a	37.39 ^c	44.10 ^b
SAMSORG-37	27.97 ^b	52.19 ^b	40.08 ^b
SAMSORG-39	18.75 ^{bc}	32.53 ^{cd}	25.64 ^{cd}
Tiksha-Mamza (Ex-Garkida	12.78 ^c	25.44 ^{de}	19.11 ^d
Ex-Tula Red Cultivar	28.36 ^b	19.94 ^e	24.15 ^{cd}
Ex-Tula Farafara Cultivar	46.19 ^a	15.19 ^e	30.69 ^c
Mean	31.42	34.73	33.07
CV(%)	66.99	53.38	59.82
SE	21.05	18.54	19.78


Means with the same letter are not significantly different at $P=.05$ using Student-Newman-Kuels (SNK) test for variables

Species complex and abundance of insect pests at Yola and Kaltungo in 2011 cropping season are presented in Table 4. At Yola, *Monolepta* sp. was the highest insect species that occurred (2.60) followed by *E. oldi* (2.38) and the least was *M. religiosa*. At Kaltungo, *Eurystylus oldi* was the highest insect species followed by *Monolepta* sp. (1.28) with the least in *Pachnoda chordata* (0.72) (Table 4). The species complex and abundance of panicle pests on eight sorghum varieties per week in location 1 (Yola) and location 2 (Kaltungo) in 2011 cropping season and the combined analysis are presented in figures 1, 2 and 3 respectively. In figure 2, the peak population of *E. oldi* occurred on 12 October with a mean population of 4.24. Population of *Monolepta* sp. was high on 19 October with a mean population of 4.24. The lowest population of species occurred at the last stage of sorghum development, from 23 to 30 November and it was the hard dough stage of the sorghum development. In figure 3, the peak population of *E. oldi* occurred on 31 October with a mean population of 4.93 and *S. zeamais* occurred on 17 October with a mean population of 6.57. These pests were found to be higher in 17 October and 31 October (milk stage and soft dough stage). 21 November and 28 November supported the lowest insect population in the sorghum (hard dough stage). In figure 3, which is the combined analysis the pests that was higher in all the weeks was *E. oldi*, *Orius* sp. and *Monolepta* sp. and where *Orius* sp. was high in the last week of September and first week of October at milk stage of sorghum the head bug population decreased especially *E. oldi*, and this validates [2]. The lowest insect population of were *M. religiosa* and *P. cordata*.

Table 4: Species complex and abundance of pests on sorghum varieties mean of 10 panicles per plot in the two locations in 2011 cropping season

Species	Yola	Kaltungo	Mean	CV (%)	SE
<i>Monolepta</i> sp.	2.60 ^a	1.28 ^{ab}	1.94	39.44	0.77
<i>Apis mellifera</i>	1.33 ^{ab}	0.86 ^b	1.09	47.07	0.51
<i>Achaearanea tepidariorum</i>	1.06 ^{ab}	1.07 ^b	1.07	26.30	0.28
<i>Spilostethus</i> sp.	0.86 ^b	1.22 ^{ab}	1.04	36.06	0.78
<i>Orius</i> sp.	0.74 ^b	0.85 ^b	0.80	22.78	0.18
<i>Forficula senegalensis</i>	0.85 ^b	0.74 ^b	0.80	22.78	0.18
<i>Eurystylus oldi</i>	2.38 ^a	2.17 ^a	2.28	42.31	0.81
<i>Nola sorghiella</i>	0.84 ^b	0.80 ^b	0.82	22.67	0.19
<i>Oebalus gugnax</i>	0.78 ^b	0.95 ^b	0.86	26.30	0.23
<i>Campylomma</i> sp.	0.78 ^b	0.75 ^b	0.77	28.46	0.22
<i>Erythroneura variabilis</i>	0.75 ^b	0.73 ^b	0.74	15.69	0.12
<i>Mirperus jaculatus</i>	0.74 ^b	0.79 ^b	0.76	17.98	0.14
<i>Messor</i> sp.	0.73 ^b	0.83 ^b	0.78	16.62	0.13
<i>Mantids religiosa</i>	0.71 ^b	0.74 ^b	0.72	13.17	0.10
<i>Pachnoda chordate</i>	0.73 ^b	0.72 ^b	0.72	11.45	0.08

Means with the same letter are not significantly different at $P=0.05$ using Student-Newman-Kuels (SNK) test for variables




I = Standard Error Bar
m in 2011

254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272

Key:

- | | | |
|--|---|-----------------------------------|
| Ins 1 = <i>Monolepta</i> sp. | Ins 8 = <i>Nola sorghiella</i> | Ins 15 = <i>Pachnoda cordata</i> |
| Ins 2 = <i>Apis mellifera</i> | Ins 9 = <i>Agnoscelis versicola</i> | Ins 16 = <i>Silidius apicalis</i> |
| Ins 3 = <i>Achaeearanea tepidariorum</i> | Ins 10 = <i>Campylomma</i> sp. | Ins 17 = <i>Apocrita polistes</i> |
| Ins 4 = <i>Spilostethus</i> sp. | Ins 11 = <i>Erythroneura variabilis</i> | Ins 18 = <i>Nezara viridula</i> |
| Ins 5 = <i>Orius</i> sp. | Ins 12 = <i>Mirperus jaculus</i> | |
| Ins 6 = <i>Forficula senegalensis</i> | Ins 13 = <i>Messor</i> sp. | |
| Ins 7 = <i>Eurystylus oldi</i> | Ins 14 = <i>Mantis religiosa</i> | |

273
274



275
276
277
278
279
280
281
282
283

Key:

- | | | |
|--|---|---------------------------------------|
| Ins 1 = <i>Monolepta</i> sp. | Ins 8 = <i>Nola sorghiella</i> | Ins 15 = <i>Pachnoda cordata</i> |
| Ins 2 = <i>Apis mellifera</i> | Ins 9 = <i>Agnoscelis versicola</i> | Ins 16 = <i>Sitophilus zeamais</i> |
| Ins 3 = <i>Achaeearanea tepidariorum</i> | Ins 10 = <i>Campylomma</i> sp. | Ins 17 = <i>Poophilus costalis</i> |
| Ins 4 = <i>Spilostethus</i> sp. | Ins 11 = <i>Erythroneura variabilis</i> | Ins 18 = <i>Kraussaria angulifera</i> |
| Ins 5 = <i>Orius</i> sp. | Ins 12 = <i>Mirperus jaculus</i> | |
| Ins 6 = <i>Forficula senegalensis</i> | Ins 13 = <i>Messor</i> sp. | |
| Ins 7 = <i>Eurystylus oldi</i> | Ins 14 = <i>Mantis religiosa</i> | |




Figure 3: Species complex and abundance of insect pests per week on sorghum in 2011 cropping season in combined analysis

Key:

Ins 1 = *Monolepta* sp.
 Ins 2 = *Apis mellifera*
 Ins 3 = *Achaearea tepidariorum*
 Ins 4 = *Spilostethus* sp.
 Ins 5 = *Orius* sp.
 Ins 6 = *Forficula senegalensis*
 Ins 7 = *Eurystylus oldi*

Ins 8 = *Nola sorghiella*
 Ins 9 = *Agnoscelis versicola*
 Ins 10 = *Campylomma* sp.
 Ins 11 = *Erythroneura variabilis*
 Ins 12 = *Mirperus jaculus*
 Ins 13 = *Messor* sp.
 Ins 14 = *Mantis religiosa*

Ins 15 = *Pachnoda cordata*

The results showed that, all the SAMSORG varieties, harbors more panicle insects, compared to the local cultivars of Ex-Garkida and Ex-Tula. These showed that, they are promising cultivars which could be resistant to panicle insect pests. Tiksha-Manza despite that it a semi-compact panicle, the red pigmentation which is high in tannin content might have been the reason that it was not favourable to panicle insect pests of sorghum, particularly *E. oldi*, which supported the earlier findings of [2 and 29] that red pigmented sorghum that are usually high in tannin content is unfavourable to panicle insect pests. The Ex-Tula(s) are tall, with loose-panicles which usually mature late and were not favourable for nesting of insects due to its pendulous panicles which matures late in the season. The height of Ex-Tula which matures late in the season usually coincides with the onset of harmattern period which comes with the strong North-east wind that is dry and cold, makes it almost impossible for the insects/pests to nest or hibernate freely in the panicles that are loose which could have been the reason for its low numbers or population. This finding also corroborates [2].

4.0 CONCLUSION AND RECOMMENDATION

The results on the species complex and abundance of sorghum panicle pests is an important step in knowing the complexity and abundance of a whole pest spectrum on sorghum at different time in the season or year they appear mostly and at what growth stage of sorghum they appear; and which of the sorghum varieties harbors the highest panicle pests population. It is also a step that reveals planting dates could be manipulated to escape high incidence and damage due to insect pests of sorghum as earlier carried by [30] on *E. oldi*. For further studies, it is recommended that, detailed studies on each of the pests be investigated on different varieties and planted at different times to find out whether planting dates could be harmonized and manipulated to escape high populations thereby escaping pest damage. All the local cultivars [Tiksha mamza (Ex-Garkida) and the Ex-Tula red and farafara] which supports fewer number of pests should be of interest to plant breeders which could be developed and released to farmers than the SAMSORG varieties support more insects even though they mature early in the season.

ACKNOWLEDGEMENTS

We are grateful to Dr. Valdiya Tamme who analysed the data statistically and made some technical adjustments in the figures. Our thanks go to Mr. Ahmed Santuraki who arranged the references and introduced and encouraged us to publish in ScienceDomain journals.

COMUPUTING INTERESTS

Authors have declared no computing interest and the research was jointly carried out.

REFERENCES

1. ICRISAT (International Crop Research Institute for Semi-Arid Tropics). Sahelian Centre (ICRISAT) West Africa Programme Annual Report 1990. Bp 12404 Niamey, ICRISAT Sahelian Centre; 1991.
2. Malgwi, A.M. Studies of Sorghum, (*Sorghum bicolor* L. Moench) head bugs. Samaru, Zaria. MSc. (Crop Protection) Thesis, Ahmadu Bello University Zaria; 1997.

3. Bello, D. Combining Ability and Heteroseres for Agronomic characters and malting Quality Traits of Grain Sorghum in Adamawa State a Ph.D Thesis submitted to Department of Crop production and Horticulture. Federal University of Technology Yola Pp 1 – 4; 2010.
4. Singh, S.P., Jotwani, M.G., Rano, B.S. and Rao, N.G.P. Stability of host plant resistance to sorghum shootfly, *Atherigora soccata* (Rondani). India Journal of Entomology. 2003; 40: 376-383.
5. Taylor, J.R.N. Overview: Importance of Sorghum in Africa. In: Belton, P.S. & Taylor, J.R.N. (Editors). Proceedings of the workshop on the proteins of sorghum and millet: enhancing nutritional and functional properties for Africa, Pretoria, South Africa, 2003; 2-4 April. Afripro (internet) Accessed 2011/01/18. <http://www.afripro.org.uk/>.
6. Leuschner, K. Sorghum Entomology Research Programme and Need in Developing World; 1985.
7. Nwanze, K.F. Sorghum Pests of West Africa in: proceeding of International Sorghum Entomology Workshop 15-21 July, 1985. College Station Texas, USA Pantecheru, pp 502324 India ICRISAT 1985; pp. 37-43.
8. McFarlane, J.H. The hemipterous insects and spiders of sorghum in Northern Nigeria. Insect Science and its Application. 1989; 10: 277-284.
9. Seshu Reddy, K.V. Insect Pests of Sorghum in Africa. Insect Science and its applications. 1992; 12 (516): 653-657.
10. Malgwi, A.M. The damage potentials of the most prominent headbug, *Eurystylus oldi* Poppius as a head bug in Nigeria. Nigerian Journal of tropical Agriculture. 1999; Vol. 1. : 21-25.
11. Tanzubil, P.B., Alen, A. and Zakaria, M. Insect infestant les panicules de sorgho (*Sorghum bicolor* (L.) Moench) dans le nord du Ghana 1.La. Ghana. Journal of Agricultural Science, 2007; 40: 43-5044.
12. Malgwi, A.M. and Ajayi, O. Population dynamics of *Eurystylus oldi* Poppius (Heteroptera: Miridae) at different stages of sorghum development at Samaru, Nigeria. Journal of Arid Agriculture. 2011; 97-103.
13. Bowden, J. Sorghum midge, *Contarinia sorghicola* and other causes of grain sorghum losses in Ghana. Bull. Ent. Res 1965; 56, 169 – 189.
14. Agyen-Sampong, M. Insect pests of sorghum head and assessment of crop loss by the major pests. Ghana Journal of Agric. Sci. 1978; 11: 109 – 115.
15. Tanzubil, P.B. Non-pesticidal approaches to managing insect pests of sorghum in Ghana. In Improvement of cropping systems in the savanna zone: The challenges ahead. Proceedings of the 3rd National workshop on improving farming systems in the interior Savanna of Ghana (ed. H. Mercer-Quarshie, K. O. Marfo, S.A. Langyintou & R. Owusu), pp 227 – 236, April 1996. Nyankpala; 1997.
16. Tanzubil, P. B. Zakaria, M. and Alem, A. Population ecology and damage potential of Mirid bugs infesting sorghum panicles in Northern Ghana. Tropical Science. 2005; 45, 58 – 62.
17. UBRBDA (Upper Benue River Basin Development Authority). Annual Report Upper Benue River Basin development Authority Yola – Nigeria. 1998; 1999: 31-32.
18. Wikipedia. Kaltungo. 2011; En. Accessed 2012/03/23. wikipedia.org/wiki/kaltungo
19. Dunuwel, D. M. Species Complex, Abundance, Damage and Control of Panicle Insect Pests of Sorghum, *Sorghum bicolor* (L.) Moench at Gombe and Yola in North Eastern Nigeria. M.Sc Thesis 90pp (unpublished) submitted to the Department of Crop Protection, Modibbo Adama University of Tech., Yola; 2012.
20. Malgwi, A.M. and Dunuwel, D. M. Damage and Control of Panicle Insect Pests of Sorghum (*Sorghum bicolor* (L.) Moench) in North eastern Nigeria. ARPN Journal of Science and Technology. 2013a; Vol. 3, No. 2: 195-202pp. February 2013. Walnut Street, USA KAGI Academic Publishers 9pp. ISSN 2225-7217 . ©2011-2013.
21. Steck, G.J., Teetes, G.L. and Maiga, S.D. Species composition and injury to sorghum by panicle feeding bugs in Niger. Insect Science and its Application. 1989; 10:199-217.
22. Sharma, H.C., Taneja, S.L., Leuschner, K. and Nwanze, K.F. Techniques to screen sorghum for resistance to insect pest. ICRISAT Information Bulletin, 1992; No. 32:4pp.
23. SAS (Statistical Analysis Systems). SAS/STAT User's Guide, Version 8 (2). Statistical Analysis Systems. Institute Inc., Cary, North Carolina, USA; 2000.
24. Dennis, S.H. Agricultural insect pests of the tropics and their control. 2nd Edn Cambridge, U.K., Cambridge University press pp 174;1983.
25. Descamps, M. Insectes nuisible aux cultures et. Insectes pre'dateurs re'cement observe's dans le Nord Cameroon. Agronomie Tropicale. 1954; 9: 174-182.
26. Ratnadas, A. and Ajayi, O. Panicle insect pests of Sorghum in West Africa pages 29-38 in. panicle Insect pests of Sorghum and pearl millet: Proceedings of an international consultative workshop 4-7 Oct. 1993.

*Tel.: +234 (0) 70 3642 0355

Email: annamalgwi@yahoo.co.uk

- ICRISAT Sahelian center Niamey, Niger (Nwanze, K.F and Youm, O. eds) Patancheru 502 324, A.P. India. ICRISAT; 1995.
27. Doumbia Y. O and Teetes, G.L. Distribution, dégâts et dynamique des populations d' *Eurystylus marginatus* (Odh.) [Heteroptera: Miridae], ravageur du sorgho au Mali. Conférence Inter-réseaux sur la recherche et la production des cultures vivrières en Afrique semi-aride. Niamey 7 – 15 Mars 1991, Niger; 1991.
 28. Doumbia, Y.O. Incidence and distribution of sorghum headbug, *Eurystylus oldi* (Hemiptera: Miridae) in West and Central Africa. Insect Science and its Application, 2001; 21: 103-117.
 29. Yadumma, J.J. The effect of Tamarind ash and Permaguard on *Sitophilus zeamais* (Moth.) (Coleoptera:Curculionidae) on stored maize and sorghum. Unpublished undergraduate project; 2000.
 30. Malgwi, A.M., Ajayi, O., Ogunlana, M.O. and Onu, I. Evaluation of headbugs Population and Damage on Sorghum, at different sowing dates at Samaru, Zaria. Entomological Society of Nigeria (ESN) Occasional Publication, 2000; 32: 107 – 112. ISBN 978-33086-2-9.