

Activity of Catambra extracts against *Meloidogyne* spp.

Aims: In this study we tested four Catambra commercial formulates (hydroalcoholic extracts) used as insecticides in Italy, against *Meloidogyne incognita* and *Meloidogyne javanica*.

Catambra plant (Ambrogios, Catalpa) a European patent belonging to Ambrogio srl, rich in an iridoid glycoside catalpol, is used to date as a mosquito deterrent, but no data exist on its biological activity against soil parasites as *Meloidogyne* sp. *Catalpa* species demonstrate various biological properties of pharmaceutical and agronomical interest, mainly due to their contents in glycosides.

Place and Duration of Study: All paralysis bioassays were performed in Pesticide Science Laboratory, of Aristotle University Thessaloniki, Greece, while chemical analysis in Department of Life and Environmental Sciences, University of Cagliari.

Methodology: The activity was tested in paralysis bioassays with freshly hatched second-stage juveniles (J2) immersed in test solutions for four time periods (12, 24, 48 and 96 hours). Catalpol contents of the extracts were measured by means of HPLC/MS and its activity was also tested against J2. The effect of exhaustive evaporation of hydroalcoholic extracts prior use in bioassays was also studied for nematicidal activity.

Results: Out of the four extracts tested without being evaporated first, two extracts (C-2 and C-4) exhibited activity and the $EC_{50/48h}$ on *M. incognita* were calculated at 640 and 728 µg/ml. On the other hand, *M. javanica* was proven sensitive only to the C-4 extract and the activity had an $EC_{50/48h}$ value at 500 µg/ml. When extracts were exhaustively evaporated before use, only the C-4 paralysed J2 and the $EC_{50/48h}$ values were 728 and 500 µg/ml against *M. incognita* and *M. javanica*, respectively. Interestingly, catalpol was not found to be nematicidal at the concentration range of 100 to 1000 µg/ml. The nematicidal activity of the extracts was therefore not correlated with their content in catalpol. This was confirmed by the chemical composition study, since the C-4 extract, exhibiting activity against both species, was of low contents in catalpol (21 µg/ml), and the C-3 extract that had the highest contents in catalpol (358 µg/ml) did not paralyse J2.

Conclusion: Catambra nematicidal activity is not based in its contents in catalpol, and further studies are conducted to delineate for activity among other constituent components.

Keywords: Catambra; catalpol; nematicidal activity; *Meloidogyne* spp.; bionematicides

1. INTRODUCTION

Plant parasitic nematodes are among the most destructive group of plant pathogens worldwide and their control is extremely challenging [1,2]. Phytonematodes parasitize a large variety of crops worldwide and their impact on yield losses has been estimated to 100 billion US\$ annually [3,4]. Specifically, Root-knot nematodes (RKN) or *Meloidogyne* sp. is the most

notorious genus [3]. Strategies to control these pests include chemical nematicides combined with cultural practices, crop rotation, and resistant cultivars. Considering that only few synthetic nematicides are still authorized [5], natural nematicides isolated from plants or microorganisms are gaining an increased interest as alternative less toxic and eco friendly biocontrol agents [1,6]. The new European legislation asks from the members to adopt (by 1 January 2014) national action plans and take all necessary measures to promote low pesticide-input farming, including integrated pest management, and to ensure that professional users of pesticides shift towards a more environmentally-friendly use of all available crop protection measures, giving priority to low-risk, non-chemical alternatives wherever possible [7,8]. This action gives a specific status to nonchemical and natural alternatives rather than conventional chemical pesticides, and requires them to be given priority wherever possible. As nematodes cause severe economic loss to many agricultural crops, a great demand for discovering environmentally acceptable substitutes for commercial nematicides exists, creating a great market opportunity for biorational products such as botanical nematicides. Recently, it has been reported that plant derived nematocidal compounds, such as furfural, (E)-2-decenal, (E,E)-2,4-decadienal, and 2-undecanone are greatly effective in controlling RKN both in vitro and in pot experiments [9-13]. Furthermore, according to the international bibliography, the control of nematodes on different vegetables has been often performed by botanical extracts [14-18]. Natural substances of plant origin that have been found to exhibit nematocidal activity are Plant Secondary Metabolites (PSMs) of various plants and belong to different chemical classes, such as aldehydes and ketones, alkaloids, glycosides, glucosinolates and isothiocyanates, limonoids, quassinoids and saponins, organic acids, phenolics and flavonoids, polyacetylenes and terpenes [1,19].

Catalpa (Bignoniaceae) species exhibit a range of biological activities of interest in pharmacy and agriculture. *Catalpa ovata* stem bark may be a useful external medicine for treatment of atopic dermatitis [20] and enhancement T cell-mediated immune responses with anticancer properties. [21] *Catalpa bignonioides*, known for its contents in naphthoquinone, catalponol, catalponone and catalpalactone [22] exhibits significant antioxidant activity [23]. D. Muñoz-Mingarro et al. [24] reported anti-inflammatory and antinociceptive actions in ethyl ether, butanolic and aqueous fractions of *Catalpa bignonioides* pod extract. In fact due to the interesting biological properties of the phenylethanoid glycosides, advances have been made so as to increase 10 times their production infecting *C. ovata* root cultures with four agropine strains of *Agrobacterium rhizogenes* [25].

Interestingly, soil treating with *C. bignonioides* has been found to act as biological soil disinfestations for 4 years against the nematode *Pratylenchus fallax* [26]. On the other hand, flavone and flavonol glycosides, as constituent compounds of *Vicia pannonica* Crantz var. *purpurascens* have been proven of considerable anthelmintic activity [27]. Nonetheless, to date no data exist on the activity of *Catalpa* extracts against *Meloidogyne* sp.

Catambra plant belonging to the Bignoniaceae's family coming from northern America, has been developed and increased in Giovanni Ambrogio's nursery, and has been protected by a European patent 2006/011 (no. EU27931). This tree is infested by few pests of any consequence because it exhibits significant insecticidal activity. Thanks to the high contents of catalpol, catambra shows a repellent action against mosquitoes and in fact the tree is sold in the market as anti-mosquito. Catalpol is an iridoid glucoside (Figure 1), which is simply monoterpene with an attached glucose molecule and its pharmacological action has not been fully established. Chemical analysis carried out from the Experimental Centre for Floriculture confirms that Catambra contains four times a quantity of catalpol contained in other *Catalpa* species [28].




Figure 1. Structure of catalpol

In this study, four commercial Catambra formulates (hydroalcoholic extracts), used as insecticides in Italy, were tested against *M. javanica* and *M. incognita*, in paralysis bioassays with 24 h second stage juveniles (J2) and we correlated the nematocidal activity with the extracts content in catalpol, determined by HPLC/MS analysis. To study for the activity of volatile and non volatile compounds in the extracts, the paralysis experiment was performed using a) the hydroalcoholic extracts and b) the dry yield of the extracts after exhaustive evaporation of the solvents.

2. MATERIAL AND METHODS

2.1. Hydroalcoholic Catambra Extracts

Four commercial Catambra formulates (hydroalcoholic extracts) were provided from Ambrogio Vivai Srl. Original Catambra is guaranteed by the European Patent and it is recognizable thanks to the anti-rip label on its foliage with a special control code and an indelible brand on its trunk [28]. The four Catambra formulates were hydroalcoholic extracts (coded as C-1 C-2, C-3, C-4), obtained from the Catambra plant and used in paralysis bioassays at concentrations of 62.5 to 1000 µg/ml, in order to evaluate their nematocidal activity..

2.2. Chemical composition analysis

Catalpol contents were measured on a HPLC-MS system, Varian tandem mass spectrometer (Palo Alto, CA) consisting of a ProStar 410 autosampler, two ProStar 210 pumps, and a 1200 L triple quadrupole mass spectrometer equipped with a 1100 series diode array detector (Agilent, Waldbronn, Germany). The column used was a Phenomenex GEMINI 3µm C18 (100 x 4.6 mm). The mobile phase consisted of aqueous 0.1% formic acid (A) and (B) methanol. The solvent gradient started at 20 % of A and 80 % of B, reaching 100 % of B at 10. The flow was 0.4 mL/min and the injection volume was 20 µL. Spectrum scan was from 200 to 600 nm. For the detector's analysis parameters set and for the HPLC data analysis, the software ChromQuest Version 4.0 Build 1120 and HP Chem, Version A.07.01 was used, respectively. LC-MS analysis was performed in electrospray ionization interface, in the positive ion mode, as follows: shield potential, 775 V; capillary potential, 55 V; needle voltage, 5700 V; nitrogen pressure, 54mTorr; housing temperature, 56 °C; drying gas temperature, 380 °C; scan time 1; detector multiplier voltage, 1200 V; isolation width of m/z 1.6 for the quadrupole 1; scan 100-1500 amu.

106 2.3. Nematode motility bioassays

107 2.3.1. Nematode cultures

108 A population of *M. incognita* and *M. javanica* originally obtained from tomato roots collected
109 from a greenhouse in Vasilika, Thessaloniki, Northern Greece, was reared on tomato,
110 (*Solanum lycopersicum* L.) cv. Belladonna, a cultivar which is very susceptible to root-knot
111 nematodes. Roots of inoculated plants, 40 days after inoculation, were uprooted, washed
112 free of soil and cut into 2 cm pieces. Eggs were extracted with sodium hypochlorite
113 according to the method of Hussey and Barker, 1973, [29]. Second-stage juveniles (J2) were
114 allowed to hatch in modified Baermann funnels at 27 °C. All J2 hatching in the first 3 days
115 were discarded and freshly hatched second-stage juveniles (J2) collected during the
116 subsequent 24 hours were used for the bioassays, according to methods described by Ntalli
117 et al., 2009, [30].

118 2.3.2. Effect of Catambra extracts and catalpol on J2 motility.

119 Effects of Catambra extracts on J2 motility were tested at the concentrations 1000, 500, 250,
120 125 and 62.5 µg/ml. Catambra formulates were directly and serially diluted in distilled water
121 containing Tween-20 (0.6% v/v) to produce test solutions, the concentration of which was
122 calculated according to the extract's yield in dry components, after exhaustive evaporation of
123 the solvents under stream of nitrogen. Additionally, weighed amounts of the dried Catambra
124 extracts, obtained after this exhaustive evaporation of the solvents, were diluted in distilled
125 water containing Tween-20 (0.6% v/v) to prepare test solutions of the above concentrations.
126 Final concentrations of Tween-20 in treatment well never exceeded 0.3%. Preliminary trials
127 showed that nematodes exposed at these concentrations of Tween-20 were not affected.
128 Plain water as well as a mixture of water, and Tween-20 (0.3% v/v) (carrier control) served
129 as controls. Nematicidal activity of commercially available catalpol was used at a
130 concentration range of 100, 250, 500, 750 and 1000 µg/ml. Catalpol stock solution was
131 prepared in methanol, whereas distilled water containing Tween-20 (0.6% v/v) was used for
132 further dilutions. Final concentrations of methanol and Tween-20 in each well never
133 exceeded 1 and 0.3 % v/v, respectively. Distilled water as well as a mixture of water,
134 methanol and Tween-20 (1 and 0.3 % v/v) (carrier control) served as controls. In all cases,
135 working solutions were prepared containing double the test concentration and were then
136 mixed in Cellstar® 96-well cell culture plates (Greiner bio-one) at a ratio of 1:1 v/v with
137 suspensions of 15 approximately J2 added to each well. Multiwell plates were covered to
138 prevent evaporation and were maintained in the dark at 27°C. Juveniles were observed with
139 the aid of an inverted microscope (Euromex, Holland) at 40× after 12, 24, 48, 96 h and were
140 ranked into two distinct categories: motile or paralysed. Each treatment was replicated 5
141 times as a split plot design, and each bioassay was performed twice.

142 2.4. Statistical analysis

143 The data of nematicidal activity are presented as percentages of paralyzed J2 in the
144 microwell assays corrected by elimination of the natural death/paralysis in the water control
145 according to the Scheider-Orelli's [31] formula: Corrected % = [(Mortality % in treatment –
146 Mortality % in control)/(100 – Mortality % in control)] × 100, and they were analyzed by
147 ANOVA combined over time. As ANOVA indicated no significant treatment by time
148 interaction, means were averaged over experiments. Corrected percentages of paralyzed J2
149 treated with extracts were subjected to Probit analysis and the effective concentrations
150 (EC₅₀) were calculated after transformation of concentrations in logarithmic scale (Figure 2).
151 The mean value of the five replicates per Catambra extract concentration and five
152 concentrations was used to calculate the EC₅₀ value.

3. RESULTS AND DISCUSSION

In order to evaluate the four Catambra extracts (coded as C-1 C-2, C-3, C-4) for nematocidal activity paralysis, bioassays were performed at concentrations 62.5 to 1000 µg/ml. These concentrations provided a linear concentration–response relationship for all extracts tested, although in most tests paralysis percentage assessed at the highest exposed dose, was less than 50%. Exhaustive evaporation was also tried before using the hydroalcoholic extracts in bioassays to study for effects on activity. According to the paralysis bioassays results, only two out of the four Catambra extracts were proved of nematocidal activity and in specific the extracts C-2 and C-4 (Table 1, Figure 2). The extract C-2 exhibited an EC₅₀ value of 640 µg/ml after 48 h of *M. incognita* J2 immersion in test solutions, but it was not found active against *M. javanica* (data not shown). Additionally, the extract C-4 was found to exhibit

Table 1. Paralysis activity (EC₅₀ values)^a of Catambra extracts against *Meloidogyne incognita* and *Meloidogyne javanica* calculated after immersion of J2s in test solutions of extracts for 12, 24, and 48 hours


extracts								
C-1			C-2		C-3		C-4	
EC ₅₀ (µg/ml)								
	<i>M.javanica</i>	<i>M.incognita</i>	<i>M.javanica</i>	<i>M.incognita</i>	<i>M.javanica</i>	<i>M.incognita</i>	<i>M.javanica</i>	<i>M.incognita</i>
12h	n.c ^b .	n.c.	n.c.	n.c.	n.c.	n.c.	n.c.	n.c.
24h	n.c.	n.c.	n.c.	n.c.	n.c.	n.c.	680	780
48h	n.c.	n.c.	n.c.	640	n.c.	n.c.	500	728

^a In cases where EC₅₀ values have not been calculated because they were outside the test concentration range, they are estimated higher than the upper concentration level (1000 µg/ml).


^b n.c.: not calculated

nematicidal activity against both *Meloidogyne* species and the respective EC₅₀ values were calculated at 728 and 500 µg/ml after 48h against *M. incognita* and *M. javanica*. Finally, the rest of the Catambra extracts tested were not found to be nematocidal at the test concentration range used for the bioassay, since the percentage of paralyzed J2 was in all cases less than 42% at the highest tested dose. When catalpol was tested against both nematode species in a dose response experiment, in the concentration range of 100 to 1000 µg/ml, it was not found active and the percentage of paralyzed J2 did not exceed 37% after J2 immersion for 96 h at the highest dose (1000 µg/ml). This fact indicated that the nematocidal activity of the extracts C-2 and C-4 could not be attributed to their catalpol contents. This result was further sustained by the chemical composition analysis results, according to which the catalpol content in the extract C-4, exhibiting activity against both nematodes species, was rather low determined at 21 µg/ml (Table 2). On the contrary, the extract yielding the highest contents in catalpol amounted to 430 µg/ml was found to be the C-3, not of any nematocidal activity whatsoever (Table 1). Interestingly, exhaustive evaporation prior to the extracts' use in bioassays, led to loss of activity of extract C-2, while only the C-4 extract paralyzed J2 (Figure 2, c). The EC₅₀ value calculated after 48 h of J2 immersion in extract C-4 test solutions was 950 and 874 µg/ml for *M. javanica* and *M. incognita*, respectively (Figure 2, c). This fact indicates that the nematocidal principles of extract C-2 were volatile compounds, but some of the active ingredients in extract C-4 were not, since activity remained after evaporation.


195
196



197
198
199



200
201
202



203
204
205
206
207
208
209
210
211
212
213

Figure 2. Regression curves of *M. incognita* (a) and *M. javanica* (b) paralysis following J2 immersion in solutions of extract C-4, for 24 h and 48 h and J2 immersion for 48 h in solution of extract C-4 after exhaustive evaporation of solvents (c). Each point represents the average % number of paralyzed J2 of five replications per treatment (two experiments replications) after elimination of natural paralysis/death observed in the control.

Table 2. Catalpol contents in Catambra extracts according to LC/MS analysis.

Extract	Catalpol mg/ml
C-1	19
C-2	358
C-3	430
C-4	21

Nevertheless, the Catambra extracts activity against *Meloidogyne* sp. is significant considering the concentration levels of other nematicidal botanical extracts, such as *Melia Azedarach* or essential oils from various plants [10, 30, 32, 33]. For this reason, it still remains to be delineated the activity among Catambra components. There was no clear trend of sensibility between the two nematode species, since at cases *M. incognita* was found more sensitive, but in other cases the contrary occurred.

4. Conclusion

On the basis of the present results, Catambra extracts exhibit promising nematicidal activity against both species *M. incognita* and *M. javanica*. Since Catambra is already used as insecticide its use could be amplified for the control of other agricultural pests, such as nematodes. Nonetheless, the nematicidal activity of the extracts is non correlated to catalpol contents. In our ongoing studies, it was focused on searching among other constituents compounds for activity.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ntalli N.G., Caboni P. Botanical Nematicides: A Review. J. Agric. Food Chem. 2012; 60: 9929–9940.
2. Ntalli N.G., Caboni P. Botanical nematicides in the mediterranean basin, Phytochem Rev. 2012; 1–9.
3. Mai W.F. Plant-Parasitic Nematodes: Their Threat to Agriculture; Sasser, J. N., Carter, C. C., Eds.; North Carolina State University Graphics: Raleigh, NC, 1985; Vol. I, pp 11–17.
4. Oka Y., Koltai H., Bar-Eyal M., Mor M., Sharon E., Chet I. New strategies for the control of plant-parasitic nematodes, Pest Manag Sci. 2000; 56: 983–988.
5. http://ec.europa.eu/sanco_pesticides/public/index.cfm.
6. Ntalli N. G., Menkissoglu-Spiroudi U. Pesticides of Botanical Origin: A Promising Tool in Plant Protection. Ch. 1, In Pesticides - Formulations, Effects, Fate, pp 3-14, Ed. Margarita Stoytcheva, 2011, InTech, Croatia.

- 257 7. European Commission 2009. Commission Directive 2009/128/EC of 21 October 2009
258 establishing a framework for Community action to achieve the sustainable use of
259 pesticides. OJ No L 309, 24.11.2009, p.71.
- 260 8. European Commission 2009. Council Regulation (EC) no. 1107/2009 of 21 October 2009
261 concerning the placing of plant protection products on the market and repealing Council
262 Directives 79/117/EEC and 91/414/EEC. OJ No L 309, 24-11-2009, pp 50.
- 263 9. Caboni P., Aissani N., Cabras T., Falqui A., Marotta R., Liori B., Ntalli N.G., Sarais G.,
264 Sasanelli N., Tocco G. Potent Nematicidal Activity of Phthalaldehyde, Salicylaldehyde,
265 and Cinnamic Aldehyde against *Meloidogyne incognita*. J. Agric. Food Chem. 2013; 61:
266 1794–1803.
- 267 10. Ntalli N. G., Vargiu S.; Menkissoglu-Spiroudi U., Caboni P. Nematicidal carboxylic acids
268 and aldehydes from *Melia azedarach* fruits. J. Agric. Food Chem. 2010; 58:
269 11390–11394.
- 270 11. Cavoski I., Al. Chami Z., Bouzebboudja F., Sasanelli N., Simeone V., Mondelli D., Miano
271 T., Sarais G., Ntalli N. G., Caboni P. *Melia azedarach* controls *Meloidogyne incognita* and
272 triggers plant defense mechanisms on cucumber. Crop Prot. 2012; 35: 85–90.
- 273 12. Caboni P., Ntalli N. G., Aissani N., Cavoski I., Angioni A. Nematicidal activity of (E,E)-2,4-
274 decadienal and (E)-2-decenal from *Ailanthus altissima* against *Meloidogyne javanica*. J.
275 Agric. Food Chem. 2012; 60: 1146–1151.
- 276 13. Ntalli N. G., Manconi F., Leonti M., Maxia, A., Caboni P. Aliphatic ketones from *Ruta*
277 *chalepensis* (Rutaceae) induce paralysis on root knot nematodes. J. Agric. Food Chem.
278 2011; 59: 7098–7103.
- 279 14. Cetintas R., Yarba M. M. Nematicidal effects of five plant essential oils on the southern
280 root-knot nematode, *Meloidogyne incognita* race 2. J. Anim. Vet. Adv. 2010; 9: 222–225.
- 281 15. Javed N., Gowen S. R., Inam-ul-Haq M., Abdullah K., Shahina F. Systemic and
282 persistent effect of neem (*Azadirachta indica*) formulations against root-knot nematodes,
283 *Meloidogyne javanica* and their storage life. Crop Prot. 2007; 26: 911–916.
- 284 16. Ladner D. C., Tchounwou P. B., Lawrence G. W. Evaluation of the effect of ecologic on
285 root knot nematode, *Meloidogyne incognita*, and tomato plant, *Lycopersicon esculenum*.
286 Int. J. Environ. Res. Public Health 2008; 5: 104–110.
- 287 17. Giannakou I. O. Efficacy of a formulated product containing *Quillaja saponaria* plant
288 extract for the control of root-knot nematodes. Eur. J. Plant Pathol. 2011; 130: 587–596.
- 289 18. Maistrello L., Vaccari G., Sasanelli N. Effect of chestnut tannins on the root-knot
290 nematode *Meloidogyne javanica*. Helminthologia 2010; 47: 48–57.
- 291 19. Chitwood D.J. Phytochemical based strategies for nematode control. Annu. Rev.
292 Phytopathol. 2002; 40: 221-249.
- 293 20. Yang G., Choi CH., Lee K., Lee M., Ham I., Choi HY. Effects of *Catalpa ovata* stem bark
294 on atopic dermatitis-like skin lesions in NC/Nga mice. J Ethnopharmacol. 2013; 145(2):
295 416-23.
- 296 21. Oh CH., Kim NS., Yang JH., Lee H., Yang S., Park S., So UK., Bae JB., Eun JS., Jeon
297 H., Lim JP., Kwon J., Kim YS., Shin TY., Kim DK. Effects of isolated compounds from
298 *Catalpa ovata* on the T cell-mediated immune responses and proliferation of leukemic
299 cells. Arch Pharm Res. 2010; 33(4): 545-50.
- 300 22. Park B.M., Hong S.Su., Lee C., Lee M.S., Kang S.J., Shin Y.S., Jung J.-K., Hwang B.Y.
301 Naphthoquinones from *Catalpa ovata* and their inhibitory effects on the production of
302 nitric oxide, Arch Pharm Res. 2010; 33: 381-385.
- 303 23. Dvorská M., Zemlicka M., Muselík J., Karafiátová J., Suchý V. Antioxidant activity of
304 *Catalpa bignonioides*. Fitoterapia. 2007; 78: 437-439.
- 305 24. Muñoz-Mingarro D., Acerob N., Llinaresc F., Pozuelod J.M., Galán de Merab A.,
306 Vicentenb J.A., Moralese L., Alguacile L.F., Péreze C. Biological activity of extracts from
307 *Catalpa bignonioides* Walt. (Bignoniaceae). J Ethnopharmacol. 2003; 87:163-167.

- 308 25. Wysokińska H., Lisowska K., Floryanowicz-Czekalska K. Transformation of *Catalpa*
 309 *ovata* by *Agrobacterium rhizogenes* and phenylethanoid glycosides production in
 310 transformed root cultures. *Z Naturforsch C*. 2001; 56: 375-381.
- 311 26. Goud J.-K.C., Termorshuizen A.J., Blok W.J., Van Bruggen A.H.C. Long-term effect of
 312 biological soil disinfestation on verticillium. *Plant Dis*. 2004; 88(7): 688-694.
- 313 27. Kozan E., Anul SA., Tatli II. In vitro anthelmintic effect of *Vicia pannonica* var.
 314 *purpurascens* on trichostrongylosis in sheep. *Exp Parasitol*. 2013; 134(3): 299-303.
- 315 28. <http://www.catambra.it/caratteristiche.htm>
- 316 29. Hussey R.S., Barker K.R. A comparison of methods of collecting inocula of *Meloidogyne*
 317 spp. including a new technique. *Plant Dis Repr*. 1973; 57: 1025-1028.
- 318 30. Ntalli N. G., Menkissoglu-Spiroudi U., Giannakou I. O. Efficacy evaluation of a neem
 319 (*Azadirachta indica* A. Juss) formulation against root-knot nematodes *Meloidogyne*
 320 *incognita*. *Crop Protection*. 2009; 28: 489-494.
- 321 31. Puntener W. (1981) Manual for field trials. In *Plant Protection*. 2nd edn. Basle,
 322 Switzerland: Ciba-Geigy Limited. 205 pp.
- 323 32. Ntalli N.G., Ferrari F., Giannakou I., Menkissoglu-Spiroudi U. Phytochemistry and
 324 nematicidal activity of the essential oils from 8 Greek lamiaceae aromatic plants and 13
 325 terpene components. *J. Agric. Food Chem*. 2010; 58: 7856-7863.
- 326 33. Ntalli N.G., Ferrari F., Giannakou I., Menkissoglu-Spiroudi U. Synergistic and
 327 antagonistic interactions of terpenes against *Meloidogyne incognita* and the nematicidal
 328 activity of essential oils from seven plants indigenous to Greece. *Pest Manag. Sci*. 2011;
 329 67: 341-351.