

**EVALUATION OF SOME PLANT EXTRACTS FOR THE CONTROL OF
BACTERIAL SOFT ROT OF TUBERS**

ABSTRACT

The use of protective chemicals on ware tubers can render it unfit for consumption as tubers undergo little processes before consumption. Hitherto, there is no effective control measure to manage this disease (*Pectobacterium* spp); therefore, selection of some plant extracts for the management the disease will be the preferred option. The objective of the study therefore was to evaluate some plant materials for the management of tuber soft rot bacteria induced by *pectobacterium* spp. Aqueous extracts of neem (*Azadirachta indica* A. Juss), Eucalyptus leaves (*Eucalyptus citrodorus* (L)), Lemon grass (*Cymbopogon citratus* (Stapf)), Garlic bulb (*Allium Sativum* (Linn)), Ginger (*Zingiber officinale* (Roscoe)) and *Aloe vera* were evaluated for the management of bacterial soft rot of some tubers (cocoyam, Irish potato, sweet cassava, sweet potato and white) induced by *Erwinia* spp. Two methods involving *in vitro* and tuber assays were used. Lemon grass, garlic, Aloe Vera, neem extracts and borax salt had the greatest inhibitory effect on *Pectobacterium* spp., and therefore recommended for the management of tuber soft rot bacteria.

Key words: Single sight, inhibitory, synthetic, serial dilution, inoculation

1.0 Introduction

All over the world, current agricultural practices are moving away from the use of synthetic chemicals due to their adverse effects on ecosystem. The adoption of practices and use of plant materials that are environment friendly constitutes a vital component of sustainable agriculture. Nigeria is endowed with abundant plant materials, which may be employed to fight plant pests and diseases. This approach has recently been the pre-occupation of many crop protectionists in Nigeria. *Pectobacteriu* spp. George *et al.*, (1) formally known as *Erwinia* spp. induces soft rot on these crops both during the vegetation cycle and on the stored tubers and therefore considered the most threatening bacterium on potatoes and

other tubers worldwide (2; 4). The bacteria are ubiquitous plant pathogen with a wide host range (tubers, vegetables and horticultural crops) (4). The typical soft rot is induced mainly by a number of species coming under the genus *Pectobacterium*. The main species are *Pectobacterium carotovorum* subsp. *carotovora*, *Pectobacterium atroseptica* subsp. *atroseptica*, and *Pectobacterium chrysanthemi* subsp. *chrysanthemi*. These species produce pectolytic enzymes that hydrolyze pectin between individual plant cells. This causes cells to separate, a disease plant pathologist term bacterial soft rot (4). The disease (*pectobacterium ssp.*) is a major factor that limits production and transportation of roots and tubers as the disease is a threat both in the field, transits and store houses (Lojkowska and Kelman, (9). The use of protective chemicals on ware tubers is not advisable, it can render roots and tubers unfit for consumption as they undergo little processes before consumption. Hitherto, there is no effective control measure to manage this disease; therefore, selection of some plant extracts for the management the disease will be the preferred option. The objective of the study therefore was to evaluate some plant materials for the management of tuber soft rot bacteria induced by *pectobacterium ssp.*

2.0 Material and Methods

2.1 Experimental materials

The plant materials used were Neem seeds (*Azadirachta indica*) (A) Juss, Eucalyptus leaves (*Eucalyptus citrodorus* L), Lemon grass (*Cymbopogon citratus*) and Aloe vera (Liliaceae sub species aloinae) and Borax salt. The tubers used were Irish potato (*Solanum tuberosum*) (L), sweet potato (*Ipomoea batatas*) (L) Poir), white yam (*Dioscorea rotundata*) (L), cocoyam (*Xanthosoma sagittifolium*) (L) Schott, sweet cassava (*Manihot esculenta* Crantz). Borax salt was obtained from National Research Institute for Chemical Technology (NARICT) Zaria. Neem fruits, Eucalyptus leaves, Lemon grass, and Aloe vera were collected from the Institute for Agricultural Research (IAR), Ahmadu Bello University, Zaria. Ginger and garlic bulbs, as well as the various tubers were purchased from Samaru Market.

2.2 Preparation of plants extracts

Five grams of each of the plant material was weighed and pounded separately in a porcelain mortar to form a paste. This was then mixed with 100ml of water and allowed to stand for 6 h., after which they were filtered through Whatman No. 1 filter papers before filter-sterilizing them by means of Millipore filters (Millipore filter corporation, Bedford, Massachusetts USA).

2.3 Preparation of bacterial inoculum

A small portion of tissue at the advancing margin of the diseased tubers was removed with a sterile scalpel and dropped in McCartney bottles containing 9ml of sterile distilled water (SDW). These were allowed to stand for 5 minutes. The content was vigorously agitated and plated on Cuppels and Kelman's medium (5) and incubated for 24 h. at a temperature of ca. 27°C Cuppels and Kelman's medium contains 30 g sodium polypectate, 0.6 g $\text{CaCl}_2 \cdot 0$, 2 g NaNO_3 , 1.5 mg Crystal violet, 3 g Agar, 1liter of distilled water. After 24 h. incubation, wells were formed on the medium showing the presence of *Erwinia carotovora* subsp. *carotovora* (5). The pathogen was picked from edges of the wells with a heat-sterilized platinum wire loop and sub-cultured on sucrose peptone Agar (SPA) following the method described by Collins and Lyne (6), until pure cultures were obtained Sucrose peptone Agar (SPA) contains 20g sucrose, 20g agar, 5g peptone, 0.5 g $\text{K}_2\text{HP04}$, 1 liter of distilled water. A loopful of bacterial growth was taken from 24-h old cultures and suspend in 9 ml of SDW from which a serial dilution of up to 10^{-7} was made and used as stock. Confirmation of the identity of the pathogen in the culture was done through the ability of the pathogen to induce rot in potato tuber within 24 h.

2.3 Evaluation of plant extract

Two methods (Single site titration assay and Tuber slice assay) were used to evaluate the plants extracts for the control of bacterial soft rot

2.3.1 Single site titration assay

Tubers free of any obvious mechanical damage or disease were washed and then immersed for 5 min. in 0.5 % sodium hypochlorite (Clorox), rinsed with SDW, sprayed with 95% ethanol and allowed to dry at room temperature. The tubers were then bruised by sterilized knife at one site per tuber following the method described by Buelow *et al.* (7). Prior to inoculation, the tubers were immersed in the extracts for

10 min., after which they were allowed to dry for 15-20 min. When this had been done, 0.1ml of the bacterial suspension adjusted to ca. 5.0×10^7 cfu/ml was then introduced at the bruised site. Sterile polythene bags containing moistened filter papers and little SDW were arranged for each treatment. These were incubated at ca. 27°C for 48 h. Tubers immersed in SDW were used as control. The experiment was laid out using a Completely Randomized Design (CRD) with three replications. After 48 hrs tubers were Tuber sliced vertically through the Single site of inoculation and the width of decayed tissue was measured. Data collected were analyzed statistically using analysis of variance (ANOVA) and means separated by means of Student Newman Kuels (SNK) Test.

2.3.2 Tuber Slice Assay

Tubers were cut 20mm thick slices into following the procedure described by Dobias (8) and Lojkowska and Kelman (9). Surfaces of the tuber slices were disinfected as described above. Directly after slicing, a nick was made in the centre of each of the tuber slices. Treatment of the tuber slices with the plants extracts was done as described above. Eight sterile Petri-dishes containing filter papers moistened with SDW were arranged for the Tuber slices. Two Tuber slices were placed in each Petri-dish. One drop of bacterial suspension 0.1m containing 5.0×10^7 cfu/ml was used to inoculate each slice. The control Petri dishes were treated with SDW. These were incubated at ca. 27°C for 48 h. After 48 hrs the width of decay tissue were measured. The experiment was laid out using a CRD with three replications on a laboratory bench. Data collected were analyses statistically using analysis of variance and means separated by means of SNK.

2.4 Testing sensitivity of the organism to the extract

The plants extracts were prepared as described above. Eight sterile Petri dishes were arranged for the set up. SPA was prepared in a flask and allowed to cool to about 45°C. Before the introduction of the medium, 0.2 ml of the aqueous plant extract was introduced into the Petri-dishes with the aid of a micropipette, after which 20 ml of molten SPA was poured into the Petri dishes. Immediately after this was done, 0.1 ml of bacterial suspension adjusted to ca 5.0×10^7 cfu/ml was then introduced into the medium with the aid of yellow-tip micropipette. The contents of the Petri dishes were mixed thoroughly

by swirling five times each in clock wise, anti clock wise and north – south directions. The plates were then allowed to solidify and incubated immediately at ca 27°C. The experiment was laid out using a CRD with three replications the experiment was repeated twice. Colony counts (cfu) were done per plate after 48 h. Data collected were analyzed statistically using analysis of variance (ANOVA) and means were separated by means of SNK.

4.0 Results

The effect of plant extracts on the population of *Pectobacterium species in vitro* using plate count methods is shown in Table 1. All the plant extracts significantly reduced the population of the bacteria compared with the control. Population reduction was however, the greatest with neem extract, which is also not significantly different from those of garlic, ginger and lemon extracts. Eucalyptus had the least inhibitory effect on the bacterial population though not significantly different from treatment with *Aloe-vera* extract. The trend of results was similar for experiments I and II. In comparison with the control and all the plant extracts, borax salt had inhibitory effect similar to that of garlic, ginger, neem and lemon extracts.

The effect of plant extracts on the inhibition of rot induced by *Erwinia ssp.* using single site titration assay is outlined in Table 2. Among the plant extracts used on potato, lemon grass extract had the greatest inhibitory effect, but it is however not significantly different from the effects of garlic, *Aloe vera*, and neem extract. Eucalyptus and ginger extract had the least inhibitory effect on rot induction. In comparison to the control and the plant extracts, borax salt had the greatest inhibitory effect. However, its effect was not significantly different from that of ginger, garlic, lemon and neem. However, all the plant extracts significantly reduced the population of the bacteria compared to the control. Similar trends were observed on other tested tubers, in the tuber slice assay there was considerable variation among the tubers (Table 3). All the plant extracts significantly reduced the population of the bacteria compared to the untreated control. On potato, garlic had the greatest inhibitory effect but was statistically similar to the effects of neem extract. This was followed by lemon grass extract, whose inhibitory effect was not significantly different from those of *Aloe Vera* and ginger. Eucalyptus had the least inhibitory effect, but its effect was comparable to that of ginger extract. Borax salt had inhibitory effect comparable to *Aloe vera*, and lemon.

On yam and sweet potato, garlic extract had the greatest inhibitory effect though its effect was not significantly different from those of Borax, lemon, Aloe vera, and neem extract. These were followed by ginger extract. Apart from the control, Eucalyptus had the least inhibitory effect. This result was similar on all the other tubers except on cassava where lemon grass and neem extracts had similar inhibitory effects.

Table 1: Effect of plant extracts on the population of *Erwinia* ssp. *in vitro* using dilution plate count method

Treatment	Mean bacterial count (cfu)/ml*	
	Experiment 1	Experiment II
Eucalyptus	2.5×10^7 ^b	2.4×10^7 ^b
Neem	7.0×10^6 ^c	7.8×10^6 ^c
Lemon	1.2×10^6 ^c	1.1×10^6 ^c
Borax	7.1×10^6 ^c	7.0×10^6 ^c
<i>Aloe vera</i>	2.3×10^7 ^b	2.3×10^7 ^b
Garlic	8.0×10^6 ^c	7.0×10^6 ^c
Ginger	1.1×10^7 ^c	1.2×10^7 ^c
Control	7.7×10^8 ^a	7.2×10^8 ^a

Means followed by the same letter are not significantly different at P = 0.05 using SNK test.

Table 2: Inhibition of rot by plant extracts (diameter of rot/mm) in the single site titration assay

Plant Extracts	Potato	Yam	Sweet potato	Cassava	Cocoyam
Eucalyptus	3.80 ^b	3.83 ^b	3.73 ^b	3.25 ^b	2.63 ^b
Neem	1.42 ^c	1.42 ^c	1.00 ^c	1.17 ^c	0.50 ^c
Lemon	0.95 ^c	0.67 ^c	0.95 ^c	0.42 ^c	0.58 ^c
Borax	0.67 ^c	0.50 ^c	0.33 ^c	0.58 ^c	0.17 ^c
<i>Aloe vera</i>	1.17 ^c	0.83 ^c	0.92 ^c	0.92 ^c	0.50 ^c
Garlic	1.00 ^c	0.75 ^c	0.67 ^c	0.58 ^c	0.33 ^c
Ginger	3.42 ^b	3.00 ^b	2.97 ^b	2.83 ^b	2.12 ^b
Control	18.83 ^a	9.17 ^a	11.00 ^a	8.50 ^a	8.50 ^a

Means followed by the same letter are not significantly different at P = 0.05 using SNK test.

Table 3: Inhibition of rot by plant extracts (diameter of rot/mm) in the tuber slice assay

Plant Extracts	Irish potato	Yam	Sweet potato	Cassava	Cocoyam
Eucalyptus	8.67 ^b	7.67 ^b	8.00 ^b	6.67 ^b	5.17 ^b
Neem	3.00 ^{de}	2.85 ^d	2.00 ^d	2.33 ^{cd}	1.33 ^d
Lemon	4.17 ^{cd}	2.00 ^d	2.33 ^d	2.33 ^{cd}	0.83 ^d
Borax	4.21 ^{cd}	1.00 ^d	2.50 ^d	1.17 ^d	0.50 ^d
<i>Aloe vera</i>	4.33 ^{cd}	0.33 ^d	2.00 ^d	1.33 ^d	0.67 ^d
Garlic	1.33 ^e	0.83 ^d	0.67 ^d	0.67 ^d	0.83 ^d
Ginger	6.67 ^b	5.50 ^c	5.50 ^c	4.33 ^c	3.50 ^c
Control	18.83 ^a	11.50 ^a	11.33 ^a	9.67 ^a	9.00 ^a

Means followed by the same letter are not significantly different at P = 0.05 using SNK

5.0 Discussion

The variability of results observed under the tuber slice assay, could not be attributed to the various extracts used but rather was due to a number of factors that may modify the reaction of tubers following slicing. For example, the location in the tubers from which tuber slices were cut may influence the susceptibility of tuber slices to the rot pathogen. The location on tuber like the cortex and medullary tissue has different resistance to rot pathogen (9).

The experiment has shown the effectiveness of lemon grass (*C. citratus*) in reducing the population of the pathogen both *in vitro* and on the treated tubers. The insecticidal and antimicrobial activity of lemon grass against many insects and pathogens has been reported by other authors (10 and 11). The antimicrobial properties could be attributed to the presence of 80% citral, citronella, geraniol, methyl-heptone, n-decyl-aldehyde and linanol in the plants, all of which are active ingredients used in the manufacture of pesticides (11).

Also confirmed was the antimicrobial activity of garlic against a broad spectrum of fungal and bacterial organisms as reported by other authors (12, 13, 14, 15 and 16). The antimicrobial properties of garlic have been attributed to the presence of an essential oil that contains allyl disulphide ($C_6H_2 S_2$) diallyl disulphide ($C_6 H_{10} S_2$) and two or more sulphur-containing compounds (16 and 17).

The wide range of antimicrobial activity of *Aloe vera* has been confirmed by this work. It significantly reduced the population of the bacteria both *in vitro* and on the tested tubers compared to control. The active principles are mainly anthroquinone, lignin, salicylic acid, anthrax quinine, aloin and emodin, which acts as painkillers; they also function as antibacterial and antivirals (18 and 19).

Furthermore the experiment confirmed the antimicrobial activity of neem, which is in agreement with the findings of previous workers (20, 21, 22, 23 and 24). This is however contrary to the report of Shenge

(16) that neem seed extract was not effective in reducing the population of (*Xanthomonas campestris* pv. *malvacearum*).

The antimicrobial activity of Eucalyptus leaves could be attributed to the presence of 0.5–3.5% volatile oil, tannins, polyphenolic acid, caffeic, ferulic, gentisic, flavonoids, quercetin, tannin, hyperoside and eucalyptin (25). Eucalyptus oil and eucalyptol reportedly have strong antibacterial properties against several strains of *Streptococcus* (26).

Finally, the experiment has also shown the effectiveness of ginger in significantly reducing the population of the bacterial soft rot both *in vitro* and on tested tubers. This could be as a result of the Zingerone substance in the plant which is the active principle (27).

6.0 Conclusion

Using plant material to control pests may alleviate the burden of heavy reliance on synthetic pesticides. However, given the low mammalian toxicity of the plant materials, they may be appropriate for low impact structural pest management as part of a complete IPM approach. These practices are labor intensive, but are economically and ecologically sound since it does not require sophisticated technology and it is appropriate for resource poor farmers in the developing countries.

7.0 REFERENCES

1. George, M., A. Julia., A. Bell., Timothy G. L. Taxonomic outline of the prokaryotes systematic bacteriology. 2nd Edition. Pp 310. 2004.
2. Hadas, R.; G. Kritzman, T. Gefen and Manulis S. Detection, quantification and characterization of *Erwinia carotovora* spp. *Carotovora* contaminating pepper seeds *Plant pathology*. 2001. (50), 117-123.
3. Lowe, J. Polyalacturonase deficient strain of *Erwinia carotovora* ssp. *atropurpurea* laes attenuated virulence. In: Lemattre M, Freigoums, plant pathogenic Bacteria. Les colloques de l'INRA 66 Paris France. INRA Editions 585-589. 1992..
4. Tavasoli, E.; A. R. Marafat and Hassazadah, N. Identity and genetic diversity of *Pectobacterium* spp., causal agents of potato soft rot in Zanjan, Iran. *Africa journal of plant science* 2011. 5(6) 329-336.
5. Cupels, D.A and Kelman, A. Evaluation of selective media for isolation of soft rot bacteria from soil and plant tissues. *Phytopathology* 1974. 64:468-475.
6. Collins, C.H. and Lyne, P.M. *Microbiological Methods*, Fourth Edition. Butterworth and Co. Ltd, England, p. 142-143. 1976.

- 238 7. Buelow, F. H., Mahar, E.A and Kelman, A. Assessment of bacterial soft rot potential in potatoes. In
239 Cargill, B. F. (ed.) *Engineering for Potatoes*, Michigan state University and America society of
240 Agricultural Engineers; St. Joseph, M.I., p. 440-455. 1986.
- 241 8. Dobias, K. Methoden Zur prufung der resistenz von karoflen gegen den erreger der knollennass
242 faull Tag-Ber, a kad land wirsth-wiss. 1976. 140:221-230.
- 243
- 244 9. Lojkowska, E and A. Kelman. Comparison of the effectiveness of different methods of
245 Screening for bacterial soft rot resistance of potato tubers. *American Potato Journal*, 1994. 71:99-
246 112.
- 247 10. Jija, M.J. Effect of lemon grass (*Cymbopogon citratus*) and black sesame (*Hyptis Spicigera*
248 products on *Callosobruchus maculatus* on stored cowpea. M. Sc. thesis submitted to the
249 Department of Crop Protection, Ahmadu Bello University, Zaria pp.81. 1999.
- 250 11. Oparaeke, A.M. Collection, Identification and screening of indigenous herbal extracts and waste
251 matter for control of insect pests of cowpea (*Vigna unguiculata* L.) Walp, Ph. D. Thesis
252 Department of crop protection, A.B.U., Zaria, 158pp. 2004.
- 253 12 Ahmed, S.R. and Agnihotri, J.P. Antifungal activity of some plant extracts. *Indian Journal of*
254 *mycology and plant pathology*, 1977.7 (2): 180-181.
- 255 13. Agrawal, P. Effect of root and bulb extracts of *Allium* spp. on fungal growth. *Transactions of the*
256 *British mycological society*, 1978. 70(3): 439-441.
- 257 14. Shashikanth, K.N., S.C. Basapa and V.S. Murthy. Studies on the antimicrobial and stimulatory
258 factors of garlic (*Allium sativum*) *Journal of food science and technology*. India 1981. 18(2):
259 44-47
- 260 15. Gondwe, B.R., R. Mabagala and L.A Femi. Effect of garlic extracts on growth of *Pseudomonas*
261 *syringae* pv. *Phaseolicola* in culture media. *Bean Improvement Cooperative Report*, 1996. 29:280-
262 281.
- 263 16. Shenge, K.C. Evaluation of plant extracts for the control of bacterial blight of cotton induced by
264 *Xanthomonas compestris* PV. *Malvacearum* (Smith) Dye. M.Sc theses Ahmadu Bello University
265 Zaria. Pp 79. 2002. .
- 266 17. Obagwu.J. Developing Biopesticides for control of citrus fruit pathogens of importance in Global
267 trade. Ph. D. Thesis, Faculty of Natural And Agricultural Sciences, University Of Pretoria, 166
268 pp. 2003.
- 269 18. Amodioha, A.C. Evaluation of some plant extracts against *Colletotrichum lindemuthianum*. *Archives*
270 *of Phytopathology And Plant Protection*, 1999. 32:141-149.
- 271
- 272 19. Ahmed. I. and Beg, A. Anti Microbial Photochemical studies on 45 Indian medicinal plants against
273 multi-drug resistant human pathogen. *Journal of Ethnopharmacological*, 2001. 7: 113-123.
- 274

- 275 20. Emechebe, A.M. Evaluation of aqueous extracts of parts of some plants and black local soap
276 solution for the control of cowpea diseases at Samaru. Cropping Scheme Meeting: Report On
277 Legumes And Oil Seeds Research Programme 26th February-1st March, 1996. Pp.79. 1996.
- 278 21. Obagwu. J. Studies on the brown blotch disease of bambara nut (*Vigna subteranea* (L.) verdc)
279 induced by *Colletotrichum capsici* (syd) Bultter and Bisby in the Zaria area of Northern Nigeria.
280 M. Sc. Thesis, A.B.U., Zaria. 105. 1997.
281
- 282 22. Emechebe, A.M. and Alabi, O. Evaluation of aqueous extracts of parts of some plants for the control
283 of cowpea diseases at Samaru. *Cropping Scheme Meeting Report on Legume and Oil Seeds*
284 *Research Programme*, 26th February – 1st March 1997, p. 77. 1997.
- 285 23. Stoll, G. Natural Crop Protection in the tropics pp 188. 1998.
- 286 24. Aliyu, S. Effect of plant extracts on the germination and establishment of sorghum seed and
287 seedlings in the glasshouse. B.Sc. Thesis, Ahmadu Bello University, Zaria, 27pp. 1999.
- 288 25. Lawrence, B.M. Biological activities of *Eucalyptus* spp and its perfume quality
289 *Chemical Technology*, 1990. 15:36-45.
290
- 291 26. Brieskon C. H. and W. Schlicht. Traditional medicine. *Pharm. Acta Helv.* 1976. 51: 133-138
- 292 27. Epidi, T.T, Alamene, A and Onuegbu, B.A. Influenecce of some plant extract on yield and insect
293 pests of cowpea (*Vigna unguiculata* (L) walp.) *Nigeria journal of plant protection*, 2005. 22 65-
294 76.
- 295
296
297