

1
2 **EFFECT OF PHOSPHORUS AND RHIZOBIA INOCULATION ON GROWTH AND**
3 **YIELD OF SOYBEAN (*Glycine max* L. Merrill)**
4

5 **MM Akpalu^{1*}, H. Siewobr¹, D. Oppong-Sekyere¹ and S.E. Akpalu²**

6 ¹ *Bolgatanga Polytechnic, Department of Ecological Agriculture, P. O. Box 767,*
7 *Bolgatanga, Upper East Region, Ghana*

8 ² *CSIR-FORIG, Bolgatanga P. O. Box 142, Bolgatanga, Upper East Region, Ghana*
9

10 **ABSTRACT**

11 **An experiment was conducted in the major and minor cropping seasons of 2012 and 2013**
12 **under field conditions at Bolgatanga Polytechnic, to study the effect of phosphorus**
13 **fertilizer and Rhizobia inoculation on growth and yield of soybean. The soybean variety**
14 **used was ‘Jenguma HSCII51’. The seed, Triple Superphosphate fertilizer and the Rhizobia**
15 **inoculant were obtained from the Ministry of Food and Agriculture (MOFA), Navrongo. The**
16 **experiment was laid in a Randomized Complete Block Design (RCBD) with three**
17 **replications. The treatments were: Soybean + phosphorus fertilizer + Rhizobia inoculation**
18 **(+P/+I), Soybean + phosphorus fertilizer only (+P/-I), Soybean inoculated with Rhizobia (-**
19 **P/+I) and the control-Soybean only (-P/-I). Parameters measured included: plant height,**
20 **nodule number, pod number per meter square, number of seeds/pods, 100 - seed weight,**
21 **shelling %, grain yield and harvest index. Results indicated that Phosphorus fertilizer was**
22 **a requirement for shoot growth, pod and seed yield. Nodulation and root growth were**
23 **significantly increased by Phosphorus + Rhizobia inoculation (+P/+I), but P fertilizer only**
24 **did not enhance root growth. Dry matter accumulation was highest between onset of**
25 **flowering and Podding (42 to 70 DAP), irrespective of the treatments.**
26 **Phosphorus fertilizer treatment only (+P/-I) and Rhizobia inoculation plus phosphorus**
27 **fertilizer (+P/+I) produced significantly more pods (3903.7 Kg/ha) and (3858.3 Kg/ha)**
28 **respectively than the Rhizobia inoculation only which recorded (2301.0 Kg/ha). The**
29 **control recorded grain yield of (2130.7 Kg/ha). Grain yield was again highest for Rhizobia**

*Corresponding author: E-mail: aa.marilyn@yahoo.com, Tel: +233(0)208134403

inoculation plus Phosphorus fertilizer (+P/+I) and Phosphorus fertilizer only (+P/-I) recording 7.61 t/ha and 7.30 t/ha respectively while Rhizobia inoculation only (-P/+I) and the control (-P/-I) produced the lowest grain yield (4.41 t/ha) and (3.80 t/ha) respectively. The treatments **did not have any significant effect** on number of seeds per pod, 100 seed weight and harvest index (HI). Shelling percentage was significant for Phosphorus fertilizer treatment only (+P/-I) (69%) and the Rhizobia inoculation only (-P/+I) (67%) but **was not significant** for the control (-P/-I) (68%) and Rhizobia plus phosphorus (+P/+I) (67%).

Keywords: *Cultivation, fertilizer, harvest index, inoculation, nodulation, phosphorus, rhizobia, soybean*

INTRODUCTION

Soybeans (*Glycine max L. merr*) is an ancient leguminous crop and possibly originated from temperate China where it was believed to be one of the five sacred grains which the county's ancient civilization depended on (Tweneboah, 2000). About 70% of the world's soybean is now produced in the USA, with the mainland China being the second largest producer. In terms of production, consumption and trade, soybean is the world's most important grain legume (Tweneboah, 2000).

Soybean is relatively a recommended crop in Ghana, referred to as the 'golden bean', 'miracle bean' and in some areas, the crop of the 'planet'. It was initially introduced into the country in 1909 for export and for animal feed but failed due to poor understanding of its cultivation and utilization; it had been tried a few more times after this in 1930s, 1950s, 1970s and the 1990s but had failed due to several reasons (such as the rainfall pattern, soil type, technique of cultivation) (Woomer, 2010).

Soybeans have many industrial uses and are important sources of vegetable oil and protein. The beans has an oil content of about 20%, protein content of 40-45% carbohydrate, 30 % and total sugars of about 10% (Haas et al. 2006). It is also a good source of calcium, phosphorus, copper, potassium, magnesium and thiamine. [Van Gerpen and Krahl \(2005\)](#); Haas *et al.* (2006) stated that soybean oils can be converted to biodiesel and glycerol. In the US, the major biofuel for transportation is soybean biodiesel, which displaces petroleum and diesel.

Soybean has a great potential for the alleviation of Protein Energy Malnutrition (PEM) in Ghana because animal sources of proteins are scarce and expensive. In Ghana, soybeans is used in the preparation of soy paste, soy milk, soy custard, soy kebab, soy cheese, soy palaver sauce and many others.

Soybean is produced by small holder farmers in Ghana, under rain fed conditions. Under these subsistence methods, soybean yield averagely 0.8 metric ton per hectare. FAO estimates that Ghana imports 2700 gallons of soybean oil. There is limited supply of soybean feed for the poultry industry in Ghana resulting in major constraints in the growth of the industry. This has resulted in significant growth of import of poultry and other meat products (protein sources) for consumption (MiDA, 2013).

Poor soils has become a problem as a result of poor land management which leads to continuous loss of soil nutrients through the activities of soil erosion, by rain, leaching, crop removal and bad cultural practices. The resultant effect is the decline in crop yields. Ohyama *et al* (2013), stated that soybean seed yield can reach 4.6 t/ha with well managed fields under good climatic and soil conditions. The highest yield of soybean was recorded in Japan at 7.8 t/ha (Ohyama *et al.*, 2013). Recently, an amazing high soybean seed yield over 10 t/ha was recorded in 2008 and 2010 by a farmer in Missouri, USA. Therefore the potential productivity of soybean should be higher than we have thought (Ohyama *et al.*, 2013).

In view of the above, measures must be taken to improve BNF because when nitrogen is lost from soils; leguminous plants however, are not able to convert atmospheric N into absorbable forms of NH_4^+ and NO_3^- before used by plants.

Natural reserves of soil N are normally low in many African soils and again phosphorus fertilizers which are important elements necessary for plants' growth are also lacking in many African soils (Hardarson *et al.* 1987). Since the crop has immeasurable potential in Africa at large and Ghana in particular, this study was conducted to provide useful information on the effects of P fertilizer application on nodulation, shoot growth and yield; to evaluate the effects of Rhizobia inoculation on nodulation and yield.

The need for Phosphorus and rhizobia inoculation treatment, particularly to a crop such as soybean, has become essential due to the immeasurable benefits the crop provides; income to resource poor farmers, an important source of protein, and its ability to fix atmospheric nitrogen to the soil.

Phosphorus is an important requirement for shoot growth and the subsequent yield of soybean. Phosphorus and Rhizobia application (to soybean) is reported to increase nodulation, pod formation and development, and a subsequent grain yield (Ajeigbe and Woomer, 2010; Cassman *et al.*, 1981).

It is therefore envisaged the study would provide an important first step at improving the crop in Ghana, particularly the North, where there is seasonal food scarcity, and also serve as a foundation to enhance their potential cultivation and use in future, thus contributing to food security.

MATERIALS AND METHODS

The study was conducted in the major and minor cropping seasons of 2012 and repeated in 2013, at the experimental field of the Bolgatanga Polytechnic which resides in Sumbrungu, located 7 km away from Navrongo. It shares boundaries with Kassena-Nankana district at the north, Bongo district at the East, south to Bolgatanga Municipality and West to Sirigu and lies between latitude 10°45' North and longitude 0°51' West of Ghana.

The land was slashed, ploughed and harrowed, and then finally raked, lined and pegged before planting at the appropriate spacing; the experiment was laid in a Randomized Complete Block Design (RCBD) with three replications. Seeds were planted in rows on flat field; spaced 5cm x 75cm apart, 3 seeds were planted per hill and were not thinned. Each plot in a block measured 3m x 3m and was contiguous to one another; there were 4 plots in each replication. The distance between the blocks was 1 meter making a total of 12 plots.

TREATMENTS: A - Control plot which is planted to soybean only (-P/-I). B - Plot planted to soybean seed which is inoculated (-P/+I). C - Plot planted to soybean and Triple Superphosphate (TSP) is applied and inoculated (+P/+I). D - Plot planted to soybean and TSP is applied (+P/-I)

SOYBEAN INOCULATION WITH RHIZOBIA

One (1 kg) of soybean seed was weighed and placed in a rubber container, 3 spoonful of sugar solution was added as an adhesive and 10 g of inoculants weighed and added to the soybean seeds. It was then stirred with a spoon until the seeds were evenly coated with the Rhizobia. After coating, the seeds were spread onto a clean surface and air dried under a shade for about one hour. The inoculated air dried seeds were then planted, three seeds per stand.

FERTILIZER APPLICATION

Triple Superphosphate (TSP) fertilizer was applied after two weeks at a rate of 139.4 g per plot in a band, 10cm away from the planting line in a 2cm deep trench and covered with moist soil after application. The application was done after seedlings developed ten to eleven leaves to enable effective use of the fertilizer.

WEED AND PESTS MANAGEMENT

All plots were weeded by hand (hoe), at least three times. The Insecticide, Cyperdicot (Cypermethrin); SPCT as an active ingredient, was used, at a recommended rate of 25ml per half knapsack sprayer, to control insects, particularly at the flowering stage.

Growth Parameters measured included plants height, number of leaves, number of nodules, leaf dry weight, stem dry weight, roots dry weight and nodule dry weight.

Yield Parameters measured also included number of pods per plant, number of pods per meter square, number of plants per meter square number of seeds per pods, 100 seeds weight and mean seed weight.

Averages, range, standard deviation and coefficient of variation were computed for the measurement data. Quantitative data was subjected to analysis of variance (ANOVA) using GenStat discovery edition 4.0 (GenStat 5 Committee, 2000). Means were separated by least significant difference at 5%.

RESULTS AND DISCUSSION

Table 1: Average Climatic data for 2012 and 2013 (Upper East Region)

Months	Average Temperature (°c)	Total Rainfall (mm)	Sunshine / hrs	Relative Humidity (%)
January	27.5	0.0	9.5	23.0
February	30.7	0.9	7.9	24.3
March	33.1	0.0	6.5	24.8
April	31.8	99.9	6.7	54.0
May	29.1	154.4	7.4	70.8
June	28.9	79.3	8.1	69.0
July	27.1	241.9	6.3	78.5
August	26.7	128.8	6.8	80.3
September	26.9	235.1	5.8	80.0
October	28.5	47.7	7.7	71.5
November	30.1	0.0	9.9	50.3
December	28.6	0.0	9.5	27.5

Source: Navrongo Meteorological Station, [year 2012 and 2013](#)

Table 2: Chemical Properties of Soil at the Experimental Site

Chemical Composition	(0-15 cm) depth
pH	6.1
*Available P (mg/kg)	2.3
Organic Carbon (%)	0.36
Organic matter (%)	0.62
Total N (%)	0.03
Exchangeable Bases (Cmol/kg/Me/100g)	
Ca	3.2
K	0.1
Mg	1.34
Na	0.04

**Available P (mg/kg), from 20.0 mg/kg and above is = Adequate to high*

EMERGENCE AND FLOWERING

Emergence was complete within 10 days after planting; all the treatments recorded 50% emergence at 4 days. The Rhizobia inoculated treatment significantly increased the number of seedlings that emerged, but there was no significant difference between the rest of the treatments, that is P fertilizer (+P/-I) application, the control (-P/-I) and both P and Rhizobia inoculated treatment (+P/+I) as shown in Table 3. This may be attributed to inoculants' effect or Rhizobia effect which enhanced emergence and germination. However, the treatments did not have any effect on 50% flowering. All the treatments took 47days to attain 50% flowering. Nevertheless, Rhizobia inoculation only recorded

the highest number of flowers of 281 followed by the other treatments and this has a relationship with emergence (Table 3).

Table 3: Effect of phosphorus and Rhizobia inoculation on emergence and flowering of soybean

Treatments	50 % Seedling Emergence (4 days)	50% flowering (47 days)
(-P/-I) No Inoculation and No P fertilizer	78.1	238
(-P/+I) Rhizobia inoculation only	109.3	281
(+P/+I) Rhizobia Inoculation + P fertilizer	80.3	264
(+P/-I) P fertilizer Only	76.0	233
Lsd (5%)	28.7	99.1
CV (%)	19.4	19.1

NS = Not Significant, P > 0.05

LEAF DRY WEIGHT

In all the treatments, leaf dry weight increased to a peak 84 DAP and declined at 70 DAP owing to onset of senescence (Figure 1). Phosphorus plus Rhizobia inoculation recorded the highest leaf dry weight of 50.7 g/plant and was followed by phosphorus application only which recorded a weight of 36.4 g/plant and peaked at 70 DAP. Rhizobia inoculation alone also took the same trend, recording a leaf dry weight of 29.5 g/plant. The control (-P/-I) recorded the lowest leaf weight of 20.70 g/plt as shown in Figure 1.

The results showed that the phosphorus application plus Rhizobia inoculation produced the greatest shoot dry matter. Field observations showed that phosphorus and rhizobia inoculation enhanced dark green colour of the leaves. It appears, therefore that rhizobia inoculation enhances P uptake leading to the development of a larger photosynthetic area. Phosphorus application has been reported to increase shoot growth or leaf growth of soybeans and other leguminous crops (Othman, 1991; Cadish *et al.*, 1992).




Figure 1: The effect of phosphorus and Rhizobia inoculation on leaf dry weight of soybean.

STEM GROWTH

Stem dry weight also followed the same trend as leaf growth but the peak stem growth was recorded at 70 DAP and at 84 DAP for all the treatments except Rhizobia inoculation only which peaked 56 DAP and remained sturdy until 70 DAP and dropped (Figure 2). Initial stems growths experienced almost the same dry matter accumulation and on 28 DAP, all the treatment begun to split as a result of stem dry weight differences. Phosphorus plus Rhizobia inoculation (+P/+I) recorded the highest dry matter residues to point at 56 DAP (Figure 2). On 56 DAP, there was stability in dry matter accumulation to 70 DAP which marked the period of vegetative growth, flowering and pods development, therefore dry matter accumulation was more rapid between 28 and 56 DAP, hence the steady weight drop because soybean stems loss their green

colour as the crop matures or senescence sets in since the level of stem enlargement ceases at onset of reproductive growth.

The acute deficiency of phosphorus can prevent nodulation by legumes. Phosphorus is required for nodule metabolism and tends to be concentrated in the nodules when the plant is deficient in these nutrients (Hara *et al.*, 1988; [Cassman et al., 1981](#)).




Figure 2: Effect of phosphorus and Rhizobia inoculation on stem growth of soybeans.

ROOT GROWTH

Comparing root growth of the treatments, the Rhizobia plus phosphorus (+P/+I) treatment again produced the highest root dry matter of 6.93 g/plt at 56 DAP. The Rhizobia inoculated treatment alone recorded the next higher value of 6.17 g/plt but at 70 DAP when the (+P/+I) treatment was gradually dropping (Figure 3). The growth may be attributed to increase in cell dry matter accumulation with well-developed tap root during growth of the plant. The control was the next highest, while phosphorus application only increased root growth up to 42 DAP and began to drop. Rhizobia inoculation might have led to increased root growth of the plant. [This finding is in line with Okagun et al., \(2004\) who observed a significant increase in root dry matter yield of inoculated improved soybean over uninoculated improved variety.](#)

The drop in root growth at 84 DAP may be due to senescence. Plant roots get dried in the soil and lose its weight as the crop matures. Therefore, as the plants mature, the

more it loses dry matter leading to significant drop in dry weight at different growth stages leading to senescence.




Figure 3: The effect of phosphorus and Rhizobia inoculation on root growth of soybean.

NUMBER OF NODULES

From Figure 4, on 14 days-after-planting, treatment (+P/-I) recorded a slightly sharp increase in number of nodules produced per plant. This may be due to bacteria inoculation having been less effective in nodulating at the second week of growth. The ineffectiveness of the Rhizobia bacteria during the two weeks was due to the period of bacteria incubation in the soil, hence at this stage, plant roots, after two weeks, did not use these bacteria much, and therefore recorded less number of nodules produced as shown in (figure 4).

On the other hand the control treatment (-P/-I) recorded the lowest number of nodules showing that the soil did not contain much bacteria to produce more nodules at this stage, but the application of P increased nodulation. On 28 days after planting, phosphorus plus rhizobia inoculation (+P/+I) treatments recorded a tremendous increase in number of nodules produced. Higher values were recorded again for the same

treatments at 42 and 56 days after planting (Figure 4), indicating effective utilization of the bacteria and fertilizer.

On 70 days after planting, there was a sharp drop in the number of nodules recorded as a result of the use of more energy to convert protein into food reserves that the plants required, therefore, nodules number degenerated or disappeared indicating the point of maturity. This agrees with the research findings of [Fournier et al, \(2008\)](#).




Figure 4: The effect of phosphorus and Rhizobia inoculation on of nodule number of soybean.

NODULE GROWTH

At different growth stages, there was [no significant increase](#) in Rhizobia effect for the treatment (-P/-I) and (-P/+I) on nodulation at 14 DAP, but treatments (+P/+I) and (+P/-I) had attained a slight increase of (0.3g/plt) and (0.29g/plt) respectively (Figure 5). The increase in growth of nodules was equivalent to increased nodule dry weight of the plants. Therefore, on 28 DAP, (+P/+I) had a significant increase in dry weight of 1.3g/plt and treatment (+P/-I) with 0.8g/plt. Nodulation therefore increased between the pre-flowering and early pod-forming stages. This suggests that, after the flowering stage, the main nitrogen-fixing activity of the whole root system reduced. The results also indicated that it was necessary to analyse the origin of nodules on the various parts of the root

system independently in order to establish clearly the effectiveness of the Rhizobia inoculation in terms of nodulation.

Several studies have reported that soybean grown under field conditions and inoculated with the rhizobia sometimes failed to show a positive response to inoculation, in terms of nitrogen fixation and crop yield (Weaver and Frederick, 1974; Cassman et al., 1981).

On 28 DAP, treatment (+P/+I) recorded an increase in dry matter residue while treatment (+P/-I) recorded the second highest dry weight of nodules produced (Figure 5). Treatments (-P/-I) and (-P/+I) obtained the lowest of 56 DAP, but as the trend continued, there was a slow drop in weight to about 1.0 g/plts for (+P/+I) and 0.83 weight g/plt for (+P/-I). This may be as a result of plant senescence or ageing because much energy is absorbed to forming pods.




Figure 5: Effect of phosphorus and rhizobia inoculation on nodule growth of soybean.

EFFECT OF PHOSPHORUS AND RHIZOBIA INOCULATION ON POD NUMBER, NUMBER OF SEEDS PER POD, 100 SEED WEIGHT AND SHELLING %.

Phosphorus fertilizer treatment only (+P/-I) and Rhizobia inoculation plus phosphorus (+P/+I) produced significantly more pods (3903.7 Kg/ha) and (3858.3 Kg/ha) than the rhizobia inoculation only which recorded 2301.0 Kg/ha and the control (2130.7 Kg/ha). But there was no significant difference between the phosphorus fertilizer treatment only

(+P/-I) and Rhizobia inoculation plus phosphorus (+P/+I). There was again no significant difference between the control and the Rhizobia inoculated treatment only (Table 4). Phosphorus fertilizer application therefore had an influence on pod formation and pod number, and P availability also encourages nodulation; this was the reflection on the high number of pods produced by the rhizobia inoculation plus phosphorus (+P/+I) treatment but rhizobia inoculation alone recorded a lower pod number. However, [the treatments did not have any significant effect on number of seeds](#) per pod and 100 seed weight (Table 4). But there were significant effects on shelling percentage with the P fertilizer treatment only.

Treatment (+P/-I) recording the highest shelling percentage of 69% followed by the control. It shows; therefore, that yield of soybean depended on pod number per unit area. This finding is in line with that of Ohyama *et al.* (2013) who stated that soybean yield depends mostly on pod number per unit area ([Cassman *et al.*, 1981](#)).

Table 4: Effect of phosphorus and rhizobia inoculation on yield components of soybean (Jenguma HSCII51); at final harvest.

Treatments	Pod No./m ²	Number of seeds/pod	100 Seed wt	Shelling %
(-P/-I) No Inoculation and No P fertilizer	2130.7	2.000	10.67	0.68
(-P/+I) Rhizobia inoculation only	2301.0	1.833	10.83	0.66
(+P/+I) Rhizobia Inoculation + P fertilizer	3858.3	1.967	10.97	0.67
(+P/-I) P fertilizer Only	3903.7	1.900	11.60	0.69
Lsd (5%)	1278.5	NS	NS	0.02
CV (%)	10.5	3.4	1.8	0.9

[NS = Not Significant, P > 0.05](#)

EFFECT OF PHOSPHORUS AND RHIZOBIA INOCULATION ON SEED YIELD COMPONENTS OF SOYBEAN.

Seed yield is determined by number of plants per unit area, the mean number of pods per plant, the mean number of seeds per pod and the mean individual seed weight.

Rhizobia inoculation plus P fertilizer (+P/+I) and P fertilizer only (+P/-I) Once again recorded significantly the highest pod dry weight of (1131.1 Kg/ha) and (1054.8 Kg/ha) respectively. But there was no significant difference between the two.

Again, there was no significant difference between the control (-P/-I) and the Rhizobia inoculation only (-P/+I) (Table 5).

Grain yield was highest for Rhizobia inoculation + P fertilizer (+P/+I) and P fertilizer only (+P/-I), recording 7.61 t/ha and 7.30 t/ha respectively, while Rhizobia inoculation only and the control recorded the lowest grain yields of 4.41 t/ha and 3.80 t/ha respectively (Table 4). Ohyama *et al.* (2013), reported that soybean seed yield can reach 4-6 t/ha with well a managed field under good climatic and soil conditions. They again stated that the highest yield of soybean of 7.8 t/ha were recorded in Japan. The yield obtained in this study was much close to that recorded in Japan and this could be considered as the highest yield reportedly obtained in the current study.

The control and the Rhizobia inoculation (-P/+I) treatments, did not have any significant increase on grain yield; hence phosphorus fertilizer can be said to have a greater influence on grain yield of soybean. This is because the impact of phosphorus fertilizer was significant across board in terms of dry matter produced, pod yield and grain yield. According to the soil analysis results of the experimental site (Table 2), the P content of the soil was very low (2.3 mg/kg) as compared to 20 mg/kg regarded as adequate or the standard.

Adequate phosphorus application and Rhizobia will lead to effective pod dry matter accumulation, and subsequently to higher grain yield.

The yield characteristics are a reflection of shoot dry weight which was increased by P fertilizer application. This may indicate that when shoot growth is limited by P unavailability, pod filling (Shelling %) and total seed yield will be accordingly limited. This

shows the importance of P supply in improving the production of soybean in Ghana. The poor yields obtained by farmers may be attributed to P unavailability in most soils particularly during the period of high shoot growth rate (42 to 70 DAP).

Table 5: Effect of phosphorus and Rhizobia inoculation on pod dry weight, grain yield and harvest index of soybean

Treatments	Pod Dry wt. (g/m²)	Grain Yield (g/m²)	Grain Yield (t/ha)	Harvest Index
(-P/-I) No Inoculation and No P fertilizer	562.2	379.9	3.80	0.54
(-P/+I) Rhizobia inoculation only	668.7	441.7	4.41	0.52
(+P/+I) Rhizobia Inoculation + P fertilizer	1131.1	763.6	7.61	0.52
(+P/-I) P fertilizer Only	1054.8	727.9	7.30	0.50
Lsd (5%)	385.2	253.6	0.30	NS
CV (%)	6.0	6.2	6.2	3.8

NS = Not Significant, P > 0.05

CONCLUSION AND IMPLICATIONS OF THE STUDY

The results have shown P requirement for shoot growth and yield of soybean. Based on the study, phosphorus and Rhizobia application to soybean played a very significant role in increasing nodulation, pod formation and development, and a subsequent grain yield.

Soybean is a relatively new crop in Ghana. Presently, inoculation is not largely practiced by farmers.

Promiscuous soybean cultivars were proposed as alternatives to the use of inoculants.

Nonetheless, there seems to be P requirement for shoot growth, nodule formation and yield.

Conversely, most savannah soils, lack of P; under the low-input agricultural systems, farmers do not apply fertilizers to soybean, they generally use the limited amount of fertilizer available for staple cereal crops like millet, sorghum, and rice and maize.

This study has shown that in order to derive full benefit from biological nitrogen fixation and hence to maximize soybean productivity in these soils, it is necessary to apply Phosphorus as part of a deliberate programme to improve soil fertility levels. It is recommended that farmers apply phosphorus fertilizer to their crop (soybean) so as to increase their harvest and income level affording them, the edge to go into large scale production of the crop. Nevertheless, for proper establishment, soybean seeds could be inoculated with appropriate Rhizobia.

Efforts should be made by extension officers and other stakeholders to increase farmers' awareness and appreciation of the technology. Because of an ever-increasing population in Africa and the world, with limited farm lands, it is very important to increase productivity of crops per unit area in order to feed the large population as well as increase protein source in diets in order to guarantee food security in Ghana particularly, and Africa at large.

REFERENCES

Ajeigbe HA and Woomer, PL (2010). Biological Nitrogen Fixation and Grain Legume Enterprise: Guidelines for N₂ Africa Lead Farmers. Tropical soil Biology and Fertility Institute of the International Centre for Tropical Agriculture. Nairobi / International Institute of Tropical Agriculture, Ibadan Nigeria. 21 pp.

Cadish, G., Bradley, RS and Nosberger, J (1992). Variation in Growth and N₂ – fixation within and among *Centrosema spp.* in Response to Phosphorus Supply. Tropical Grasslands, **26**: 226 – 234.

Cassman, KG., Whitney, AS and Fox, RL (1981). Phosphorus Requirements of Soybean and Cowpea as affected by Mode of N Nutrition. Agron. J. **73**: 17-22.

358 Fournier J, Timmers ACJ and Sieberer BJ (2008). Mechanism of Infection Threads
 359 Elongation in Root Hairs of *Medicago truncatula* and Dynamic Interplay with Associated
 360 Rhizobial Colonization. *Plant Physiology* **148**:1985-1995.
 361 Hardarson, G., Golbs, M and Danso, SKA (1987). Nitrogen Fixation in Soybean (*Glycine*
 362 *max* L. Merrill) as Affected by Nodulation Patterns. *Soil Biology and Biochemistry*,
 363 **21**:783-787.
 364 Millennium Development Authority (MiDA), (2013). Investment Opportunity in Ghana:
 365 Maize, Rice, and Soybean. Accra, Ghana.
 366 Ministry of Food and Agriculture of Ghana (MoFA), (2010). 2010 MoFA annual program
 367 Review. MoFA, Accra, Ghana.
 368 Ohyama Takuji, Ritsuko Minagawa, Shinji Ishikawa, Misaki Yamamoto, Nguyen Van Phi
 369 Hung, NO, Kuni, S., Takashi, S., Yoshifumi, N and Yoshihiko, T (2013). Soybean Seed
 370 Production and Nutrition. A comprehensive Survey of International Soybean Research –
 371 Genetics, Physiology, Agronomy and Nitrogen Relationships. Pp 115-121.
 372 Okagun JA, Otuyemi BT and Sangiga N (2004). Soybean Yield Determinants and
 373 Response to Rhizobia Inoculation. In: an On-Farm Trial in Northern Guinea Savannah of
 374 Nigeria. *West African Journal of Applied Ecology*, Vol. **6** Pp 30-34
 375 Othman, WN, Lie, TA, Mannetje, L and Wassink, GY (1991). Low Levels of Phosphorus
 376 Supply Affecting Nodulation, N₂ – fixation and Growth of Cowpea (*Vigna unguiculata* L.
 377 Walp). *Plant and Soil*, **135**:67-64.
 378 Singleton PW, Abaidoo, RC and Keyser, HH (1990). Highlights of Nitrogen Fixation
 379 Research. New York: Kluwer Academic publishers, 1990. Applied BNF Technology. Pp.
 380 44
 381 Tweneboah, CK (2000). Legumes. In *Morden Agriculture in the Tropics with Special*
 382 *Reference to Ghana*, Co-Wood Publishers, pp.185-3237.
 383 Van Gerpen, JH and Krah, J (2005). *The Biological Handbook ACS Chemical Biology*.
 384 Pp. 89
 385 Weaver, RW and Frederick, LR (1974). Effect of Inoculums Rates on Competitive
 386 Nodulation of *Glycine max* L. Merrill. II. Field Studies. *Agron. J.* **66**: 229-232.
 387 Woome, PL (2010). Biological Nitrogen Fixation and Grain Legume Enterprise:
 388 Guidelines for N₂ Africa Lead Farmers. Tropical Soil Biology and Fertility Institute of the

389 International Centre for Tropical Agriculture. Nairobi / International Institute of Tropical
390 Agriculture, Ibadan Nigeria. 21 pp.

391

392

393

394

395

396

397

398

399

400