

Upland Rice and Dry Season Sorghum Production under Double Cropping Systems in a Tropical Semi-Arid Region

J.P. Mvondo-Awono^{1*}, A. Boukong² and H. Beyegue-Djonko¹

¹Department of Crop Sciences, Faculty of Agronomy and Agricultural Sciences, University of Dschang. P.O. Box 222 Dschang, Cameroon.

²Department of Soil Sciences, Faculty of Agronomy and Agricultural Sciences, University of Dschang. P.O. Box 222 Dschang, Cameroon.

* Corresponding author; E-mail address: jpmvondoa@yahoo.fr

jpmvondoa@yahoo.fr

ABSTRACT

Aims: To investigate the influence of upland rice cultivar and dry season sorghum planting date on crop yield under rainfed double cropping systems as an option to increase cereal production for food security and income generation to farmers.

Study design: The experimental design was a randomized complete block design with 20 replicates for rice cultivars evaluation the first year (2009). Part of the experimental set up was subsequently used the following year (2010) to evaluate dry season sorghum production as second crop in the system. Treatments were the nine combinations of previous rice cultivars and sorghum transplanting dates replicated four times.

Place and Duration of Study: Experiments were conducted in 2009 and 2010 in a tropical semi-arid area, at Salak in the outskirts of Maroua, the main city of the Far-Northern Region of Cameroon.

Methodology: The three rice cultivars of Brazilian origin tested in this experiment were B22, Primavera and SEBOTA 281.2. They were followed by the local sorghum variety called "Guelendeng", sown at three different dates at the onset of dry seasons. Factors evaluated were the paddy yields of rice cultivars and grain yield of dry season sorghum.

Results: The rice cultivars were able to grow and score paddy yields ranging between 2.6 and 5.2 t ha⁻¹, with a highly significant interaction between years and cultivars (P<0.001). Upland rice grown in the rainy season was followed by the production of 832 to 1819 kg ha⁻¹ of dry season grain sorghum depending on year or preceding rice cultivar. Planting date did not reveal a significant effect on dry season sorghum yield but the latter was significantly determined by preceding rice cultivar (P<0.001). Dry season sorghum yield was significantly higher, when the preceding rice cultivar was Primavera the rainiest year (2010), or significantly lower the driest year (2009).

Conclusion: The choice of an upland rice cultivar to be included in double cropping systems with dry season sorghum is important but questions remain concerning the sustainability of the system in relation to moisture and nutrient management. Designing adequate fertilization schemes for cereals in double cropping systems and the availability of relevant short cycle

upland rice cultivars to be included as dry season sorghum preceding crop are main challenges to be addressed.

Keywords: Cropping systems, double cropping, upland rainfed rice, dry season sorghum, transplanted sorghum, Vertisols, food security.

1. INTRODUCTION

Efforts to increase national cereal production in Cameroon have led to the investigation on double cropping systems of rice and dry season sorghum (*Sorghum bicolor* (Linn.) Moench), on vertisols in the Far-Northern Region of the country [1, 2]. Sorghum, rice and other cereals are staple foods for local people but their supply is inadequate due to low yields or frequent crop failure [3, 4]. Also, local people cannot afford purchasing cereals from the market due to poverty and limited income. Consequently, famine and malnutrition often plague in the region [5]. Dry season sorghum locally called "Muskwari", is a reliable back-up against the failure of rainfed sorghum [6, 7]. About 46% of households cultivate dry season sorghum as sole crop on farms averaging 1.4 ha [8].

Dry season sorghum is transplanted on heavy-clayed vertisols at the beginning of the dry season and is capable to uptake residual water from non tilled soils to complete its cycle [2, 9]. The potential of vertisols for rice production is high and they are regarded as soils where rice can be introduced, without using the land reserved for other food crops [1, 4]. Vertisols characteristically swell, become sticky and impermeable when wet. As a result, large land tracks of vertisols in the Far-Northern Region of Cameroon are left flooded and abandoned during rainy season.

Introducing rice in the rainy season on wet or inundated vertisols before planting dry season sorghum may be a good strategy to increase domestic rice production and reduce rice imports. The latter have been on the rise in recent years and they exceeded 500 000 [10]. About 95% of the population eats rice at least once a week in Cameroon [3], but little has been undertaken to improve domestic production. National rice production was estimated at 65 000 tons over a cultivated area ca. 50 000 ha [10]. Rice is mainly produced under irrigation (87%) and rainfed lowland (9%); the rest (4%) is cultivated under upland conditions where the sole water supply is rainwater [10]. Farmers are becoming more and more interested in upland rice production due to the availability of improved early maturing cultivars. Some 30 cultivars of that type have been supplied for trial in various ecosystems in the Far-Northern Region of Cameroon between 2000 and 2004 [11, 12].

The period within which dry season sorghum can be successfully transplanted at the onset of the dry season is short and is determined empirically by farmers based on the pattern of late rains [4, 13]. This practice is well established in Northern Cameroon since the 19th century [14] and has assisted greatly to food security [4, 9]. However, dry season sorghum as second crop may be affected by preceding rice variety or by relative late plantings on drying vertisols.

Vertisols cover more than 800 000 ha in the Far-Northern Region of Cameroon alone [15] and more than 50% are suitable for upland rice production [13]. However, properties of vertisols vary depending on their location on the landscape. In lowland position, they are inundated for periods varying from days to weeks. In upland position, inundations barely exceed a few hours or days. Double cropping of rice and dry season sorghum was examined under rainfed lowland conditions [1, 2]. Such investigations are needed in other ecosystems including upland situations. It is useful to investigate farming systems that could benefit farmers both for food security and income generation. A system that would

jeopardize dry season sorghum systems may not be accepted by local farmers [7]. This paper examines the possibility of growing rice and sorghum in double culture cropping systems on upland vertisols in a semi-arid tropical region of Cameroon.

2. MATERIAL AND METHODS

2.1 Experimental site

Experiments were conducted in 2009 and 2010 in a tropical semi-arid area named Salak in the outskirts of Maroua, the main city of the Far-Northern Region of Cameroon. The site is a flat upland traditionally used for dry season sorghum production exclusively. Monthly precipitations records from the local meteorological station, and soil characteristics determined by routine laboratory methods [17] are indicated in Table 1 and Table 2, respectively.

Table 1. Monthly precipitations (in mm) over the experimental site at Salak in 2009 and 2010.

Year	J	F	M	A	M	J	J	A	S	O	N	D	Total
2009	0	0	0	24	35	57	186	199	96	0	0	0	597
2010	0	0	0	14	61	93	227	248	94	88	0	0	825

Source: [16].

Table 2. Properties of the topsoil (0-20 cm) of the experimental site as determined by routine laboratory methods

Particulars	Value	Methods
Sand (%)	22	Pipette Method
Silt (%)	48	Pipette Method
Clay (%)	30	Pipette Method
pH (1 : 2.5:: Soil : Water)	6.24	Glass Electrode, pH Meter
Organic carbon (%)	1.78	Walkley and Black
Total N (g kg ⁻¹)	1.76	Modified Kjeldahl Method
CEC (meq 100g ⁻¹ of soil)	4.54	Ammonium acetate at pH 7
Available Phosphorus (mg kg ⁻¹)	53	Bray II

2.2 Crop varieties

The three rice cultivars of Brazilian origin tested in this experiment were B22, Primavera and SEBOTA 281.2 whose summary descriptions are given in Table 3 [11]. They were selected based on their adaptation to rainfed conditions [16]. They need less than 120 days to be harvested. The sorghum variety under investigation is "Guelendeng". The main distinctive feature of this local variety is its short cycle and suitability for late plantings.

Table 3. Characteristics of the three rice cultivars of Brazilian origin (SEBORA 281.2, B22 and Primavera) used in the experiment

Rice cultivars	Plant Characteristics	
	Aptitude	Length of growing cycle (days)
SEBOTA 281.2	Upland, lowland or irrigated	115-120
B22	Upland	105-110
Primavera	Upland	105-110

Source: [11].

101 2.3 Treatments and experimental design

102 Treatments were the nine combinations of the three rice cultivars and the three dry season
103 sorghum transplanting dates replicated four times. The experimental design was a
104 randomized complete block design with 20 replicates for rice cultivars evaluation the first
105 year (2009). Part of the experimental setup was subsequently used the subsequent year
106 (2010) to evaluate dry season sorghum production as second crop in the system.
107 Experimental units were 5 m by 8 m plot of land. Spacing was 1.5 m between blocs and 1 m
108 between adjacent experimental units. Factors evaluated were the paddy yields of the three
109 rice cultivars and grain yield of the dry season sorghum subsequently transplanted at three
110 different dates.

111 2.4 Cropping practices

112 A non-selective herbicide (Glyphosate) was applied over the experimental units before
113 planting. Rice was seeded on July 2nd in 2009 and on June 29th in 2010 at 20 cm by 20 cm.
114 After each band of four rice lines, a space of 40 cm wide was provided for relay planting of
115 sorghum later in the season.

116 Rice seeds were directly hand planted on non tilled vertisols after their cracks were closed
117 following the first rains. Holes of 5 cm depth were opened using a sharp stick and 5 rice
118 seeds were dropped into planting holes. Then, 15 days after emergence, 150 kg ha⁻¹ of urea
119 were applied on rice experimental units. This was followed 30 days later by the application
120 of 200 kg ha⁻¹ of NPK fertilizer 22-10-15 as locally recommended. Rice plots were hand
121 weeded twice during the growing season. After rice harvest, sorghum was transplanted
122 at 1m by 1m in experimental units. In 2009, sorghum nurseries were planted on August
123 24th and 30th. In 2010, the planting dates of nurseries were August 25th and September
124 1st. Sorghum was transplanted on November 6th, 12th and 18th in 2009 and September
125 30th, October 10th and October 20th in 2010. No fertilizer was applied to dry season
126 sorghum and its weeding was not necessary.

127 2.5 Data collection and analysis

128 Rice yields were estimated from 1 m² subplots at the centre of experimental units
129 while sorghum yields were obtained from the two central lines. Yields from experimental
130 units were translated to amounts per hectare and analyzed. The grain humidity of rice
131 varieties was near 8%. Data were introduced on Excel sheets, and then transferred to
132 Statistical Analysis System (SAS) software analysis. Procedures GLM and MEANS of SAS
133 aided to perform the analysis of variance and the means' comparison. The probability level
134 was 5% for significant differences and 1% for highly significant differences.
135

136 3. RESULTS AND DISCUSSION

137 3.1 Paddy rice production

138 Rice was the first crop in the system and all three cultivars under study were able to grow
139 and provide a yield on upland vertisols. Paddy rice yields were between 2.6 and 5.2 t ha⁻¹
140 (Table 4). The analysis of variance indicates a highly significant interaction between years
141 and cultivars for rice production (Table 5). The rice cultivar SEBOTA 281.2 scored higher
142 yield of paddy than Primavera and B22 in overall. In 2009 the driest year, Primavera
143 produced more paddy than B22 but in 2010 under wetter conditions, the opposite trend was
144 observed (Table 4).
145
146

Table 4. Mean rice paddy yields in kg ha⁻¹ of the three rice cultivars SEBOTA 281.2, Primavera and B22 obtained in 2009 and 2010

Years (Y)	Rice cultivars (R)	Rice paddy yield (kg ha ⁻¹)	
		Mean	Standard Error
2009	SEBOTA 281.2	4201	163
	Primavera	3776	188
	B22	2653	108
2010	SEBOTA 281.2	5204	131
	Primavera	3701	125
	B22	3921	109

Table 5. Summary of analysis of variance of data from paddy production of three rice cultivars B22, Primavera and SEBOTA 281-2 obtained in 2009 and 2010 on upland Vertisols at Salak in the Far North Region of Cameroon

Sources of variation	DF	Mean Square	F value	Pr > F
Bloc	19	3032329	2.81	0.0001**
Year	1	48246854	44.65	<.0001**
Rice cultivar	2	62720413	58.04	<.0001**
Year*Rice cultivar	2	15200062	14.07	<.0001**

****:** Significant at the 1% level.

Yields obtained (Table 4) are in the range of the 2 to 6 t ha⁻¹ harvested by local farmers when they grow rice alone under lowland conditions [4]. However, yields observed in this experiment were low as compared to those obtained with the same varieties under more favorable conditions. SEBOTA 281.2 produced more than 10 t ha⁻¹ of paddy rice in the Lake Alaotra area in Madagascar where rainfall is 1000 to 1200 mm over 06 months [11]. In general, yields were better in 2010 compared to 2009 due to improved moisture conditions. On the experimental site, rainfall was 597 mm and 825 mm in 2009 and 2010, respectively. It is noteworthy to observe that prevailing climatic conditions were not conducive to achieving the full potential of tested varieties (Table 3).

Although all three rice varieties are considered suitable for upland rainfed conditions, it appears that when moisture conditions are better, B22 produces more paddy than Primavera and yields of SEBOTA 281.2 increase at a greater magnitude. SEBOTA 281.2 cultivar with a longer cycle (115 to 120 days) can be grown under rainfed or irrigated conditions [11]. In a similar experiment on lowland vertisols in another locality and in combination with other varieties, SEBOTA 281.2 produced up to 7.6 t ha⁻¹ of paddy rice [1]. Primavera is a strict upland rice cultivar which is very sensitive to water-saturated soils. This is probably why it yielded less than B22 under wetter conditions in 2010. They roughly have the same cycle (105 to 110 days). The cultivar B22 is also upland but more tolerant to soil water saturation than Primavera. B22 took advantage of the late rains that occurred in October 2010 (88 mm). The behavior of the two rice cultivars differs depending on growing conditions. This suggests that the amount of rainfall and availability of soil moisture are determinant factors for the choice of upland rice varieties, especially in areas with relatively short growing seasons as it is the case in the semi-arid area of the Far-Northern Region of Cameroon.

The shrinkage of the rainy season in the semi-arid area of the Far-Northern Region of Cameroon was documented from compiled records of the beginnings and ends of

successive rainy seasons since 1940 [18]. There is a need for crops and cultivars with short growing cycles. These can be found among the SEBOTA [11] and NERICA [12] rice cultivars. The cultivar SEBOTA 281.2 performed well in both years. The choice of rice cultivar has a significant impact on both individual crop yield and the system yield, when overall grain production is considered. Moreover, cultural practices that prolong rice cycle should be avoided. Application of nitrogen in excessive dosage often delay rice maturity [19] and would delay the planting of dry season sorghum. However, low rates of fertilizer application would jeopardize the sustainability of the double cropping system. Both rice and dry season sorghum need large amounts of nitrogen depending on crop yield [20].

3.2 Dry season sorghum production after rice

Dry season sorghum was the second crop in the double cropping system. Average dry season sorghum yields varied from 832 to 1819 kg ha⁻¹ depending on year and preceding rice cultivar (Table 6). The analysis of variance revealed highly significant effects of year and preceding rice cultivar on the production of dry season sorghum (Table 7). The average yield of sorghum was 1537 kg ha⁻¹ in 2009 and 920 kg ha⁻¹ in 2010. With regard to the preceding rice cultivar, dry season sorghum yield was higher (1094 kg ha⁻¹) when the preceding rice cultivar was Primavera the rainiest year (2010) and lower (1145 kg ha⁻¹) with B22 as preceding rice cultivar in 2009, the driest year (Table 6).

Table 6: Mean dry season sorghum yield of the local variety "Gueleldeng" grown after rice cultivars in 2009 and 2010 on Vertisols on the experimental site at Salak in the semi-arid area of the Far North Region of Cameroon.

Year (Y)	Preceding rice cultivar (R)	Dry season sorghum yield (kg ha ⁻¹)	
		Mean	Standard Error
2009	SEBOTA 281.2	1819	248
	B22	1145	98
	Primavera	1646	175
2010	SEBOTA 281.2	834	73
	B22	832	58
	Primavera	1094	81

208

209

210

Table 7 : Summary of the analysis of variance of yield data of the local dry season sorghum cultivar "Gueleldeng" after three rice cultivars B22, Primavera and SEBOTA 281.2 in 2009 and 2010 on upland Vertisols at Salak in the Far North Region of Cameroon

214

215

Sources of variation	Df	Mean Square	F value	Pr > F
Replicate	3	431685	2.89	0.04*
Year	1	2386384	15.99	0.0002**
Preceding rice cultivar	2	1081674	7.25	0.0017**
Planting date	2	291782	1.95	0.15
Year * Preceding rice cultivar	2	66773	0.45	0.64
Year * Planting date	2	78201	0.52	0.59
Preceding rice cultivar * Planting date	4	265158	1.78	0.14
Year * Preceding rice cultivar * Planting date	4	52672	0.35	0.84

216

217

*: Significant at the 5% level

**: Significant at the 1% level

218
219 Producing rice in the rainy season did not prevent cultivation of subsequent dry season
220 sorghum on upland vertisols. After paddy rice harvest, it was possible to grow a sorghum
221 crop on upland vertisols successfully. Subsequent dry season sorghum grain yields ranged
222 between 832 and 1819 kg ha⁻¹ within the two years of observation. Dry season sorghum
223 yields obtained under upland conditions are more important compared to the 540 to 811 kg
224 ha⁻¹ harvested after rice under lowland conditions [1]. In general, sorghum production as
225 second crop was better on the upland site compared to the lowland location where plantings
226 had to be delayed due to excess soil moisture and late rice harvest. On the upland site, early
227 sowing was achieved by relay plantings of dry season sorghum. Farmers' dry season
228 sorghum yields in sole cropping systems vary from 0 to more than 3 t ha⁻¹ in Northern
229 Cameroon, depending on soil and climatic conditions. Reported mean yields of dry season
230 sorghum grown as sole crop under experimental conditions varied from 500 to 1500 kg ha⁻¹
231 [13].

232 **3.3 Influence of rice-sorghum double cropping**

233 Sustained production of rice and sorghum on available vertisols would increase total cereal
234 availability. Hence, double cropping of rice and sorghum is interesting for local farmers
235 because the system maintains the production of the latter which is the traditional staple food
236 [4]. This scheme can also provide rice for households' consumption or commercialization [1,
237 7]. Rainfed sorghum is dominant but growing dry season or transplanted sorghum on vertic
238 soils is an opportunity to strengthen food security. Some farmers succeed in generating a
239 surplus. Income-based strategies increasingly involve the sale of part of the grain harvest.
240 Dry season sorghum is overall the most appreciated cereal and is sold in larger proportions
241 at a higher price, especially when it is supplied to towns [5].

242 Another interesting feature of the rice-sorghum double cropping system is that grain
243 production is increased without need for additional land. This form of agricultural
244 diversification or intensification, although in an embryonic state in the Far-Northern Region of
245 Cameroon, deserves a particular attention because of yield fluctuations and risks of crop
246 failure. In 2009 rice yields were relatively lower due to reduced rainfall. The following year
247 with improved rainfall, there was a substantial paddy rice yield increase. The risk of failure
248 also concerns dry season sorghum, the second crop in the system, and farmers are
249 reluctant to miss its transplanting period [7]. Since the preference of local farmers is on dry
250 season sorghum, the few who produce rice minimize the risk by using less than 30% of
251 cultivated vertisols for rice production [4].

252 Dry season sorghum production is mainly determined by soil water content at transplanting.
253 Because the dry season comprises most of its growing cycle, one may suggest the adoption
254 of early transplanting or the use of cultivars adapted to late plantings. Under upland
255 conditions and with plantings taking place before the 20th of November, no significant
256 difference among the dry season sorghum sowing dates was observed. Under lowland
257 conditions however, differences were observed among the sowing dates [1]. In 2010, higher
258 yields were obtained with the two early sowings while best performances were recorded with
259 late sowings in 2009 under lowland conditions. These differences could be attributed to
260 fluctuations in soil moisture throughout sites and years [6], but soil water content was not
261 monitored.

262 The time for transplanting dry season sorghum is determined empirically by growers after
263 the last rains, when the soils are still moist to ensure seedling regrowth, but are still dry
264 enough to avoid seedling decay [7]. Abundant water from rains is not favorable to dry
265 season sorghum establishment and at the same time, low soil moisture with late plantings is

not desirable as seedlings and plants thrive to complete the growing cycle. Monitoring soil moisture to determine the best transplanting period was suggested but there is high uncertainty on the occurrence of the late rains [13]. One of the solutions adopted by local farmers is the use of successive plantings in nurseries [7, 21]. Another option is the use of varieties adapted to late plantings such as "Guelendeng" which may have contributed to mask the effects of planting date in this experiment. The recourse to relay plantings of dry season sorghum before rice harvest is also a potential explanatory factor to the non-significance of the planting date effect.

The success of dry season sorghum requires good co-ordination of the nursery plant production period and the replanting date. Farmers perform staggered sowing in nurseries every five to 10 days from August onwards to obtain seedlings suitable for transplantation throughout September and October [21]. In general, rice produced more in the wetter year and dry season sorghum in the driest year. Although differences observed in yields are explained using the contrast in rains between 2009 and 2010 or the influence of preceding rice cultivar, other factors are also important for the phenology of dry season sorghum including the response to photoperiod or to low temperatures [9].

In extensive dry sorghum production systems, the crop is not fertilized. Nutrients are mainly provided by natural processes such as sedimentation, nitrogen fixation, and atmospheric deposition. Researches on agricultural practices have demonstrated that fertilization at planting or transplanting stage has no significant effect on dry season sorghum yield in Northern Cameroon [21, 22]. This was explained by low soil moisture during the dry season [22]. Dry season sorghum could better benefit of residual fertilizer applied to rice as first crop in the double cropping system. The annual application of 150 kg ha⁻¹ of urea and 200 kg ha⁻¹ NPK 22-10-15 (for a total of 113 kg N ha⁻¹) as presently recommended for rice production would not sustain both rice and dry sorghum needs in a long term. In this double crop cereal system (rice-sorghum), nitrogen fertilizer applications need to be substantial to prevent a rundown of soil fertility. Similar observations were made under lowland conditions [1]. Further work should determine adequate fertilizer levels for the rice-sorghum double cropping systems that ensure a positive nutrient balance for a sustained production.

This study demonstrates that double cropping of upland rice and dry season sorghum is a sound option to increase cereal production in the semi-arid region in Northern Cameroon. Growing rice as a precedent crop to traditional dry season sorghum ensures three to four times more output of cereals from the cultivation of the same plots than when dry season sorghum alone is produced. The local short-cycle-dry-season sorghum variety is very suitable to the double cropping system investigated as it tolerates late plantings and thus buffers efficiently counter effects that may result from delayed planting of sorghum to allow rice mature prior to its harvest. Since preceding rice cultivar revealed a significant effect, it is worthy dedicating efforts in choosing rice cultivars that suit better to the double cropping system. Versatile rice varieties that can be grown under various regimes (upland, lowland or irrigated conditions) are of great interest owing to their ability to cope with the variability of climatic conditions in the semi-arid region in Northern Cameroon.

4. CONCLUSION

Investigation of the possibility of producing rice and sorghum following a double cropping system on uplands vertisols in the Far-Northern Region of Cameroon has shown that the three rice cultivars SEBOTA 281.2, B22 and PRIMAVERA are suited for upland rainfed production on vertisols. As first crops in the system, they produced 2.6 to 5.2 t ha⁻¹ of paddy. In both years cultivar SEBOTA 281.2 performed better than the other cultivars. The choice of rice cultivar has a significant impact on both individual crop yield and the system yield.

317 Subsequent dry season sorghum grain yields ranged between 832 and 1819 kg ha⁻¹. Hence,
318 producing rice in the rainy season did not hinder subsequent dry season sorghum
319 production. The adoption of double cropping of rice and dry sorghum production on upland
320 vertisols can significantly increase the availability of cereals in the Far North Region of
321 Cameroon where cereal shortages, famine and low income are common among farmers.
322 However, the needs for adequate fertilization to sustain the cereal double cropping system
323 and the availability of short cycle rice cultivars are the main challenges.

324 **COMPETING INTERESTS**

325
326 Authors have declared that no competing interests exist.
327

328 **REFERENCES**

- 329
- 330 1. Mvondo-Awono, JP, Lawane, Boukong A., Beyegue-Djonko H, Abou Abba Abdoulaye,
331 Adjil Abadji and Tchikoua C. The influence of rice cultivar and sorghum planting date on
332 crop yield in lowland rainfed double cropping systems in northern Cameroon. *African*
333 *Journal of Agricultural Research*. 2013; 8(1): 57-63.
334
 - 335 2. Ambassa-Kiki R, Aboubakar Y, Boulama T. Zero-tillage for rice production on
336 Cameroonian Vertisols. *Soil and Tillage Research*. 1996; 39(1): 75-84.
337
 - 338 3. ACDIC (2006). Food sovereignty in Cameroon: a study based on the rice sector.
339 Association Camerounaise de Défense des Intérêts des Consommateurs, Yaoundé,
340 Cameroun. French.
341
 - 342 4. Gnassamo J, Kolyang. Mutations agricoles : adoption du riz pluvial dans les plaines
343 Tupuri, In Jamin JY, Seiny Boukar L, Floret C, editors. *Savanes africaines : des espaces*
344 *en mutation, des acteurs face à de nouveaux défis*, Actes du colloque, mai 2002,
345 Garoua, Cameroun. PRASAC, N'Djamena – CIRAD, Montpellier, France ; 2002. French.
346
 - 347 5. Madi A. Les prix des produits et le système productif dans la zone cotonnière de
348 l'Extrême Nord Cameroun. *Cahiers Agricultures*. 2000; 9: 125-130. French.
349
 - 350 6. Vernier P, Aboubakar Y, Tchari B. The upland rice-sorghum muskwari double cropping
351 system in Cameroon. Preliminary experiments. *Agronomie Tropicale*. 1987; 42(4): 280-
352 288.
353
 - 354 7. Abolgo P. Valorisation des Karé par la double culture Riz-Muskuwaari à Barlang dans la
355 Province de l'Extrême Nord Cameroun. Mémoire d'Ingénieur Agronome. Université de
356 Dschang, Dschang, Cameroon; 2005. French.
357
 - 358 8. Njomaha C. Agricultural change, food production and sustainability in the far North of
359 Cameroon. Phd Thesis, Leiden University, The Netherlands; 2004.
360
 - 361 9. Seignobos C, Moukouri HK. Potentialités des sols et terroirs agricoles. In *Atlas de la*
362 *Province de l'Extrême Nord Cameroun*. MINREST, INC-IRA, Paris; 2000.
363
 - 364 10. Piebiep Goufo. Rice Production in Cameroon: a Review. *Research Journal of Agriculture*
365 *and Biological Sciences*. 2008; 4(6): 745-756.
366
 - 367 11. Bouzinac S, Taillebois J, Seguy L. La Saga SEBOTA : les riz poly-aptitudes SEBOTAS
368 créés dans et pour les systèmes de culture durables en semis direct sur couverture

369 végétale (SCV) permanente au service des rizicultures alternatives permanentes
370 diversifiées, "propres" et à faible coût. CIRAD, France; 2009.
371
372 12. Nguyen VN. Recent developments in rice production in Sub-Saharan Africa and the
373 dissemination of NERICA. Paper presented on October 6th, 2002.at the FAO Japanese
374 Symposium on food security in Africa, Nihon University, Fujisawa, Japan; 2003.
375
376 13. Mathieu B. Le sorgho repiqué au Nord Cameroun: De l'analyse des pratiques culturelles
377 à l'accompagnement technique des producteurs. CIRAD-DPGT, France; 2000.
378
379 14. Couty P (1987). La production agricole en Afrique subsaharienne : manières de voir les
380 façons d'agir. Cahier des sciences humaines, ORSTOM, vol 23, N° (3-4) : 391-408.
381 French.
382
383 15. Brabant P, Gavaud M. Les sols et les ressources en terres du Nord Cameroun. Carte et
384 notice explicative N° 130. MESRES-IRA, Yaoundé. ORS TOM, Paris; 1985. French.
385
386 16. PCS ESA II. Rapports d'activités. Campagnes agricoles 2009 et 2010. SODECOTON,
387 Garoua, Cameroun; 2010. French.
388
389 17. Pauwels JM, Van Ranst E, Verloo M, Mvondo Ze AD. Méthodes d'analyses de sols
390 et de plantes, équipement, gestion de stocks de verrerie et de produits
391 chimiques. Publications Agricoles–28. MESIRES/CUDs/INADER Dschang, Cameroun.
392 AGCD Bruxelles, Belgique; 1992. French.
393
394 18. Beauvilain A. Tableau de la pluviométrie dans les bassins du Tchad et de la Bénoué, de
395 la création des stations à Décembre 1994. Documents pour la recherche N°3.
396 Ndjamena, Centre National d'Appui à la Recherche (CNAR), Tchad; 1995.
397
398 19. WARDA (Africa Rice Center)/FAO/SAA. NERICA®: the New Rice for Africa – a
399 Compendium. EA Somado, RG Guei and SO Keya (eds.). Cotonou, Benin: Africa Rice
400 Center (WARDA); Rome, Italy: FAO; Tokyo, Japan: Sasakawa Africa Association; 2008.
401
402 20. Sys CE, Van Ranst E, Debaveye J, Beernaert M. Land evaluation, part III: Crop
403 requirements. General Administration for Development Cooperation, Brussels, Belgium;
404 1993.
405
406 21. Barrault J, Eckebil JP, Vaille J. Point des travaux de l'IRAT sur les sorghos repiqués du
407 Nord- Cameroun. Agronomie Tropicale. 1972; 27 (8): 791-814. French.
408
409 22. Carsky RJ, Ndikawa R, Singh L, Rao MR. Response of dry season sorghum to
410 supplemental irrigation and fertilizer N and P on Vertisols in Northern Cameroon.
411 Agricultural water management. 1995; 28 (1): 1-8.
412
413