

**Induction of Resistance in Cocoyam
(*Xanthosoma sagittifolium*) to *Pythium*
myriotylum by Corm Treatments with
Benzothiadiazole and its Effect on Vegetative
Growth**

ABSTRACT

Root rot disease of cocoyam (*Xanthosoma sagittifolium*) caused by *Pythium myriotylum* (Oomycete) is a major constraint in cocoyam production in many countries in the world. Benzothiadiazole (BTH), a systemic acquired resistance inducer, was used in two concentrations (15 and 50 mg/L) as corm soaking treatment for 24 h duration to test for its effect on root rot disease resistance and vegetative growth of cocoyam under greenhouse conditions. The fungicide Metalim 72 WP (metalaxyl + cuprous oxide) was used as standard chemical control. Corm treatment with BTH significantly reduced cocoyam root rot disease (CRRD) severity compared with the pathogen-inoculated control. This was expressed by a disease index percentage of 37 and 20 for 15 and 50 mg/L BTH respectively after 28 days post-inoculation as opposed to the control with a disease index of 60. BTH at 50 mg/L was found to be more effective than the fungicide in inducing significant levels of resistance against *P. myriotylum*. In the absence of the pathogen, BTH and the fungicide applied as corm soaking treatment did not significantly influence the chlorophyll content. BTH significantly reduced shoot and corm/root dry weights compared to the control and fungicide respectively. However no visible phytotoxic effect was observed. BTH was tested for possible effects on *P. myriotylum* in vitro. There was no inhibition of mycelial growth with 15mg/L of BTH in the growth media in comparison with the control. No remarkable antifungal activity was observed with 50 mg/L. The promising effect of BTH against CRRD indicates that this compound may be future alternatives to traditional chemicals for disease control.

Keywords: Benzothiadiazole; induced resistance; corm treatment; *Xanthosoma sagittifolium*; *Pythium myriotylum*; vegetative growth

1. INTRODUCTION

Cocoyam (*Xanthosoma sagittifolium*), one of the most important edible tuber crops to millions of people in the tropics and subtropics, has been severely affected by a root rot disease caused by a soil-borne fungus, *Pythium myriotylum* [1, 2]. Symptoms of cocoyam root rot disease (CRRD) include: slow rate of leaf production, premature yellowing of young leaves and initial water soaked lesions at the root tips that gradually spread to the entire root system resulting in root rot. Plants become stunted when attacked at the early growing

stage. Yield losses in some infected cocoyam plantings in Cameroon were reported to be as high as 90% [1]. Acceptable cocoyam varieties resistant to CRRD are not available. Current control methods are based on cultural practices such as long interval of bush fallowing, roughing of diseased plants, crop rotation, etc and the use of fungicide which cannot consistently suppress the disease [3]. In the absence of cocoyam cultivars resistant to this disease and lack of efficient control measures, development of alternative or complementary approaches for management of this disease is highly desirable.

In this context, the induction of resistance using the host defence mechanisms is of practical application against plant diseases. This mode of resistance is increasingly attracting attention because of its potential usefulness in crop protection. The prospect of broad-spectrum disease control using the plant's own resistance mechanism has led to increasing interest in the development of agents which can mimic natural inducers of resistance [4]. Systemic acquired resistance (SAR) inducers have been shown to control many plant pathogens in agricultural production. BTH or acibenzolar-S-methyl (ASM) is among the most commonly used elicitors. When applied to the plant, it can induce accumulation of pathogenesis related-proteins (such as chitinases, glucanases, peroxidases) and lead to reduced incidence and severity of several diseases on many crops [5].

However, the scope of resistance activation by BTH is crop-specific. Optimal integration of BTH in disease control measures, therefore, needs to be carefully assessed for each crop. Very limited information is available about SAR inducers that may provide protection of cocoyam plants against CRRD. Previous studies indicated that BTH applied as foliar sprays was an efficient elicitor of some defence reactions in cocoyam and enhanced the activities of peroxidase, polyphenoloxidase as well as phenolic compounds [6]. BTH is usually applied as foliar sprays, but it can also induce systemic resistance when applied as seed imbibition in oilseed rape [7], cashew [8] and melon [9]. In cocoyam, the corms are the most common material used by farmers for cultivation. These corms are primary sources of *P. myriotylum* inocula. Consequently, inducer application by corm treatment suggests a suitable procedure for obtaining maximum protection against CRRD during seedling emergence and early growth stages. Therefore, the aim of this study was to test the effect of corm application of BTH on resistance to *P. myriotylum* in cocoyam and on plant vegetative growth. The direct effect of BTH on in vitro mycelial growth of *P. myriotylum* was also investigated.

2. MATERIAL AND METHODS

2.1 Experimental location and materials

The work was carried out in 2011-2012 in the greenhouse at the Jay P. Johnson Biotechnology Laboratory of the Institute of Agricultural Research for Development (IRAD), Ekona Regional Centre, in the South West Region of Cameroon.

Plant materials composed of corms of the highly susceptible cocoyam cultivar (*X. sagittifolium* cv. White) were obtained from farmers' fields in Ekona (South West Region, Cameroon). Corms were trimmed with a knife to an average weight of 71g, disinfected in 5% (v/v) sodium hypochlorite (commercial bleach, 9% chlorine) for 30 minutes and thoroughly rinsed.

The chemical inducers and fungicide used in this study included:

Bion (WG 50) with 50% BTH was dissolved in distilled water. Metalaxyl (1 metalaxyl: 5 cuprous oxide, w/w) is a product distributed by AGROCHEM Douala (Cameroon) and was prepared by dissolving the required quantity in distilled water. It is a systemic and contact

fungicide which is effective against *Pythium* spp. Concentrations of BTH used in this study are concentrations of the active ingredients of the product.

2.2 Effect of BTH as corm treatment on severity of CRRD

Disinfected corms were soaked into different concentrations of BTH (15 and 50 mg/L) for 24 h. The concentrations were based on the results from preliminary trials in the laboratory. Corms soaked for 12 h in the suspension of the 3.3 g/L of Metalm 72 WP were used as standard. Corms of healthy and disease control plants were soaked in distilled water for 24h.

Plant experiments were carried out in disease-conducive volcanic soil collected from Ekona. Soil was sterilized at 121°C for 20 minutes in an autoclave and cooled for 2 days prior to use. Chemically treated and control corms were planted in polybags (30.5 x 21.2 cm) containing 3 kg of sterilized soil. Each polybag contained 1 corm. The experimental set-up was a completely randomized design (CRD) with 5 corms per treatment, including a healthy and a disease control. Plants were grown in greenhouse at 28°C and watered when necessary to maintain moisture content of the soil substrate. The experiment was conducted three times.

Four-week-old plantlets, grown on sterilized soil, were used for challenge-inoculation with the pathogen. 5-7 day-old *Pythium* cultures were blended using a commercial blender in distilled water to produce a stock suspension of mycelia stands. The inoculum concentration was adjusted to about 1250 propagules per gram of soil. Inoculation consisted in spreading 100 ml of mycelia suspension on the soil of the potted plants.

Disease severity was determined the 28th day after inoculation on fully-opened leaves based on a rating scale from 0 to 4: 0 = 0%, 1 = 1 to 25%, 2 = 26 to 50%, 3 = 51 to 75%, and 4 = 76 to 100% of foliage with yellowing [1] and disease index percentage (DI%) was estimated using the formula as follows:

$$DI\% = ((n_1 + 2n_2 + 3n_3 + 4n_4) / 4N) \times 100$$

Where:

n₁ = number of plants with scale 1

n₂ = number of plants with scale 2

n₃ = number of plants with scale 3

n₄ = number of plants with scale 4

N = total number of plants sampled

2.3 Effect of corm treatment with BTH on plant vegetative growth

Disinfected corms were soaked into BTH or fungicide solution as described previously. For control plants, corms were soaked in distilled water for 24 h. The experimental set-up was a completely randomized design (CRD) with 10 corms per treatment, including a control. The experiment was repeated three times. Data was collected on the following parameters:

The daily numbers of sprouted corms were recorded until a constant count was achieved and the time taken for 50% sprouting of corms (E50) calculated according to the following formula [10]:

118 $E50 = t_i + (N/2 - n_i) \cdot ((n_j - n_i) / (t_j - t_i))$

119 Where:

120 N = final number of emerged seeds

121 n_i and n_j = cumulative number of seeds emerged by adjacent counts at times t_i and t_j

122 t_i and t_j = Initial time and final time, respectively

123 Chlorophyll content: Chlorophyll content in fresh leaves was estimated using the method
124 described in [11]. For chlorophyll extraction, 1g fresh tissue from interveinal leaf area was
125 cleaned several times with distilled water and homogenized in mortar and pestle with 90%
126 (v/v) acetone. The extract was then centrifuged at 2500 rpm for 10 minutes. The optical
127 density (OD) of the supernatant was read and recorded at 663nm and 645 nm
128 (corresponding to chlorophylls a and b respectively) using a spectrophotometer.

129 Chlorophyll content was calculated using the formula described by [12] as follows:

130 $\text{Chl a (mg/ml)} = 11.64 (A_{663}) - 2.16 (A_{645})$

131 $\text{Chl b (mg/ml)} = 20.97 (A_{645}) - 3.94 (A_{663})$

132 Where: A_{663} is the optical density (absorbance) at wavelength 663nm for chlorophyll a

133 A_{645} is the optical density at wavelength 645nm for chlorophyll b

134 Dry weight of shoots and corms/roots: At the end of the experiment, dry weight of shoots and
135 corms (including roots) were recorded. Samples were dried in an oven at 105°C for 72 h.
136 Possible phytotoxic effects of BTH were also evaluated.

137

138 **2.4 In vitro effect of BTH on mycelial growth of *P. myriotylum***

139 To investigate an eventual direct antagonism of BTH on *P. myriotylum*, the effect of BTH on
140 mycelial growth was assessed in vitro using PDA amended with two concentrations (15 mg/L
141 and 50 mg/L). Non-amended PDA was used as control. BTH was incorporated into the PDA
142 medium before autoclaving. One mycelial plug, 5 mm in diameter, was removed from
143 actively growing culture (5-7 day old cultures) of *P. myriotylum* and placed fungus-side down
144 in the centre of each 9 cm-PDA plate. The plates were incubated at 28°C (optimal growth
145 temperature of *P. myriotylum*) in an incubator. Each treatment had three replicates and the
146 experiment was repeated three times.

147 Radial growth was assessed by measuring the diameter of fungal colonies in two
148 perpendicular directions and calculating the mean diameter at 40 h after incubation which
149 corresponded to the time taken for the colony to grow up to the edge of the medium in the
150 control plates. The diameter of the plug (5 mm) was subtracted from the colony diameter
151 before estimating mean diameter of the colony.

152 The in vitro percentage inhibition was calculated according to the formula:

153 $\text{Percentage of inhibition} = ((C - T) / C) \times 100$

154 Where: C = colony diameter (mm) of the control

T = colony diameter (mm) of the test plate

2.5 Data Analysis

Data were subjected to tests of analysis of variance (ANOVA). After rejecting null hypothesis of equal means using ANOVA F-test, Fisher's Least Significant Difference (L.S.D.) was used for comparing treatment group means at 0.05. Statistical software Genstat computer package 5 Release 3.2 was used for Windows. Values were means $\pm$ standard deviation.

3. RESULTS AND DISCUSSION


3.1 Effect of BTH on CRRD severity

Chemical activation of disease resistance in plants could represent an additional option for crop protection. This may be a good option for controlling soil borne pathogens where chemical application is difficult. Resistance inducers may also provide a complementary option for chemical control against some pathogens where genetic resistance is not available or not sufficient. CRRD is a devastating disease and given the lack of effective control measures, corm treatment with BTH was evaluated as a preventive management strategy during early attack. BTH applied as corm treatment induced various levels of CRRD protection, as identified by disease index in plants inoculated with *P. myriotylum*. Disease symptoms were observed in all inoculated plants but greatly reduced in plants treated with BTH. The two tested levels of BTH were found to be effective in inducing statistically significant levels of resistance against *P. myriotylum*. This was expressed by a disease index percentage of 37 on the 28th day post- inoculation for the low concentration tested (15 mg/l). Disease index decreased to 20 with increase in concentration of BTH (Fig. 1). Plants from control corms sown in inoculated soil showed the highest disease index percentage. In comparison, seed soaking in solutions of BTH at 25 or 50 mg/L consistently reduced disease severity of black root rot caused by *Thielaviopsis basicola* in cotton seedlings against [13]. Applications of BTH have been shown to be effective in decreasing susceptibility in several plant species [14, 15, 16].

In earlier study, [6] showed that foliar application of BTH stimulated defence reactions in *X. sagittifolium*. The resistance was noted as a decrease in CRRD incidence and severity. However, BTH application as corm treatment, as described in this study, suggests an easy, rapid and cheaper procedure for obtaining maximum protection against CRRD during sprouting, emergence and the early establishment phase of the crop. Using a combination of these two modes of BTH application may offer a more effective approach in managing CRRD.

Although BTH did not completely suppress the CRRD, the reduction and delay of symptoms by this elicitor could be exploited by giving possibility to cocoyam growers to apply and combine other CRRD control methods such as the use of tolerant varieties and composts.

The fungicide also significantly reduced CRRD when applied as corm treatment with a disease index percentage of 36. But BTH at 50 mg/L was significantly more effective than the fungicide. BTH is considered an environmentally friendly crop protection agent and plant defence activator and has been tentatively employed to reduce the use of agrotoxic substances while maintaining crop yields [17]. The results presented here showed that exogenously applied BTH to corm provided protection to cocoyam plant similar or better to that of the standard fungicide. These observations shed more light on the potential of BTH as a valuable alternative means of CRRD control.



202

203 **Fig.1. Effect of corm treatment with BTH to control *P. myriotylum* in cocoyam**
 204 *Bars with common letter do not differ significantly according to Fisher's L.S.D.*

205 **3.2 Effect of BTH on cocoyam vegetative growth**

206 The effect of BTH on cocoyam growth is presented in the Table 1. Sprouting was observed
 207 in all the cocoyam corms soaked in different treatments. The chemicals tested affected only
 208 the time to sprouting. Sprouting time was significantly increased in corms treated with 15
 209 mg/L of BTH as compared to control ones. Corms treated with 50 mg/L of BTH or with the
 210 fungicide showed sprouting time of 8.34 and 8.63 days respectively. No evidence of
 211 phytotoxic symptoms in terms of disruption of plant growth, leaf stunting or scorching was
 212 observed within days following all treatment application.

213 There was no significant difference in the chlorophyll content of the leaves between means
 214 of treated and control plants. But shoot dry weight of plants treated with 50 mg/L BTH was
 215 significantly reduced when compared with fungicide treated plants. BTH significantly reduced
 216 the dry weight of corms/roots as compared to the control.

217 The deleterious effects were only observed as reduced growth but not foliar damage or
 218 reduced chlorophyll content; therefore it might be possible to overcome the reductions in dry
 219 mass via fertilization. The negative influence of BTH on plant growth is well documented and
 220 has been explained as a consequence of the allocation cost arising from the metabolic
 221 competition between the processes involved in plant growth and the de novo synthesis of
 222 defence-related compounds [18]. Thus it is likely that BTH controls root rot by activating the
 223 defence responses rather than by directly affecting root development and function. Such
 224 growth defects triggered by BTH may also depend on abiotic conditions [19] and host
 225 genetics, since some plants are much more sensitive to BTH-induced toxicity than others
 226 [20]. These findings of negative impacts of BTH applications on the vegetative growth of
 227 cocoyam plants are consistent with previous study on bean and cucumber in which foliar
 228 spray of BTH led to adverse effects on growth [21].

Table 1. Effect of BTH corm treatment on growth parameters in cocoyam

Treatment	Sprouting time (days)	Chlorophyll content (mg/ml)	Shoot dry weight (g)	Corm/root dry weight (g)
Control	7.06 b	30.39 a	5.72 ab	8.42 a
15mg/L BTH	9.14 a	30.35 a	5.82 ab	6.02 b
50mg/L BTH	8.34 ab	29.49 a	4.62 b	4.96 b
Fungicide	8.63 ab	30.35 a	7.26 a	7.51 ab
LSD	2	2	2	2

In each column, means with similar letter(s) are not significantly different from each other. Bars with common letter do not differ significantly according to Fisher's L.S.D.

3.3 Effect of BTH on in vitro mycelial growth of *P. myriotylum*

To address whether BTH has a direct action on *P. myriotylum*, possibly acting as a fungicidal compound, we checked if mycelial growth was affected. BTH did not reduce mycelial growth at 15mg/L after 40 h of culture. But the in vitro growth of *P. myriotylum* depended partially on the concentration of BTH tested, because by increasing the concentration from 15 to 50 mg/L, growth of the pathogen was inhibited by only 12% which is not significant. These results confirmed that BTH possessed little or no direct antimicrobial activity when used at lower concentrations. Its protective effect against pathogen was mainly due to the induction of the plant's own defence mechanisms [22].

Although BTH at low concentration (15 mg/L) did not exhibit any fungitoxicity against *P. myriotylum* in the form of reduced mycelia growth, when applied to the corm at the same concentration, BTH reduced CRRD severity on inoculated plants. The induction of the defence mechanisms without the antifungal activity was enough to suppress the infection in cocoyam plants treated with BTH and inoculated with *P. myriotylum*. These results showed that the protective effect of BTH against CRRD could only due to its host defence-inducing activity not to adverse effect on growth of the pathogen. This was also demonstrated in our study by temporal separation of the inducer and the pathogen inoculation. Induced systemic resistance normally requires a time-lapse period between inducer treatment and pathogen inoculation for gene activation of defense responses to take place [23].

4. CONCLUSION

Based on the present work, the use of BTH was efficient in protecting greenhouse-grown cocoyam plants against *P. myriotylum*, in a way similar to the standard fungicide. The protective effect of BTH cannot be attributed to the toxic effect of the chemical on the fungus, as the product did not show fungistatic activity against *P. myriotylum* in vitro. This work also showed that in the absence of pathogen, treated plants experienced reduced growth. Additionally, neither the BTH nor the fungicide treatments affected chlorophyll content in cocoyam. If confirmed in field tests, resistance induced by BTH has potential for control CRRD possibly in combination with existing crop protection practices.

CONSENT (WHERE EVER APPLICABLE)

No manuscripts will be peer-reviewed if a statement of patient consent is not presented during submission (wherever applicable).

This section is compulsory for medical journals. Other journals may require this section if found suitable. It should provide a statement to confirm that the patient has given their

informed consent for the case report to be published. Journal editorial office may ask the copies of the consent documentation at any time.

Authors may use a form from their own institution or SDI Patient Consent Form 1.0. It is preferable that authors should send this form along with the submission. But if already not sent during submission, we may request to see a copy at any stages of pre and post publication.

If the person described in the case report has died, then consent for publication must be collected from their next of kin. If the individual described in the case report is a minor, or unable to provide consent, then consent must be sought from their parents or legal guardians.

Authors may use the following wordings for this section: "All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.'"

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

This section is compulsory for medical journals. Other journals may require this section if found suitable. If human subjects are involved, informed consent, protection of privacy, and other human rights are further criteria against which the manuscript will be judged. It should provide a statement to confirm that the authors have obtained all necessary ethical approval from suitable Institutional or State or National or International Committee. This confirms either that this study is not against the public interest, or that the release of information is allowed by legislation.

All manuscripts which deal with animal subjects must be approved by an Institutional Review Board (IRB), Ethical Committee, or an Animal Utilization Study Committee. , and this statement, and approval number, must accompany the submission. If required, author should be ready to submit a scanned copy of the IRB or Ethical Committee Approval at any stage of publication (Pre of post publication stage). The manuscript should contain information about any post-operative care and pain management for the animals.

For manuscripts involving animal experiments, Authors may use the following wordings for this section "All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

All manuscripts which deal with the study of human subjects must be accompanied by Institutional Review Board (IRB) or Ethical Committee Approval, or the national or regional equivalent. The name of the Board or Committee giving approval and the study number assigned must accompany the submission. If required, author should be ready to submit a scanned copy of the IRB or Ethical Committee Approval at any stage of publication (Pre of post publication stage).

For manuscripts involving human experiments, Authors may use the following wordings for this section: "All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki."

327

328 REFERENCES

329

330 1. Tambong JT, Poppe J, Höfte M. Pathogenicity, electrophoretic characterization and in
331 planta detection of the cocoyam root rot disease pathogen, *Pythium myriotylum*. European J
332 Plant Pathol. 1999;105:597-607.

333 2. Perneel M, Tambong JT, Adiobo A, Floren C, Saborio F, Levesque A, Höfte M.
334 Intraspecific variability of *Pythium myriotylum* isolates from cocoyam and the host crops.
335 Mycol Res. 2006;110:583-593.

336 3. Adiobo A, Oumar O, Perneel M, Zok S, Höfte M. Variation of *Pythium*-induced cocoyam
337 root rots severity in response to soil type. Soil Biol Biochem. 2007;39:2915-2925.

338 4. Walters D, Walsh D, Newton A, Lyon G. Induced resistance for plant disease control:
339 maximizing the efficacy of resistance elicitors. Phytopathology. 2005;95:1368-1373.

340 5. Oostendorp M, Kunz W, Dietrich B, Staub T. Induced disease resistance in plant by
341 chemicals. Eur J Plant Pathol. 2001;107:19-28.

342 6. Mbouobda H D, Fotso, Djocgoue PF, Omokolo ND, El Hadrami I, Boudjeko T. Benzo
343 (1,2,3)-thiadiazole-7-carbothioic S-methyl ester (BTH) stimulates defense reactions in
344 *Xanthosoma sagittifolium*. Phytoparasitica. 2010;38:71-79.

345 7. Jensen BD, Latunde-Dada AO, Hudson D, Lucas JA. Protection of Brassica seedlings
346 against downy mildew and damping-off by seed treatment with CGA 245704, an activator of
347 systemic acquired resistance. Pest Sci. 1999;52:63-69.

348 8. Lopez AMQ, Lucas JA Lucas. Effects of plant defence activators on anthracnose disease
349 of cashew. Eur J Plant Pathol. 2002;108:409-420.

350 9. Buzi A, Chilozi G, De Sillo D, Magro P. Induction of resistance in melon to *Didymella*
351 *bryoniae* and *Sclerotinia sclerotium* by seed treatments with acibenzolar-S-methyl and
352 methyl jasmonate but not with salicylic acid. J Phytopathology. 2004;152:34-42.
353

354 10. Farooq M, Basra SA, Hafeez K, Ahmad N. Thermal hardening: a new seed vigor
355 enhancement tool in rice. J Integr Plant Biol. 2005;47:187-193.

356 11. Lichtenthaler K, Buschmann C. Chlorophylls and carotenoids: measurement and
357 characterization by UV-VIS spectroscopy. In: Wrolstad RE, Acree TE, An H, Decker EA,
358 Penner MH, Reid DS, Schwartz SJ, Shoemaker CF, Sporns P, editors. Current protocols in
359 food analytical chemistry (CPFA). New York: John Wiley & Sons; 2001.

360 12. Strain HH, Svec WA. Extraction, separation, estimation and isolation of chlorophylls. In:
361 Vernon LP, Seeley GR, editors. The chlorophylls. Academic; 1966.

362 13. Mondal AH, Nehl DB, Allen SJ. Acibenzolar-S-methyl induces systemic resistance in
363 cotton against black root rot caused by *Thielaviopsis basicola*. Australasian Plant Pathol.
364 2005;34(4):499-507.

- 365 14. Rohilla Rashmi, Singh US, Singh RL. Mode of action of acibenzolar-S-methyl against
366 sheath blight of rice, caused by *Rhizoctonia solani* Kühn. Pest Manag Sc. 2001;58:63-69.
- 367 15. Amzalek E, Cohen Y. Comparative efficacy of systemic acquired resistance-inducing
368 compounds against rust infection in sunflower plants. Phytopathology. 2007;97:179-186.
- 369 16. Faessel L, Nassr N, Lebeau T, Walter B. Effects of the plant inducer, acibenzolar-S-
370 methyl, on hypocotyl rot of soybean caused by *Rhizoctonia solani* AG-4. J Phytopathology.
371 2008;156:236-242.
- 372 17. Iriti M, Faoro F. Does benzothiadiazole-induced resistance increase fitness cost in bean?
373 J Plant Pathol. 2003;85:41-47.
- 374 18. Heil M, Hilpert A, Kaiser W, Linsenmair KE. Reduced growth and seed set following
375 chemical induction of pathogen defence: does systemic acquired resistance (SAR) incur
376 allocation costs? J Ecol. 2000;88:645-654.
- 377 19. Dietrich R, Ploss K, Heil M. Growth responses and fitness costs after induction of
378 pathogen resistance depend on environmental conditions. Plant Cell Environ. 2005;28:211-
379 222.
- 380 20. Walters DR, Bingham IJ. Influence of nutrition on disease development caused by fungal
381 pathogens: implications for plant disease control. Ann Appl Biol. 2007;151:307-324.
- 382 21. Azami-Sardooei Z, Seifi HS, De Vleeschauwer D, Höfte M. Benzothiadiazole (BTH)-
383 induced resistance against *Botrytis cinerea* is inversely correlated with vegetative and
384 generative growth in bean and cucumber, but not in tomato. Australasian Plant Pathol.
385 2013;42(4):485-490.
- 386 22. Vallad GE, Goodman RM. Systemic acquired resistance and induced systemic
387 resistance in conventional agriculture. Crop Sci. 2004;44:1920-1934.
- 388 23. Ryals JA, Neuenschwander UH, Willits MG, Molina A, Steiner HY, Hunt MD. Systemic
389 acquired resistance. Plant Cell. 1996;8:1809-1819.
390
391

APPENDIX