

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23

5
6
7
8
9
10

6

7

8

9

10

12

13
14
15

16
17

18
19

20
21
22
23

treatment plot for data collection on growth, yield and soil chemical composition. The data were subjected to analysis of variance F- test and the means separated by Duncan Multiple Range Test at 5%

Result. The results showed that there were significant ($P=0.05$) increases in plantain growth ,yield parameters and soil chemical composition under different treatments. The mixtures of neem and wood ash extracts increased significantly [$P= 0.05$], the leaf area and stem girth of plantain by 49.24% and 12% respectively compared to neem leaf extract. Also, the mixtures of the two extracts increased plantain bunch weight, finger weight, diameter, length and finger population by 39%, 41%, 3%, 36% and 41.6% compared to neem leaf extract. When compared with NPK 15-15-15, the treatment increased plantain bunch weight , fingers weight, diameter ,length and finger population by 29.7%, 31%, 6%, 8%, and 25% .Besides, the mixture of the water extracts of neem leaf and wood ash increased soil pH, organic matter , K, Ca, and Mg by 14%, 78%, 31%, 95.5% and 94% compared to NPK 15-15-15 fertilizer except N and P. The soil K/Ca, K/Mg and P/Mg ratios were higher under NPK 15-15-15 fertilizer than the liquid extracts signifying nutrient imbalance.

Conclusion; Mixture of water extracts of neem leaf and wood ash gave the best growth and yield parameters of plantain and this was due to its nutrients superiority compared to the other treatments.

KEY WORDS: Growth, improved soil chemical composition, low cost neem leaf, wood ash and 1:1 water extracts, Plantain (*Musa sapientum L*), and yield performances..

moyinjesu2004@yahoo.com

47 **GSM : 08038512581**

48

49

50

51 **INTRODUCTION**

52 Plantain (*Musa sapientum* L.) belongs to the family Musaceae and it originated
53 from Malaysia. It is a herbaceous plant with short lived stalks which are corm like
54 rhizomes usually 1.8m tall. Plantain is a food crop eaten by people in the West, East
55 and Central Africa, it has become exportable crop which is consumed in Europe,
56 America and Asia. Plantain has high iron content, minerals and vitamins which
57 provide dietary needs of people for better health living and it is processed into
58 different food stuff such as plantain flour, snacks and chips which are highly sought
59 for in both local and international markets [1]. Therefore, there is an enormous
60 increase in demand for the crop.

61 Total world production of the plantain and banana is estimated to be over 16
62 million metric tons out of which estimated 12 metric tons are produced in Africa
63 annually. About 70 million people in African region are estimated to derive more than
64 one quarter of their food energy requirements from plantain. [2]. In Nigeria, there are
65 some yield reduction constraints such as decline in soil fertility, pests and diseases and
66 weeds infestation that limit expansion [3].

67 Apart from medicinal value, neem leaf extract has many uses in plant
68 protection and production. Neem leaf extract can inhibit soil nematodes (4). Products
69 derived from neem leaf can act as powerful Insect Growth Regulation (IGR) which

reduce insect growth in crops and plants. Thus, neem products can be used as neem insecticide, neem pesticide, neem fumigant, neem fertilizer, neem manure, neem compost, neem urea coating agent and neem soil conditioner [5], [6].Neem seed extracts can reduce nitrification [7] of applied ammonium sulphate fertilizers and leaching loss of nitrogen. Pot experiments showed that neem extract could increase the use efficiency of chemical nitrogen significantly in soils [8].

In Bangladesh, wood ash is regularly used in banana plantations in terrace areas. Farmers say wood ash prevents diseases and improves water use efficiency. Wood ash is a good source of potash and lime; it also improves the pH of acidic soils in a manner similar to agricultural lime [9]. Wood ash had long been regarded as a source of potassium for plant growth (1-13% K by weight) but it also contains appreciable quantities of Ca (20-30%), P (0.3 - 0.4%), Mg (1- 3%) and smaller amounts of S, Mn, Fe, B, Cu and Zn [10].

Despite the economic importance of plantain to the nation, it is to note that maximum yields of the crop have not been attained. This is because of the continuous cultivation on the same piece of land without fertilizer application which has led to decline in soil fertility [11]. Besides, the present hectareage of plantain is very low because of subsistence farming and poor agronomic knowledge on the part of farmers.

Efforts aimed at improving the yield and increasing the soil fertility led to the uses of inorganic fertilizers but their uses are limited by high cost especially by poor farmers and the destruction of soil properties on continuous use [12].Further attempts to find substitute for the use of chemical fertilizers led to the use of solid organic fertilizers such as poultry, goat, pig and cattle manures to achieve sustainable

production but the problems of bulkiness and difficulty in the transportation of such materials affected their adoption by farmers [13].

In view of the above facts, water extracts of wood ash, neem leaf and mixed neem leaf and wood ash were applied at present to grow plantain. Except the works of [14], [15] and [16] on the use of neem leaf and wood ash extracts to control pests in maize and cowpeas and as source of fertilizers, there is scarcity of research information on the uses of water extracts of neem leaf and their mixtures on the growth and yield parameters of plantain.

The objectives of this study are to

- i. Determine the effects of water extracts of wood ash, neem leaf + wood ash on the growth and yield parameter of plantain.
- ii. To determine the effect of the extracts on the soil chemical composition after 16 months of planting plantain.

2.0 MATERIALS AND METHODS

2.1 Site description

A field experiment to evaluate the effects of neem leaf extract , wood ash extract and mixed extracts of neem leaf and wood ash on plantain in the rainforest zone of Akure, Nigeria was conducted during 2006-2009.

The soil is sandy clay loam, skeletal, kaolintic, isohyperthermic oxic paleustalf (Alfisol) [17] The annual rainfall is between 1100 and 1500mm while the temperature ranged between 29-32°C

2.2 Collection and analysis of the extracts

115 Wood ash and neem leaf were collected from the cassava processing unit and
116 neem leaf plantation in Federal College of Agriculture, Akure while the NPK 15-15-
117 15 fertilizer was purchased from Ondo State Agricultural Input Supply Agency. The
118 plantain suckers were collected from the Nigeria Institute for Agricultural Research,
119 Ibadan.

120 The preparation of neem leaf extract was done by weighing 10 kg of fresh
121 leaves, chopped into bits using a knife, immersed in a plastic container containing 50
122 litres of water, kept under a shade and properly covered. The solution was stirred
123 every three days to allow proper leaching of the nutrients into the water until the 14th
124 day. Thereafter, the leaf were carefully removed using a sieve of 2mm to obtain clean
125 neem leaf extract and diluted at a ratio of 1:1 to reduce the concentration of extracts as
126 well as preventing scorching of the plants. The application of the extracts is at 3L per
127 plot [833.3 Lha⁻¹]. wood ash extract was prepared by weighing 10kg of sieved wood
128 ash into 50 litres of water in a plastic container and thoroughly stirred with paddle
129 every three days to enhance proper leaching of nutrients. This continued until the 14th
130 day of setting up the experiment and the suspension was properly sieved to obtain
131 clear wood ash extracts.

132 The wood ash extract was also diluted at a ratio of 1:1 with water to reduce the
133 concentration and applied at 3L per plot. The remnants of wood ash solid components
134 were properly disposed to prevent contamination of the environment.

135 Mixed water extract of neem leaf and wood ash was prepared by taking 50% of
136 undiluted neem leaf extract and 50% of undiluted wood ash by volume and diluted at
137 a ratio of 1:1 with water and applied at 3L per plot while the NPK 15-15-15 fertilizer

138 was applied at 300 kg ha^{-1} . 5ml each of the filtered extracts of neem leaf, wood ash and
 139 mixed neem leaf + wood ash was analyzed for %N using microkjedahl method [18]
 140 while soluble P, K, Ca and Mg contents were determined [19].
 141
 142

143 **TABLE 1: The chemical analysis of the extracts used for the experiment**

Treatments	N (%)	P (%)	K (%)	Ca (%)	Mg (%)	Quantity applied per plot (36m²)	Quantity applied Per hectare
Neem leaf extract	3.56	0.83	1.67	0.77	0.75	3L	833.3L
Wood ash extract	0.15	0.53	2.60	15.00	1.00	3L	833.3L
Mixed neem leaf + wood ash extract	3.69	1.10	3.20	15.66	1.53	3L	833.3L

144
 145 Table 1 presents the data on chemical composition of the extracts used for the
 146 experiment. Mixed water extract of neem leaf and wood ash had the highest values of
 147 % N, P, K, Ca and Mg compared to wood ash and neem leaf extracts (sole forms). The
 148 sole form of neem leaf had better values of % N and P than wood ash extract while the
 149 wood ash extract had higher values of % K, Ca and Mg than the neem leaf.
 150
 151
 152

TABLE 2: Total amount of nutrients kg ha^{-1} by each extract and NPK 15-15-15 fertilizer

Treatments	N	P	K	Ca	Mg
	Total kg ha^{-1}				
Neem leaf extract	148.3	34.6	69.6	32.08	31.3
Wood ash extract	6.3	22.9	108.3	625.0	41.7
Mixed neem leaf + wood ash extract	153.7	45.8	133.3	625.5	63.8
NPK 15-15-15	45.0	45.0	45.0	0.2	0.1

Formatted: Highlight

The amount of nutrients [total kg/nutrient] supplied by each extract; mixed neem leaf + wood ash, neem leaf, wood ash and NPK 15-15-15 fertilizer is also presented in Table 2. More nutrients were supplied by the extracts particularly the mixed extract of neem leaf+ wood ash than the NPK 15-15-15 and control treatments.

The quantities of nutrients supplied by the extracts were adequate for sustaining growth and yield of plantain in the field as reflected in the soil chemical composition after harvest of plantain.

2.3 Field experiment

The land was cleared, ploughed and harrowed, divided into plots and each plot size is 6m X 6m (36m²). Hybrid plantain suckers treated with nemagon (a.i.1, 2 Dibromo-3-chloropropane) to control nematode were planted in early July, 2006 at a spacing of 2m X 2m.

There were five treatments namely: neem leaf extract, wood ash extract, mixed extract of neem leaf + wood ash applied at 3 litres/36m² plot, NPK 15-15-15 fertilizer applied at 300 kg/ha and a control treatment (no fertilizer nor extract), replicated three times and arranged in a randomized complete block design.

Weeding operation was first done after three weeks of planting and continued at three weeks interval until harvest. Five plants of sprouted plantain suckers were sampled in each plot on which parameters for plant height (cm), leaf area (cm²), stem girth (cm) and leaf population or number were taken every week starting from 2 weeks after establishment and this continued till 20 weeks after establishment.

The plantain started bearing inflorescence between 10 12 months after planting and the first harvest for matured plantain bunches was done at 15th month after establishment (November, 2007) and continued till early 2008. The following yield parameters were taken weight of plantain fingers, length of fingers (cm) plantain fingers diameter and plantain finger population. After harvesting, the plantain plots were properly weeded and the growth continued till the second harvest in late October, 2008 and April 2009.

Necessary agronomic practices such as propping up the plantain with sticks to prevent being fallen off during storms in early year, also, the plants were sprayed with Avesthrin (Cypermethrin 10EC at 10 ml/10 Litres of water) to prevent insect attack on the leaves.

2.4 Soil collection and analysis

Thirty core samples were collected randomly from 0-15cm depth on the site using soil auger, mixed thoroughly and the bulk sample was taken to the laboratory, air-dried and sieved to pass through a 2mm screen for chemical analysis.

The soil pH (1:1 soil:water) and (1:2 soil:0.01M CaCl_2) solution was determined by using a glass calomel electrode system[20] while organic matter was determined by the wet oxidation chromic acid digestion[21]. The total nitrogen was determined by the microkjedahl method [19] while available soil phosphorus (P) was extracted by the Bray P-1 extractant and measured by the Murphy blue colouration and determined on a spectronic 20 spectrophotometer at 882nm[22]. Exchangeable soil K, Ca and Na were extracted with 1M NH_4OAc [Ammonium acetate] pH 7 solution. The K, Ca and Na contents were determined with flame photometer while Mg was determined with an atomic absorption spectrophotometer [AAS] [23].

The soil exchangeable acidity [H^+ and Al^{3+}] were determined using 0.01M HCL extractant and titrated with 0.01M NaOH[24] while the micronutrients (Mn, Cu, Fe and Zn) were extracted with 0.1M HCl [25]. The mechanical analysis of the soil % silt, % clay and % sand textural determination was done by the hydrometer method [26]. The soil bulk density was determined by core method as described by [27].

214 After the experiment, soil samples were taken from each plot, air dried, sieved
215 with 2mm sieve, and analyzed for soil pH, organic matter, total N, available P,
216 exchangeable K, Ca and Mg as described ear The soil properties before application
217 of the treatments (initial soil fertility) were presented in Table 3. According to the
218 established critical levels for soil in South West Nigeria, the soil is acidic pH 5.80 and
219 low in soil organic matter compared to critical levels of 3%
220 [28]

221 . The soil nitrogen (%N) is 0.07 which was less than 0.15% total nitrogen considered
222 as optimum for crops [29]. The soil available P is 5.10 mgkg¹ considered as adequate
223 for crop production [28].

224 Exchangeable K, Ca, Mg and Na values were much lower than the critical
225 level of 0.2 mgkg⁻¹ while the micronutrients were more than the established critical
226 levels of 5 mgkg⁻¹ Fe, 1.0 mg/kg⁻¹ Cu, 5.0 mgkg⁻¹ Mn and 3.0 mgkg⁻¹ Zn [30]. The
227 soil bulk density was 1.60 mgm⁻³ while the soil textual class is sandy loam.

228

229 **TABLE 3: Soil properties before planting plantain**

Particulars	Values	Methods
Soil pH (H ₂ O)	5.8	Glass calomel electrode system (1)
Soil pH (0.01M CaCl ₂)	5.3	
Organic matter (%)	0.4	Wet oxidation chromic acid digestion (2)
Nitrogen (%)	0.1	Microkjedahl method (3)
Available P (mg/kg)	5.3	Bray P extractant (4)
Exchangeable Bases		1M NH ₄ OACpH7 (5)
K (mmolkg ⁻¹)	0.1	
Ca ²⁺ (mmolkg ⁻¹)	0.1	
Mg ²⁺ (mmolkg ⁻¹)	0.1	
Exchangeable Acidity		0.01M HCL extractant (6) and titrated against 0.1M NaOH
H ⁺ (mmolkg ⁻¹)	4.3	
Al ³⁺ (mmolkg ⁻¹)	1.5	
Micronutrients		0.1M HCL extractant (7)
Fe (mgkg ⁻¹)	8.5	
Zn (mgkg ⁻¹)	3.8	
Cu (mgkg ⁻¹)	2.0	
Mn (mgkg ⁻¹)	1.8	
Particle Size Analysis (Soil Texture)		Bouycous Hydrometer method (8)
Sand (%)	79.5	
Silt (%)	14.7	
Clay (%)	5.7	
Bulk density (Mgm ⁻³)	1.6	
% Porosity	41.8	Core method (9)

230

231

232

2.5 Statistical analysis

Analysis of variance followed by Duncan Multiple Range Test was done according to [31].

3.0 RESULTS

3.1 Treatment effects on growth and yield parameters of plantain

There were significant increases ($P \leq 0.05$) in the plant height, leaf area, leaf population/number and stem girth (Table 4) as well as plantain bunch weight, diameter of plantain fingers, length and population for two cropping years (Table 5) under different fertilizer extracts compared to the control treatment.

Among the leaf extracts, the mixed of neem leaf + wood ash had the highest values of plantain leaf area and stem girth while the neem leaf and wood ash extracts (sole forms) had the highest values of plant height and leaf population/number respectively. For example, mixed extract of neem leaf + wood ash treatment increased the leaf area and stem girth by 49.24% and 12% respectively compared to neem leaf extract. When compared with NPK 15-15-15 Fertilizer, the mixed extracts of neem leaf + wood ash treatment increased slightly plantain plant height and stem girth by 1.0%, 32.60%, 13.8% and 13.8% respectively.

In 2007/2008 cropping season, the mixed extract of neem leaf + wood ash treatment increased the plantain bunch weight, finger weight, diameter length and finger population significantly ($P=0.05$) compared to other extracts. For example, mixed extract of neem leaf + wood ash increased plantain bunch weight, finger weight, diameter, length of fruit and population by 29.7%, 31%, 6%, 8% and 25% respectively.

Generally, the plantain bunch weight, diameter and length of fingers increased significantly in 2008/2009 cropping season under the mixed extract of neem leaf + wood ash more than 2007/2008 except plantain finger population and weight of fingers.

TABLE 4: Effect of the treatments on the growth parameters of plantain

Treatments	plant height(cm)	Leaf area(cm²)	leaf population/ number	Stem girth(cm)
Control	45.6a	1454.6a	4.6a	17.7a
Wood ash extract	81.3b	4790.6d	8.2cd	29.4bc
NPK 15-15-15	85.9bc	3601.5c	6.3b	28.4b
Neem leaf extract	91.2d	2712.8b	6.8b	28.1bc
Mixed neem leaf + wood ash extract	86.5c	5344.2e	7.3c	33.0d

Treatment means within each group followed by the same letters are not significantly different from each other using Duncan Multiple Range test at 5% level.

265 **Table 5: Effect of the treatments on the yield parameters of plantain**

Treatments	Cropping year	Weight of plantain bunch (kg ha ⁻¹)	Weight of plantain fingers (kg ha ⁻¹)	Diameter of plantain fruit finger (cm)	Length of fruit finger (cm)	Plantain finger population
Control	Nov 2007	2722.7a	2221.6a	2.2a	10.9a	9.0a
Wood ash	/Mar 2008	11529.5c	10555.3d	6.5b	20.3dc	28.0cd
NPK15-15-15		12496.5d	11215.7d	6.4b	17.8b	27.0c
Neem leaf		10833.3b	9580.7b	6.6bc	18.6bc	21.0b
Mixed neem		17777.5d	16495.5e	6.8d	19.3d	36.0c
leaf + wood ash extract						
Control	Nov 2008	2752.7a	2220.6a	2.8a	13.6a	10.0a
Wood ash	/Mar 2009	11941.1c	10833.33c	6.9bc	29.8c	27.0d
NPK15-15-15		12080.0 d	10833.3c	5.8b	26.3b	26.0c
Neem leaf		9580.7 b	8611.0b	6.3d	27.8c	23.0b
Mixed neem		24166.0e	11944.1d	7.7d	28.3cd	29.0c
leaf + wood ash extract						

266 Treatment means within each column followed the same letters are not
 267 significantly different from each other using Duncan Multiple Range Test at 5%

268 **3.2 Treatment effects on soil properties after harvest of plantain**

269 There were significant increases ($P < 0.05$) in the total N, available P,
 270 exchangeable K, Ca, Mg and organic matter under different fertilizer extract compared
 271 to the control treatment after harvest of plantain (16 months after establishment) Table
 272 6.

273 Mixed extract of neem leaf + wood ash treatment increased the soil organic
 274 matter, N, P, K, Ca and Mg by 16.00%, 15.7%, 19% and 16.9% compared to the neem

leaf extract. When compared with NPK 15-15-15 fertilizer, the mixed extract of neem leaf + wood ash increased soil pH, O.M, K, Ca and Mg by 14%, 2.78%, 31%, 95.3% and 94% except N and P.

The NPK 15-15-15 fertilizer decreased significantly the soil Ca and Mg while the control treatment had the least values of the soil parameters (soil pH, O.M, N, P, K, Ca and Mg). The soil K/Ca, K/Mg and P/Mg ratios were 5:1, 6:1 and 346:1 respectively under NPK 15-15-15 Fertilizer compared to soil K/Ca 1:3, 1:2, 10:1 under mixed extract of neem leaf + wood ash respectively.

Table 6: Effect of the treatments on soil properties after harvest of plantain

Treatments	Soil pH	O.M %	N %	P mgkg ⁻¹	K mmolkg ⁻¹	Ca mmolkg ⁻¹	Mg mmolkg ⁻¹
Control	5.20a	0.28a	0.03a	1.60a	0.03a	0.03a	0.06a
Wood ash extract	6.60e	1.45c	0.06b	7.00cd	0.36cd	1.15d	0.75c
NPK15-15-15	5.36ab	0.37d	0.23d	11.73ab	0.29b	0.06ab	0.05a
Neem leaf extract	6.20c	1.4c	0.06b	6.83c	0.34c	0.85c	0.69b
Mixed neem leaf + wood ash extract	6.24d	1.68d	0.08c	8.10e	0.42e	1.29e	0.83de

Treatments within each column followed by the same letters are not significantly different from each other using Duncan Multiple Range Test at 5%

4.0 DISCUSSION

The fact that NPK 15-15-15 fertilizer improved the vegetative growth parameters such as plant height, leaf area, leaf population of plantain is consistent with its better soil N and P nutrients which are made much more readily available in mineralised forms than the organic forms in the neem leaf, wood ash extracts and mixed extract of neem leaf + wood ash.

The decrease in soil Ca and Mg in plots treated with NPK 15-15-15 fertilizer could be due to nutrient imbalances reflected in high K/Ca, K/Mg and P/Mg ratio which consequently affected uptake of Ca and Mg by plantain. In addition, the low values of soil organic matter, Ca and Mg under the NPK 15-15-15 fertilizer plot signified low base saturation, and the sandy loam texture also encouraged soil leaching activities and poor nutrients retention.

These observations could explain the reduction in values of plantain growth and yield parameters i.e. plant height, leaf area, stem girth, leaf population, weight of plantain bunches, weight of plantain fingers and fingers population under the NPK 15-15-15 fertilizer compared to the mixed extract of neem leaf + wood ash treatment which contained Ca, Mg and organic matter and subsequently increased soil fertility status for higher values of growth and yield parameters of plantain. (32) reported that continuous use of ammonium sulphate and NPK 15-15-15 Fertilizer decreased soil pH, K and macronutrients in okra, also (33) reported that arbitrary use of inorganic fertilizers resulted into signs of toxicities, poor yield responses and deterioration of soil properties in okra.

The performance of neem leaf extract in improving the growth and yield parameters of plantain could be traced to its high N, P, K, Ca and Mg nutrients

311 supplied to the soil which subsequently increased the soil organic matter, total N,
312 available P, exchangeable K, Ca and Mg.

313 This also reflected in the increased values of the plant height, leaf area, stem
314 girth, leaf population, plantain bunch weight, fingers weight, length, diameter and
315 population. The finding was also supported by [8] who reported that neem seed extract
316 enhanced the nitrogen use efficiency in soils as well as increasing the yield of crops.
317 Also, [34] also reported the use of neem leaf extract increased soil nutrients, growth
318 and yield parameters of tomato.

319 In addition, the moderate potassium content of neem leaf extract also
320 contributed to better performance of plantain yield parameters and this was supported
321 by [35] who reported the vital role of potassium in crop production.

322 The wood ash extract increased the values of plantain bunch weight, weight of
323 fingers, diameter, length and finger population more than neem leaf and this could be
324 traced to its very high K, Ca and Mg nutrients which subsequently increased the soil
325 organic matter, K, Ca and Mg for uptake by plantain. This finding was supported by
326 [36] who reported that wood ash is a good source of potash and lime which
327 encouraged the growth and yield of plantain.

328 The increase in soil pH under wood ash extract compared to others was traced
329 to its high K, Ca and Mg and could be effective as a liming material [37] unlike the
330 NPK 15-15-15 fertilizer which with the continuous use could decrease soil pH. The
331 soil pH been reported to influence nutrient uptake and availability. [38] reported that
332 oil palm bunch ash, wood ash and cocoa pod husk improved soil K, Ca and Mg

nutrients and corrected soil acidity in an Alfisol grown to coffee and maize. [39] also reported that industrial wood ash was a good soil amendment for crop production.

The highest values of plantain bunch weight, weight of fingers, diameter of finger, length of fingers and finger population under the mixed extract of neem leaf + wood ash treatment could be due to the fact that it had the combined nutrient superiority of P, K, Ca Mg and N compared to neem leaf and wood ash extracts applied in sole forms. Therefore, the positive impact of extract of neem leaf + wood ash also reflected in the increase in soil pH, organic matter, N, P, K, Ca and Mg and the subsequent uptake by the plantain.

Hence, the impact of the mixed extract on the soil properties has been responsible for high yield parameters, plantain bunch weight, finger weight, diameter and length of fingers and finger population which further signified higher economic returns, food security and profitability for plantain farmers as well as providing raw materials to establish plantain flour, chips and biscuit industries. It is believed that this would promote plantain values chain development for export trade.

The observation was supported by [34] who reported superior performance of mixed extract of neem leaf + wood ash treatment in maize and water melon intercrop. Besides, the uptake of nutrients (N, P, K, Ca and Mg) in a liquid form from the mixed extract of neem leaf + wood ash could also be responsible for better plantain yield performance over the solid organic fertilizers which must be mineralized with adequate moisture before uptake by plant roots. Also, it has residual values which would have contributed to plantain bunch weight in the 2nd cropping year which NPK 15-15-15 fertilizer did not possess for subsequent cropping.

The poor growth and yield performance of plantain in the control treatment was consistent with the fact that the soil was very low in nutrient contents and this observation was supported by [13] who reported poor growth and yield responses of crops in soils that are not fertilized. The lowest values of soil N, P, K, Ca and Mg in the control treatment were a reflection of poor soil fertility status, hence, there is need to ensure fertilization of soils for bumper crop productivities in the tropical region where soils are low in organic matter, K, Ca, Mg and low soil pH [12].

In term of nutritional status, plantain has high iron, minerals and vitamins contents which helped to build up both physiological and metabolic activities in the body when consumed by people. This will reduce money spent on purchase and consumption of iron and vitamins based drugs and this observation was in line with meeting millennium development goals (MDGs) in food security and sound health for people in the world.

5.0 CONCLUSION

It could be concluded from the research work that mixed extract of neem leaf + wood ash treatment gave the highest values of plantain growth and yield parameters and also improved the soil nutrients (soil N, P, K, Ca, Mg and organic matter) compared to NPK 15-15-15 fertilizer, neem leaf and wood ash extracts (sole forms) respectively. It is recommended that for better performance of plantain bunch weight, finger weight, finger diameter and length, growth parameters and improvement of soil fertility status, application of mixed extract of neem leaf + wood ash at 833.3 Lha^{-1} is appropriate.

379 Hence, the use of mixed extract of neem leaf + wood ash could substitute for
380 300 kg ha^{-1} of NPK 15-15-15 Fertilizer and it also reduced the problems of high cost of
381 purchasing the inorganic fertilizer as well as solving problems of bulkiness and
382 difficulty in the transportation of solid organic waste. Finally, it will help to prevent
383 environmental pollution and deterioration of soil properties associated with
384 continuous use of inorganic fertilizers such as NPK 15-15-15 fertilizers.

385 |

386

387 **6.0 REFERENCES**

- 388 1. **Wilson GF:** Status of banana and plantain in West Africa. Proc workshop on
389 Banana and Plantain breeding strategies. Cairns Australia .1986: 27 – 35 .
- 390 2. **Swennen R, Wilson GF:** Responses of plantain to mulch and fertilizer.
391 International Institute of Tropical. Agriculture. Annual Report, IITA Ibadan,
392 Nigeria .1983: 187.
- 393 3. **Shaibu AA, Maji EA, Ogburia MN:** Yield evaluation of plantain and banana
394 lanraces and hybrids in humid agro-ecological zone of Nigeria. Journal of
395 Agric Research and Development. 2012:2(1):74 – 79.
- 396 4. **Agbein NO, Emechebe AM, Marley PS, Akpa AD:** Evaluation of
397 Nematicidal Action of some Botanicals on Meloidogyne Incognita In vivo and
398 In Vitro. Journal of Agriculture and Rural Development in the Tropics and
399 subtropics 2005: 106 (1): 29 – 39
- 400 5. **Hossain MB, Poehling HM, Thoming G, Borgemeister BE:** Effects of soil
401 application of neem (Neem Azaal ® - U) on different life stages of
402 Liriomyzasativae (Diptera: Agromyzidae) on tomato in the humid tropics.
403 Journal of Plant Diseases and Protection. 2008:115 (2): 80 – 87.
- 404 6. **SubbalakshmiLokanadhan, Muthukrishnan O, Jeyaraman S:** Neem
405 products and their agricultural applications. JBiopest.2012:.5 (Supplementary).
406 :72-76 .
- 407 7. **Santhi SR, Palaniappan SP, Puroshothaman D:** Influence of neem leaf on
408 nitrification in a lowland rice Soil. Plant and Soil. 1986: 93(1):133-135.

- 409 | 8. **Zhang X, Shen Q, Tan J, Mao Z.** Effect of neem seed extract on nitrogen use
410 | efficiency in two different soils (in Chinese). Ying Yong Sheng Tai XueBao.
411 | 2002 : 11(1): 1417 – 1420.
- 412 | 9. **Alberta E.** Standard and guidelines for the use of wood ash as a liming
413 | material for agricultural soils. 2002: Pub No: T/666. ISBN: O-7785-2280-6
414 | (Printed Edition. ISBN: 0-7785-2281-4 (On-line Edition: Website:
415 | <http://www.gov.ab.ca/env/>
- 416 | 10. **Etiegni L, Mahler RL, Campbell AG. :Effect of wood ash source of**
417 | **nutrient for seleted crops** Soil Sci. Plant Anal. 1991: 22(1): 257-259.
- 418 | 11. **Moyin-Jesu EI.** Incorporation of Agro-Industrial Biomass and their effects on
419 | growth and nutrient content of Arnaranthus. Pertanika Jour of Trop. Agric
420 | Science Malaysia .2003.:26(1): 49-58.
- 421 | 12. **Moyin-Jesu EI.** Evaluation of different organic fertilizers on the soil fertility,
422 | leaf chemical composition and growth performance of coffee seedlings.
423 | African Journal of Science. and Technology. 2007:2 (2):1-6
- 424 | 13. **Moyin-Jesu EI, Atoyosoye B:** Utilization of Agro-Agricultural Wastes for the
425 | growth, leaf and soil chemical composition of cocoa seedling in the nursery.
426 | Pertanika Journal of Tropical Agric Science. 2002.:25(1):53-63.
- 427 | 14. **Ivbijaro MF.** Preservation of cowpea with neem seeds. Fuss ProtecoEcology
428 | Journal. 1983: 1(5). 127 – 132..
- 429 | 15. **Moyin-Jesu EI:** Evaluation of modified neem leaf, wood ash and neem leaf,
430 | wood ash and neem leaf extracts as pest control and seed treatment in maize.
431 | Emirate Journal of Agric and Food. 2010: 22(1): 1-8.

- 432 | **16. Moyin-Jesu EI:** Comparative evaluation of modified neem leaf, neem leaf and
433 | wood ash extracts on soil fertility improvement, growth and yield of maize
434 | (Zea mays L.) and watermelon (sole and intercrop). Scientific Research
435 | Publishing. 2012a;11 (1):1-8. DOI identifier: 10. 4236.
- 436 | **17. Soil Survey Staff.** Soil Taxonomy A basic system for soil classification for
437 | making and interpreting soil surveys.1999:USDA Handbook No.436,
438 | Washington, D.C, USA.
- 439 | **18. Jackson ML.** Soil chemical analysis .Englewood cliffs N.J. 1958 Prentice
440 | Hall :59 – 67.
- 441 | **19. Association of Analytical Chemists:** Official methods of analysis 12th Edition.
442 | Association of Official Analytical Chemists, 1970 Arlington,VA.
- 443 | **20. Crockford L, Nowell R.** Laboratory manual of physical chemistry.. Exp 31
444 | and 32. John Wiley and Sons, New York. 1956: 58 – 59.
- 445 | **21. Walkley A, Black IA.** An examination of Degtjaroff method for determining
446 | soil organic matter and a proposed modification of the chromic acid filtration
447 | soil Sci.1934: 37: 29 – 38.
- 448 | **22. Murphy J, JP Riley.** A modified single solution method for determination of
449 | phosphate in natural waters. Analytical Chem. Acta. 1962:27,:31 – 36.
- 450 | **23. Jackson ML.** Soil chemical analysis .Englewood cliffs N.J. Prentice Hall.
451 | 1954: 59 – 67.
- 452 | **24. Mclean EO.** Aluminum:Methods of soil analysis Part 2, Agronomy 9,
453 | Americal Society of Agronomy. Madison Wisconsin, U.S.A.MCA Black
454 | (eds).1962:927-932.

- 455 | 25. **Ogunwale JA, Udo EJ.** A laboratory manual for soil and plant analysis.
456 | Agronomy Dept. University of Ibadan, Nigeria.1978: 201 – 206.
- 457 | 26. **Bouycous H.** Mechanical analysis of soil using hydrometer method. Analytical
458 | Chem Acta.1951: 22(1):32-34.
- 459 | 27. **Ojeniyi SO.** Effect of replacement of grass and forest with crops on soil
460 | textural and chemical properties. A paper presented at international conference
461 | on soil fertility, soil tilth and post clearing land degradation in the humid
462 | tropics. Eds. Babalola A. Proceedings 11th International Conference held at
463 | University of Ibadan. 1985: 20 – 27.
- 464 | 28. **Agboola AA , Corey,RB.** Soil testing N, P, K for maize in the soils derived
465 | from metamorphic and igneous rocks of Western State of Nigeria. Nigeria
466 | Journal of West African Science Assoc. 1973: 17(2): 93-100.
- 467 | 29. **Sobulo RA, Osiname OA.** Soils and fertilizer use in Western Nigeria. Tech
468 | Bull, No 11, Institute of Agricultural Research and Training, University of Ife,
469 | Nigeria. 1981: 8-9 .
- 470 | 30. **Adeoye GO.** Comparative studies of Ammonium bi fluoride chelate extractants
471 | and some convectional extractant for sedimentary soils of S. W. Nigeria. PhD.
472 | Thesis, Faculty of Agric and Forestry, University of Ibadan, Nigeria.1986:120-
473 | 124.
- 474 | 31. **Gomez KA,Gomez AA.** Statistical procedure for Agricultural Research 2nd
475 | edition, John Wiley and sons. New York.1984 : 20-24.

- 476 | **32. Aduayi EA.** Effect of ammonium sulphate fertilization on soil chemical
477 | composition, fruit yield and nutrient content of okra. *Ife Journal of Agric.*1980
478 | : 2 (1):16-33.
- 479 | **33. Moyin-Jesu EI, Ojeniyi SO.** Effects of sole and amended plant residues on
480 | soil nutrient contents and yield of Okra (*Abelmoschus esculentum* L).
481 | *Discovery and Innovation Journal.*2006:18(4): 318 – 326
- 482 | **34. Moyin-Jesu EI:** Comparative evaluation of modified neem leaf, neem leaf and
483 | wood ash extracts on soil fertility status, growth and yield of
484 | tomato(*Lycopersicon esulentum* L.).*Global Journal of Bio-Science and*
485 | *Biotechnology*, 2012b: 1(2):271-276.
- 486 | **35. Adams SA.** The role of potassium in Agriculture.*American Society of*
487 | *Agronomy*, Maldison, WI .1986:3(1):22-25.
- 488 | **36. Etiegni L, Mahler RI, Campbell AG .Potential of wood ash as source of**
489 | **fertilizer for crop.** *Soil Science Plant Analysis.*1991 :22(1) :257
- 490 | **37. Gordon W.** Coffee Tropical Agricultural Series In: **H.Murray (Eds).**
491 | *Macmillian Publishing Ltd. London.*1988 :1-20.
- 492 | **38. Obatolu, CR. 1995:** Nutrient balance sheet of Alfisol grown to coffee and
493 | maize using organic fertilizers. In: *Proceeding of third annual conference of All*
494 | *African Soil Science Society on Soil management and environmental*
495 | *protection held at University of Ibadan, Ibadan, Nigera.(eds*
496 | *A.A.Agboola).*1995 :250 – 256.
- 497 | **39. Meyers NL, Kopecky MJ.** Industrial wood ash as a soil amendment for crop
498 | production. 1998. *TAPPI Journal.*1998: 81 (4): 123-130.

499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517

ACKNOWLEDGEMENT

The author acknowledged the support received from God almighty and my family.

COMPETING INTERESTS

There is no competing interests exist.

AUTHORS' CONTRIBUTION

ETHICAL APPROVAL

The study is not against the public interest.