

Original Research Article**Contributions of Rhizobium Inoculants and Phosphorus Fertilizer to Biological Nitrogen
Fixation, Growth and Grain Yield of three Soybean varieties on a Fluvic Luvisols****ABSTRACT**

An experiment to identify an efficient strategy to optimize biological nitrogen fixation (BNF) in three soybean varieties {Jenguma (TGx1448-2E), Anidaso and Quarshie (TGx1445-2E)} was conducted in the experimental field of the CSIR-Savanna Agricultural Research Institute, Nyankpala, Northern Region, Ghana. . The experiment had a split-split plot design with three replications and interactions tested were inoculation rate (0, 50 and 100% inoculation) (main plot), soybean variety (sub-plot) and phosphorus rate (0, 22.5 and 45.0 kg P_2O_5 /ha) (sub-sub plot) using maize as a reference crop. The soybean was sown at two seeds/hill at a spacing of 50 cm x 10 cm. The results showed that inoculating soybean with the commercial inoculant Legumefix (Rhizobium strain 532c) had no effect on plant height, nodule number, nodule dry weight, shoot dry weight, pod number, grain yield and 100-seed weight. However, the soybean varieties showed significant differences in pod number, 100-seed weight and phosphorus uptake efficiency (PUE). The amount of N_2 fixed (measured by Total Nitrogen Difference method) ranged between 52.3-71.0 kg N/ha. Phosphorus applications of 22.5 and 45.0 kg P_2O_5 /ha increased grain yield by 35.4 and 33.9%, respectively and also increased N_2 fixation by 49.39 and 69.82%, respectively over the unfertilized control. The interactions among these treatments did not significantly influence the parameters measured except PUE for which there were significant differences among the soybean varieties and the phosphorus rates.

Inoculation did not therefore increase nodulation and BNF of the three soybean varieties but phosphorus application increased the growth and grain yield of the soybeans.

Keywords: *Inoculation rates, soybean varieties, phosphorus rates, BNF, grain yield*

INTRODUCTION

Soybean (*Glycine max* (L.) Merrill) is a legume plant belonging to the botanical family *leguminosae*. It is an economically important leguminous crop worldwide and also the most important legume in Ghana (Plahar, 2006). According to Dugje *et al.* (2009), soybean is more protein-rich than any of the common vegetable or legume food sources in Africa. It is a source of edible oil (20-25%) with 42-45% protein content (Alam *et al.*, 2009). Soybean is a promising pulse crop proposed for the alleviation of the acute shortage of protein and oil worldwide (Mahamood *et al.*, 2009). It used as a good source of unsaturated fatty acids, minerals (Ca and P) and vitamins A, B, C and D (Alam *et al.*, 2009).

Despite these good characteristics, its growth and productivity is affected mainly by nitrogen even though this element is the most abundant element on the earth and about 78% of the earth's atmosphere is nitrogen gas. Despite the abundance of nitrogen in the atmosphere, plants are unable to use it directly because it is present in an inert form (N₂) and the nitrogen in the soil is lost through microbial dinitrification, soil erosion, leaching, chemical volatilization, removal of nitrogen- containing crop residues from the land. Nitrogen is therefore the most limiting plant nutrient for crop production in West Africa (Sangakara *et al.*, 2003). Most legumes, however, through symbiosis with rhizobia have the ability to reduce N₂ through biological nitrogen fixation (BNF) into a plant-usable form.

Currently, BNF contribution on majority of smallholder farms rarely exceeds 5 kg N/ha/year with nitrogen-fixing legumes in Ghana (Mapfumo, 2001). A measure of more than 240 kg N/ha of fixed N₂ in soybean in southern Africa on smallholder farms with associated grain yield of more than 3.5 t/ha has been recorded (Giller, 2001). This shows that the potential rates of BNF in soybean are not only limited by the efficiency of legume-rhizobium symbiosis but environmental and management factors could also limit the efficiency. These factors may include temperature, rainfall, soil or fertilizer nutrients like nitrogen and phosphorus, soil pH, etc. Phosphorus is an essential ingredient for Rhizobium bacteria to carry out BNF processes. Inadequate P restricts root growth, the process of photosynthesis, translocation of sugars and other such functions which directly influence N fixation by legume plants.

The general objective of this study was therefore to identify efficient strategies that will optimize biological nitrogen fixation in soybean for increased crop yield. The specific objectives were: (i) To evaluate the effect of Rhizobium inoculation and P fertilizer application rate on nodulation and BNF of three soybean varieties. (ii) To evaluate the effect of Rhizobium inoculation and P application on growth and yield of three soybean varieties. (iii) To determine the effect of Rhizobium inoculation and P application on P uptake efficiency of three soybean varieties.

MATERIALS AND METHODS

Study area: The study was carried out on the experimental field of the Savanna Agricultural Research Institute at Nyankpala in the Northern Region of Ghana from June to December 2012. Nyankpala is located the interior Guinea Savanna agro-ecological zone of Ghana about 16 km west of Tamale and lies on latitude N 09° 24' 15.9" and longitude W 01° 00' 12.1'. The climate is relatively dry with a mono-modal rainfall pattern. Average ambient temperatures are high year

round (about 28°C) but the months of December and January are characterized by minimum temperatures that may fall to 13°C at night while March and April may experience temperatures of 40°C in the early afternoon. The mean annual rainfall of the area is 1000 mm with a total of 1022 mm. The soil at the trial site is Fluvic luvisols which is classified as the Tingoli series and the vegetation type is a tree Savannah.

Soil sampling, preparation and analysis: Soil samples were taken before the field was ploughed with a tractor. Five soil cores were taken at the depth of 0–15 cm from each replicate plot and bulked to obtain a composite samples Soil samples were air-dried and sieved through a 2-mm mesh before subjecting them to chemical and physical analyses using standard laboratory procedures. Particle size analysis was done by using the hydrometer method (Anderson and Ingram, 1993). Soil pH was measured in the supernatant suspension of 1: 2.5 soils and water mixture using a pH meter. Soil organic carbon was determined by the method of Walkley and Black (1934). Whilst total nitrogen of the soil was determined by the Kjeldahl method (Bremner and Mulvaney, 1982) available phosphorus was determined by the Bray 1 method (Bray and Kurtz, 1945). Cation Exchange Capacity (CEC) was determined by leaching the soil with neutral 1 N ammonium acetate (FAO, 2008).

Soil Characteristics: The physical and chemical analyses revealed that the experimental site is loamy-sand in texture and neutral in reaction (pH 5.5-7.0) with a very low organic carbon content (<20 g/kg) (Table 4.1) Total nitrogen and exchangeable cations were less than 1 g/kg and 5 cmol ⁽⁺⁾/kg, respectively with effective cation exchange capacity and available P being less than 5 cmol ⁽⁺⁾/kg and 10 mg/kg soil, respectively according to Landon (1991).

91 **Table 1: Some physical and chemical soil properties of the study area**

Parameter	Test Value
Sand (g/kg)	855.00
Silt (g/kg)	83.00
Clay (g/kg)	62.00
Texture	Loamy-sand
pH (1:2.5; soil:H ₂ O)	6.70
Organic carbon (g/kg)	11.00
Total N (g/kg)	0.50
Available P (mg/kg)	2.20
Exchangeable cations (cmol ⁽⁺⁾ /kg):	
Ca	1.50
Mg	0.70
K	0.08
Na	0.06
Exchangeable acidity (cmol ⁽⁺⁾ /kg)	0.73
Effective cation exchange capacity (cmol ⁽⁺⁾ /kg)	3.07

92

93 **Experimental design and treatments:** The experiment was designed in a split-split plot with
94 three replications and treatments tested were inoculation rate (0, 50 and 100 % inoculation) as
95 the main plot, soybean variety {Jenguma (TGx-1448-2E), Anidaso, Quarshie (TGx-1445-2E)} as
96 the sub-plot and phosphorus rate (0, 22.5 and 45 kg P₂O₅/ha) as the sub-sub plot. The inoculants
97 was a commercial peat-based one of *Bradyrhizobium japonicum* strain 532c. The test soybeans
98 are medium-maturing (95-110 days) with 45 days to 50% flowering and have a potential grain
99 yield of 1.8-2.5t/ha.

100 The plot size was 3 m x 4 m and spacing between plots and blocks were 1 m and 2 m,
101 respectively. For the purpose of assessing BNF, a non-fixing reference crop (maize) was added

to the treatment combinations. Phosphorus was applied as triple super phosphate to the test crops at the rate of either 22.5 or 45 kg P_2O_5 /ha but not to the reference crop. However, muriate of potash (MoP) was applied to all experimental plots at 30 kg K/ha. The fertilizers were deposited in holes 5 cm away from the plants and covered with the soil at two weeks after sowing.

Seeds inoculation: The inoculants were added to the soybean seeds in a container after moistening the seeds. The inoculants and the seeds were mixed thoroughly until the seeds were adequately coated with the inoculants and allowed to air-dry on a sheet of canvas in the shade for a few minutes after which they were planted on the ridges. The treatment with 100% inoculation rate received 5 g of inoculants per 1 kg of seed whilst the half recommended inoculation rate (50%) was done by mixing 1 kg of the seeds with 2.5 g of inoculants.

Data collection: Plant height, shoot dry weight, nodule dry weight and nodule number per plant were recorded from ten representative plants at eight weeks after planting (8 WAP). Plant height was taken from the ground level to the apex of the plant with a graduated pole and the average was calculated for each plot. After the root systems of the 10 plants were cut and gently washed on a 2-mm mesh sieve under a jet of tap water, the nodules were detached, counted and oven-dried at 65°C for 48 h. Pods on these ten plants were removed and counted to obtain the pod load (i.e. pod number/plant). The shoots of five plants sub-sampled from the ten plants harvested were also oven-dried at 65 °C for 48 h and the weights recorded.

After threshing the pods harvested in the harvest area of each treatment plot, the grains were sufficiently sun-dried on a concrete platform and weighed on an electronic balance. Hundred

124 seeds from each treatment were randomly picked and weighed. This was replicated three times
125 and the average 100-seed weight determined.

126 **Estimation of N₂ fixed:** The technique used to estimate N fixation was the Total Nitrogen
127 Difference (TND) method. This was done by comparing total nitrogen of the legume with that of
128 a non-legume (Murray *et al.*, 2008). The amount of N fixed was calculated by subtracting total
129 nitrogen of the reference crop (maize) from that of the legume (soybean), and the difference
130 value is assumed as N derived by BNF (N₂ fixed).

Thus, N₂ fixed = Total N in legume - Total N in reference crop

$$\text{where Total N in plants} = \frac{(\text{Dry matter weight } \left(\frac{\text{kg}}{\text{ha}}\right) \times \% \text{ N in plants})}{100}$$

131 Then shoot nitrogen content was analyzed using the Kjeldahl procedure (FAO, 2008).

132 **Phosphorus Uptake efficiency:** Phosphorus uptake efficiency (PUE) was estimated using an
133 equation by Khair *et al.* (2002).

$$\text{PUE} = 100 \times \left[\frac{\text{P in fertilized plants} - \text{P in control plants } \left(\frac{\text{kg}}{\text{ha}}\right)}{\text{P applied } \left(\frac{\text{kg P}_2\text{O}_5}{\text{ha}}\right)} \right]$$

134
135 **Statistical analysis:** All data collected were subjected to statistical analysis using Genstat
136 Discovery Edition 10. Nodule count was transformed before the analysis. Analysis of variance
137 (ANOVA) was done to determine differences in means among treatments. All treatment means
138 were compared using the Least Significant Difference (LSD) at 5 % level of significance.

139

140 **RESULTS AND DISCUSSION**

Plant height: Application of inoculants, soybean variety and their interactions had no significant (P>0.05) effect on plant height at 8 WAP unlike phosphorus application which significantly (p<0.05) affected it (Table 2). Phosphorus applications of 22.5 and 45 kg P₂O₅/ha significantly increased plant height by 2.76% and 6.87%, respectively over the control treatment (0 kg P₂O₅/ha). This finding is in agreement with reports of Bekere *et al.* (2012) that phosphorus application of 60, 120 and 180 mg P/kg significantly influenced soybean height under controlled environments.

Shoot dry weight: Shoot dry weight (biomass) at mid flowering was not significantly influenced by inoculation rate, variety and their interactions (Table 2). Similar shoot dry weights produced by the test soybean varieties at mid-flowering under similar conditions of growth shows that they have equal growth and biomass production potentials as stated by Salisbury and Ross (1992). Contrarily, phosphorus application of 22.5 and 45 kg P₂O₅/ha significantly enhanced biomass production in the soybean crops by 35.6 and 53.6%, respectively as compared to the no P treatment with the 45 kg P₂O₅/ha rate yielding higher biomass than the lower P rate (Table 2). This result is in agreement with Bekere *et al.* (2012) who concluded that omission of P from optimum nutrition of soybean dramatically reduced shoot dry matter yield of soybean.

Table 2: Effects of inoculation rate, variety and phosphorus rate on plant height and shoot dry weight of soybean

Treatment	Plant height (cm)	Shoot dry weight (kg/ha)
<u>Inoculation rate (%)</u>		
0	69.99	5604
50	69.79	4984
100	69.24	5089

166	Pr (I)	0.826	0.269
167	Lsd (0.05)	3.26	956.7
168	<u>Variety</u>		
169	Jenguma	69.34	5198
170	Quarshie	69.13	5647
171	Anidaso	70.55	4833
172	Pr (V)	0.443	0.595
173	Lsd (0.05)	2.53	1704.8
174	<u>Phosphorus rate (kg P₂O₅/ha)</u>		
175	0	67.38	3331
176	22.5	69.29	5171
177	45	72.35	7176
178	Pr (P)	<0.001	<0.001
179	Lsd (0.05)	1.05	646.5
180	Pr (I x V)	0.950	0.762
181	Pr (I x P)	0.851	0.835
182	Pr (V x P)	0.304	0.184
183	Pr (V x I x P)	0.371	0.383
184	CV (%)	6.94	22.4

185

186 **Nodule number and dry weight:** There was no significant ($P > 0.05$) effect on nodule number

187 and dry weight of soybean due to application of inoculants, variety and their interactions (Table

188 3). The lack of nodulation response to rhizobium inoculation might suggest that N was not a

189 limiting nutrient at this site. Other nutrients such low phosphorus and molybdenum could affect

190 inoculation response and also, native rhizobia can prevent the rhizobia introduced in the

191 inoculants from forming nodules on the crop. It could also be possible that in this study the

192 native rhizobia might have also fixed as much nitrogen as the plant needs making inoculation

193 unnecessary. Contrarily, significant positive effects of high rates of inoculation (high numbers of

rhizobia per seed) have been demonstrated by Roughley *et al.* (1993) and Date (2001). With the application of P, however, there was a significant ($P < 0.05$) enhancement in nodulation of soybean and the degree of enhancement was dependent on the rate of P applied (Table 3). Phosphorus application of 22.5 and 45 kg P_2O_5 /ha significantly increased number of nodules by 11.85 and 21.71%, respectively over the control treatment. This significant positive effect of phosphorus on nodulation underlines the influence phosphorus has on nodule development through its basic functions as an energy source (Singleton *et al.*, 1984).

Table 3: Effects of inoculation rate, variety and phosphorus rate on nodule number and dry weight of soybean

Treatment	Nodule Number (no/plant)	Nodule Dry Weight (mg/plant)
<u>Inoculation rate (%)</u>		
0	1.37	2233
50	1.35	1904
100	1.34	2049
Pr (I)	0.89	0.67
Lsd (0.05)	0.18	978.2
<u>Variety</u>		
Jenguma	1.38	2237
Quarshie	1.42	2289
Anidaso	1.26	1659
Pr (V)	0.106	0.109
Lsd (0.05)	0.16	658.4
<u>Phosphorus rate (kg P_2O_5/ha)</u>		
0	1.19	686
22.5	1.35	2304
45	1.52	3196
Pr (P)	<.001	<.001
Lsd (0.05)	0.05	1186.4

223	Pr (I x V)	0.956	0.944
224	Pr (I x P)	0.719	0.417
225	Pr (V x P)	0.896	0.451
226	Pr (V x I x P)	0.103	0.250
227	CV (%)	7.0	23.4

228

229 YIELD AND YIELD COMPONENTS

230 **Pod number:** Inoculation rate and the interactions had no significant ($P > 0.05$) effect on pod
 231 number though inoculated treatments (50 and 100% inoculation) apparently had more pods than
 232 noninoculated ones whereas phosphorus rate and varietal differences significantly ($p < 0.05$)
 233 affected the number of pods (Table 4). Application of 22.5 and 45 kg P_2O_5 / ha increased pod
 234 number by 33.39 and 47.87%, respectively over the control treatment and this result was
 235 supported by the findings of Mohan and Rao (1997) that higher numbers of pod were produced
 236 when higher doses of phosphorus were applied

237 The observation with inoculation contradicts the findings of Shahid *et al.* (2009) that seed
 238 inoculation of soybean produced more pods per plant than un-inoculated control. Bhuiyan *et al.*
 239 (2008) and Malik *et al.* (2006) also reported that pod number of mung bean and soybean were
 240 significantly increased by inoculating with *Bradyrhizobium*. The variety Quarshie produced
 241 mean pod number of 66.1 which was statistically ($P < 0.05$) higher than the mean pod number of
 242 Anidaso but statistically similar to that of Jenguma . Genetic factors of the soybean varieties may
 243 have contributed to the significant differences in their pod loads which may be affirmed by the
 244 assertion of Bouquet (1998) that genotype selection is one of the most important factors for
 245 increasing pod yield in soybean.

246 **Grain yield and grain size:** Inoculant had no significant ($P > 0.05$) influence on both grain yield
 247 and seed size (100-seed weight) (Table 4). The lack of grain yield response to inoculation can be

attributed to the fact that inoculation did not increase nodulation (Table 3) and N₂ fixation (Table 5) and this may indicate an abundance of effective native soybean rhizobia at this location. Other factors may include cultivar and strain interaction and drought. Graham, (1992) reported that drought affects symbiosis between host and rhizobia and this influences rhizobial survival in the soil, the host or the process of nodulation and grain yield. Lack of significant effect on yield improvement by inoculation has also been reported by other workers (Chemining'wa *et al.*, 2004; Otieno *et al.*, 2007). Varietal differences were observed to be significant ($p < 0.05$) for seed size only (Table 4) and the seeds of inoculated Jenguma were statistically bigger (12.57 g/100 seeds) than the seed size (10.8 g/100 seeds) of Anidaso but statistically similar to that of Quarshie (12.6 g/100 seeds). Seed weight has been noted to affect seedling vigour and it was reported that genotypes with heavier seeds produced more vigorous seedlings and the relationship between seed weight and seedling vigor was more pronounced in the earlier growth stage (Qiu and Mosjidis, 1993). Similar to this result, Solomon *et al.* (2012) reported significant difference ($p \leq 0.01$) in 100-seed weight by the main effect of variety. Phosphorus rate significantly ($p < 0.05$) increased both grain yield and 100-seed weight (Table 4). Phosphorus applications at 22.5 P₂O₅ /ha and 45 kg P₂O₅ /ha induced significantly similar responses with respect to grain yield and seed size but both were significantly higher than that of the control (0 kg P₂O₅/ha). Soybean has a relatively high requirement for phosphorus and yield and seed quality can be enhanced by phosphorus fertilizer in soils testing low in phosphorus (Shahid *et al.*, 2009). Bakere and Hailemariam (2012) also reported that seed yield of soybean increased with phosphorus fertilizer applications of 20, 40 and 60 kg P₂O₅/ha.

Table 4: Effects of inoculation rate, variety and phosphorus rate on pod number, grain yield and 100-seed weight of soybean

272	Treatment	Pod Number	Grain Yield	100-seed Weight
273		(no./plant)	(kg/ha)	(g)
274	<u>Inoculation rate (%)</u>			
275	0	55.6	1545	12.1
276	50	57.9	1646	11.8
277	100	59.3	1815	12.1
278	P (I)	0.31	0.26	0.39
279	Lsd (0.05)	5.83	385.60	0.13
280	<u>Variety</u>			
281	Jenguma	60.5	1881	12.7
282	Quarshie	66.1	1560	12.6
283	Anidaso	46.2	1566	10.8
284	P (I)	0.018	0.078	<.001
285	Lsd (0.05)	13.3	318.10	0.58
286	<u>Phosphorus rate (kg P₂O₅/ha)</u>			
287	0	39.1	1233	11.5
288	22.5	58.7	1909	12.2
289	45	75.0	1864	12.36
290	P (P)	<.001	<.001	0.002
291	Lsd (0.05)	7.32	258.80	0.49
292	P (I x V)	0.874	0.265	0.993
293	P (I x P)	0.916	0.901	0.195
294	P (V x P)	0.337	0.267	0.653
295	P (I x V x P)	0.595	0.402	0.908
296	CV (%)	23.0	28.10	7.40
297				

298 **N₂ fixation:** The amount of N₂ fixed in soybean was not significantly (p > 0.05) affected by
 299 inoculation, variety and the interactions (Table 5). The failure of inoculation to enhance the
 300 amount of N₂ fixed might be due to the inability of the inoculation to increase nodulation in

soybean though Ruark (2009) has reported of an increased N_2 fixation in soybean due to increased nodulation. This observation is contrary to the report that the amount of N_2 fixed in *Phaseolus vulgaris* L. was increased significantly with *Rhizobium* inoculation in all organs (roots, shoots, pods and whole plant) both in glasshouse and field experiments compared with the un-inoculated control treatment (Bambara, 2009). The non-significant difference of variety on the amount of N_2 fixed supports the facts that soybean maturity date affects the amount of N_2 fixed in soybean. N_2 fixation has been reported to increase with increasing crop duration (Ogoke *et al.*, 2006) because longer growth duration allows for a longer period of N_2 -fixation in the nodules. Increased crop duration in the field means a longer period of nodule activity. The soybean varieties used in this study have the same maturity period (they are medium-maturing varieties). This might explain why the difference in the amount of N_2 fixed was not significant among the three varieties. .However, the amount of N_2 fixed at the experimental site (52.1 to 71 kg N / ha) in the soybean varieties studied was within the range of 15-162 kg N/ha estimated for soybean by Giller and Wilson (1991) and Larue *et al.*(1981) in the tropics. . This is higher than the 41-50 kg N / ha reported by Yusuf *et al.* (2006) but lower than the 61-109 kg N/ha obtained by Ogoke *et al.* (2006). Unlike with the varieties, phosphorus application significantly enhanced N_2 fixation at both rates of 22.5 and 45 kg P_2O_5 / ha resulting in increases of 49.39 and 71.90 % of fixed nitrogen, respectively over the control (Table 5). This indicates that P deficiency which limits plant growth can also seriously limit symbiotic N_2 fixation as the latter has been noted to have a higher P requirement for optimal functioning than either plant growth or nitrate assimilation (Ogoke *et al.*, 2006).

Phosphorus uptake efficiency: Whereas application of the commercial inoculants had no significant effect on phosphorus uptake efficiency (PUE), remarkable differences in PUE were

observed among the three soybean varieties used as well as the P levels applied (Table 5). Jenguma was the most efficient in taking up the fertilizer P applied followed by Anidaso and Quarshie. The differences in the PUEs of the test legume varieties may suggest that differences in PUE could occur among plant species or genotypes due to differences in the abilities of their root systems to acquire P from the soil and accumulate it in the shoots (Rao *et al.*, 1997). Soybean varieties are known to differ in their ability to grow under low-P conditions and the more P-efficient varieties may have internal and/or external mechanisms that allow greater soil P extraction and grain yield (Neue, 1991). In this work, the higher grain yield of Jenguma compared to the other two varieties (Table 4) can be attributed to its significantly higher PUE (Table 5). This genetic PUE superiority of Jenguma may, however, be adversely affected if more P than the optimum required by the variety is applied since the higher rate of P in this study led to a significant reduction in PUE (Table 5). The lower PUE observed with the higher P application rate in this investigation is similar to the findings of Khair *et al.* (2002) who reported that P uptake efficiency decreased with the rate of applied P in un-inoculated plots but increased in inoculated plots at NIFA.

Phosphorus uptake efficiency was significantly affected by the interaction of variety and phosphorus rate (Figure 1). Jenguma had a statistically higher mean phosphorus uptake efficiency (41.11%) than Quarshie and Anidaso when phosphorus was applied. Among the interactions tested, the application of P and variety proved to be the best combination with respect to phosphorus uptake efficiency. This observation contradicts the work Daoui *et al.* (2012) who found no significant interaction of genotype and phosphorus (G * P) on P uptake efficiency among Moroccan faba bean varieties.

Table 5: Effects of inoculation rate, variety and phosphorus rate on Amount of N₂ fixed and Phosphorus uptake efficiency of soybean

Treatment	Amount of N ₂ fixed (kg/ha)	P uptake efficiency (%)
<u>Inoculation rate (%)</u>		
0	59.40	16.67
50	59.40	21.11
100	63.60	17.78
P(I)	0.132	0.503
Lsd (0.05)	5.08	10.02
<u>Variety</u>		
Jenguma	59.10	23.70
Quarshie	71.00	13.33
Anidaso	52.33	18.52
P (I)	0.228	0.003
Lsd (0.05)	22.58	5.04
<u>Phosphorus rate (kg P₂O₅/ha)</u>		
0	29.00	-----
22.5	57.30	33.33
45	96.10	22.22
P (P)	<0.001	<0.001
Lsd (0.05)	9.74	2.74
P (I x V)	0.697	0.537
P (I x P)	0.711	0.202
P (V x P)	0.238	<0.001
P (I x V x P)	0.919	0.697
CV (%)	29.00	26.80




Figure 1: Interaction effect of variety and phosphorus application rate on phosphorus uptake efficiency of soybean

CONCLUSION

Inoculation of soybean with the commercial inoculants *Legumefix* at the rates of 2.5 and 5 g/kg seed had no significant effect on plant height, nodulation, shoot dry weight, pod number, pod yield, grain yield, grain size, P uptake efficiency and N₂ fixation. The combined application of P fertilizer and inoculation also had no significant effect on nodulation and BNF of the test soybean varieties. Inoculation of soybean with *Legumefix* on Fluvic luvisol therefore did not favour nodulation and BNF of the three soybean varieties. The soybean varieties were similar in height, nodule formation and development, biomass and grain yields and N₂ fixation but differed significantly with respect to pod load, seed size and phosphorus uptake efficiency in the order Jenguma > Quarshie > Anidaso. The application of phosphorus fertilizer, however, significantly increased all the parameters measured but it was observed that supplying soybean with 45 kg P₂O₅/ha increased growth and grain yield more than applying 22.5 kg P₂O₅/ha though the latter

was was more efficiently used by the crop. It is therefore concluded that P application increases growth and yield of the three soybean varieties and care should be taken to apply the P in the right amount.

REFERENCES

- Alam, M. A., Siddiqua, A., Chowdhury, M. A. and H. Prodhan, M. Y. 2009.** Nodulation, yield and quality of soybean as influenced by integrated nutrient management. Bangladesh Agric. Univ. 7(2): 229–234
- Anderson, J. M and Ingram, J. S. I. 1993.** Tropical soil biology and fertility. A handbook of methods. 2nd Edition pp. 93-95. CAB International U.K
- Bekere, W. and Hailemariam, A. 2012.** Influence of inoculation methods and phosphorus rates on nitrogen fixation attributes and yield of soybean (*Glycine max L.*) at Haru, Western Ethiopia.
- Bekere, W., Endalkachew, W. and Tesfu, K. 2012.** Growth and nodulation response of soybean (*Glycine max L.*) to *Bradyrhizobium* inoculation and phosphorus levels under controlled condition in South Western Ethiopia. African Journal of Agricultural Research Vol.7 (30), pp. 4266-4270.
- Bhuiyan, M. M. H, Rahman, M.M., Afroze, F., Sutradhar, G.N.C. and Bhuiyan, M.S. I. 2008.** Effect of phosphorus, molybdenum and Rhizobium inoculation on growth and nodulation of mungbean. 1Scientific Officer, RARS, Hathazari, Chittagong. J. Soil Nature. 2 (2):25-30
- Bouquet, D. J. 1998.** Yield and Risk Utilizing Short-season Soybean Production in the Mid-Southern USA Crop Sci. 38: 1004-1010.

- 412 **Bray R. H., & Kurtz, L. T. 1945.** Determination of total, organic and available forms of
413 phosphorus in soils. Soil sci., 59: 39-46
- 414 **Bremner, J. M. and Mulvaney, C. S. 1982.** Nitrogen-total, in methods of soil analysis. Page,
415 A. L. (Ed). Methods of soil analysis. Agronomy Monography 9. American Society of
416 Agronomy. Madison, WI. pp. 595-624.
- 417 **Chemining' wa, G. N., Muyhomi, J. W. and Obundho, E. O. 2004.** Effect of rhizobia
418 inoculation and urea application on nodulation and dry matter accumulation of green
419 manure legume at Katumani and Kabeti sites of Kenya. Legume Research Network
420 Project Newsletter Issue No. 11pp 12-18.
- 421 **Date, R., A. 2001.** Advances in inoculant technology: a brief review. Australian Journal of
422 Experimental Agriculture 41, 321-325.
- 423 **Dugje, I. Y., Omoigui, L. O., Ekeleme, F., Bandyopaolhyay, Lava Kumar, R. P. and**
424 **Kamara, A. Y. 2009.** Guide to soybean production in northern Nigeria. IITA, Ibadan. 21
425 pp.
- 426 **F. A. O., 2008.** F. A. O. fertilizer and plant nutrition bulletin. FAO, Rome, Italy, page:220
- 427 **Giller K. E. and Wilson K. J., 1991.** Nitrogen fixation in tropical cropping systems. CAB
428 International, Wallingford, U.K. 313 p.
- 429 **Giller, K. E. 2001.** Nitrogen fixation in tropical cropping system. Wallingford CAB
430 International Wallingford, UK.448p
- 431 **Graham, P. H. 1992.** Stress tolerance in rhizobium and Brady rhizobium and nodulation under
432 adverse soil conditions. Canadian Journal of Microbiology 38, 475-484
- 433 **Hussein, A.H.A. 2009.** Phophorus use efficiency by two varieties of corn at different phosphorus
434 fertilizer application rates. Research journal of applied sciences 4(2):85-93.

- 435 **Khair, M.K.**, Moammed, T., Faisal, H.Tand Khalid, N.2002. Phosphorus use efficiency as
 436 affected by phosphorus application and inoculation. parkistan Journal of Agronomy 1(1)
 437 49-50.
- 438 **Landon, 1991.** Tropical book of soil manual. 2nd edition
- 439 **Larue, T. A. and Patterson, T. G. 1981.** How much N do legumes fix? Advances in agronomy,
 440 34, 15 - 38. In Ogoke, I.J., Togun, A.O.,Carsky, R.J.,and Dashiell, K.E. 2006. N₂ fixation
 441 by soybean in the Nigerian moist savanna: Effects of maturity class and phosphorus
 442 fertilizer. Tropiculture 24, 4193-199.
- 443 **Mahamood, J., Abayomi, Y. A., and Aduloju, M. O. 2009.** Comparative growth and grain
 444 yield responses of soybean genotypes to phosphorous fertilizer application. African
 445 Journal of Biotechnology Vol. 8 (6), pp. 1030-1036. Available online at
 446 <http://www.academicjournals.org/AJB>
- 447 **Malik, M. A., Cheema, M. A., Khan, H. Z., and Ashfaq, M. W. 2006.** Growth and yield
 448 response of soybean (glycine max l.) to seed inoculation and varying phosphorus levels.
 449 J. Agric. Res., 44:47-53
- 450 **Mapfumo, P., Campbell, B.M., and Mpeperek, P. 2001.** Legumes in soil fertility
 451 management: the case of pigeon pea in smallholder farming systems of Zimbabwe.
 452 African crop soil science journal 9: 629-644.
- 453 **Murray U., David H., Mark P., George C., Bob B., Ken G., Bruno A., and Philip C., 2008.**
 454 Measuring plant associated nitrogen fixation in agricultural systems. Australian center for
 455 international agricultural research. www.aciar.gov.au

- 456 **Ogoke, I.J., Togun, A.O., Carsky, R.J., and Dashiell, K.E. 2006.** N₂ fixation by soybean in the
457 Nigerian moist savanna: Effects of maturity class and phosphorus fertilizer. *Tropiculture*
458 24, 4193-199
- 459 **Otieno, P. E., Muthomi, G. N., Chemining Wa, and Nderitu, J. H. 2007.** Effect of rhizobia
460 inoculation, farmyard manure and nitrogen fertilizer on growth, nodulation and yield of
461 selected food grain legumes. *African Crop science Conference Proceedings* vol. 8. Pp.
462 305-312.
- 463 **Plahar, W. A. 2006.** Overview of the soybean industry in Ghana. CSIR-food research institute,
464 P. O. Box m.20, Accra, Ghana. Workshop on soybean protein for human nutrition and
465 health. www.csir.org.gh/fri.html
- 466 **Qiu, J. and Mosjidis, J. A. 1993.** Genotype and planting depth effects on seedling vigor in
467 sericea lespedeza. *Agr. Exp. Station., Auburn University. Pp* 46:309-312
- 468 **Rao, I. M., Borrero, V., Ricaurte, J., Garcia R., and Ayarza, M. A. 1997.** Adaptive attribute
469 of tropical forage species to acid soils. 11 Differences in phosphorus acquisition and
470 utilization as influences by varying supply and soil types. *J. plant nutrition*, 20: 155-180
- 471 **Roughley, R. J., Gemell, L. G., Thompson, J. A. and Brockwell, J. 1993.** The number of
472 *Bradyrhizobium* sp. (*Lupinus*) applied to seed and its effect on rhizosphere colonization,
473 nodulation and yield of *Lupin*. *Soil Biology and Biochemistry* **25**, 1453-1458.
- 474 **Ruark, M. 2009.** Nitrogen and soybeans. assistant professor of soil science state extension Soil
475 Scientist. Area Soil, Water, and Nutrient Management Meetings
- 476 **Salisbury, F. and Ross, C. W. 1992.** Plant physiology. Wadsworth publishing company
477 California, U.S.A. pp540.

- 478 **Sangakkara, U. R., Richner., W., Schnelder, M. K., Stamp, P. 2003.** Impact of intercropping
479 beans (*Phaseolus vulgaris* L.) and sun hemp (*crotalaria juncea* L.) on growth, yields and
480 nitrogen fixation of maize (*Zea mays* L.) grown in the humid and tropics during the minor
481 season. Maydike 48:233-238
- 482 **Shahid, Q. M., Saleem, M. F., Khan, H. Z. and Anjum S. A. 2009.** Performance of soybean
483 (*Glycine max* l.) under different phosphorus levels and inoculation. Department of
484 agronomy, university of agriculture, Faisalabad, Pakistan. Pak. j. agri. sci., vol. 46(4),
485 <http://www.pakjas.com>
- 486 **Singleton, P. M. and Bohlool, B. B. 1984.** Effect of salinity on nodule formation by soybean.
487 Plant and Physiology. 74: 72-76.
- 488 **Solomon, T., Lalit, M. P., & Tsige, A. 2012.** Effects of inoculation by Brady rhizobium strains
489 on nodulation, nitrogen fixation, and yield of soybean (*Glycine max* l., *merill*) varieties
490 on Nitisols of Bako, Western Ethiopia. Department of natural resource management,
491 colleague of agric and env'tal science, Haramaya University. International scholarly
492 research network
- 493 **Walkley, A., and Black, I. A. 1934.** An examination of the degtjareff method for determining
494 soil organic matter and a proposed modification of the chromic and titration method. Soil
495 Sci. 37:29-38.
- 496 **Yusuf, A. A., Iwuafor, E. N. O., Olufajo, O. O, Abaidoo, R. and Sanginga, N. 2006.**
497 Genotype effects of cowpea and soybean on nodulation, N₂-fixation and N balance in the
498 northern Guinea Savanna of Nigeria. In Proceedings of the 31st Annual Conference of
499 the Soil Science of Nigeria 13-17th November, 2006. Ahmadu Bello University Zaria.
500 147-154.