

Disease severity on cabbage protected from arthropod pests with insecticide treated Agronets in Kenya

Abstract

Cabbage (*Brassica oleracea* var. *capitata* L.) is one of the most important vegetables in Kenya, mainly grown by smallholder farmers for food and source of income. Its production is hampered by insect pests and diseases, leading to economic yield losses. Row cover insect proof nets have been used in the past to manage insect pests of cabbage and were recently introduced to Kenya. This study planned to document disease incidence and severity levels on cabbage grown under insecticide impregnated Agronets as a tool to manage insect pests and microclimate. The study was conducted at KALRO-Kabete and PTC (Practical Training Centre)-Thika, Kenya both at the nursery and in the field. Treatments included 1) use of insecticide (alpha cypermethrin) impregnated Agronets with (0.9 mm mesh), 2) untreated 0.4 mm mesh Agronet, 3) grass shading as a farmer practice (at nursery) and 4) control where there was no Agronet use. The treatments had five replications in a completely randomized block design. Plots measured 2 x 6 m, with a 1 m path between consecutive plots and 2 m between consecutive blocks. Three major diseases were recorded: damping off (*Rhizoctonia solani*), downy mildew (*Peronospora parasitica*) and black rot (*Xanthomonas campestris* pv. *campestris*). Treatments showed significant ($P<0.05$) differences on severity of damping off, which was lowest on seedlings covered with Agronets compared with the other treatments at nursery. Comparably, the downy mildew disease severity was higher on seedlings covered with Agronets than those with no cover or shading at the nursery. The findings show that frequent disease scouting is crucial when using Agronets for cabbage pests management. This would allow farmers to take action when diseases are noticed.

Further supportive studies to understand threshold limits that can trigger growers to employ disease control application may be needed.

Key words: cabbage seedling, downy mildew, damping off, black rot, disease incidence

1.0 INTRODUCTION

Cabbage is one of the most important vegetables in Kenya and is grown by smallholder farmers as a major source of income. The crop provides proteins comprising of all essential amino acids, minerals such as calcium, iron, magnesium, sodium, potassium and antioxidant and is reported to have anti-carcinogenic properties (Singh *et al.*, 2009). It is also a rich source of ascorbic acids, carotene (Pro Vitamin A) and has a fiber content and calcium which may reduce the risk of colon cancer (Swarup, 2006). Being a major source of income for smallholder farmers, it has enabled them to support their family needs and remain financially viable, especially in the rapidly growing Kenyan peri-urban farming community. However, its production is hampered by several biotic and abiotic factors. Major constraints in Kenya, and other tropical countries, are insect pests and diseases (MOA, 2006). Among cabbage diseases, the most serious are black rot caused by *Xanthomonas campestris* pv. *campestris*, turnip mosaic virus and soft rot caused by several different pathogens such as *Erwinia carotova*, *Erwinia chrysanthemi* and *Pseudomonas flourescens* (MOA, 2006). Downy mildew (*Peronospora parasitica* (Pers)), is also reported as a key disease, limiting cabbage production (Nyambo and Lohr, 2005). Black rot is usually a destructive disease of a wide range of crucifer plants and cabbage is one of the main hosts. Black rot has been a major disease constraint of cabbage production in smallholder farming systems in Africa where substantial losses are experienced, especially during the warm and wet seasons (Day *et al.*, 1992; Onsando, 1992; Mguni, 1996; Massomo, 2002). It has been reported that under severe outbreak of this disease, the crop is subjected to attack by a variety of pathogens both at

nursery as well as in the field leading to substantial reduction in yield to tune of 5 to 70% (Jorgensen and Walter, 1995). Agronets have been used extensively in various cultivations. They not only contribute to increased production but they also have a positive effect on the quality of the produce by moderating the microclimate under cover. It has also been used to enhance productivity, quality and homogeneity of plants and fruits by generating moderate greenhouse and windbreak effects. They have also been used for protection against adverse climatic action by protecting cultivations from damages due to hail, strong winds and rain (Briassoulis *et al.*, 2007). In Benin, Agronets are used to protect crops against infestation by pests (Martin *et al.*, 2006; Licciardi *et al.*, 2008). In Kenya, it has been tested and proved as an effective pest management tool against various insect pests (Gogo *et al.*, 2012; Muleke *et al.*, 2013). It is thus prudent to understand the effects that this technology may have on diseases of cabbages, which are not targeted for control by the technology. Since the Agronets modify the microclimate, retaining higher moisture and relative humidity, and increased temperatures, there is likelihood of severe disease growth if uncontrolled. This study was carried out to document the key cabbage diseases, their incidence and severity on cabbage grown under the Agronet technology both under nursery and field conditions in Kenya.

2.0 MATERIALS AND METHODS

2.1 Study site: The experiment was conducted at the KALRO-Kabete and PTC (Practical Training Centre) Thika- Kenya experiment stations, from March to September 2011. Kabete station is located Northwest of Nairobi 36⁰ 41'E and 01⁰ 15'S and at an altitude of 1747 m above sea level. The annual mean temperature ranges from 11⁰c to 23⁰c. The average rainfall received is 1000 mm per year and the soil type is Nitisol.

Practical Training Centre-Thika is located at latitude, 1°06'S and longitude, 37° 01'E. The altitude is 1520m. The mean annual rainfall at Thika is 856mm and is bimodal. Mean annual maximum and minimum temperatures are 26.1°C and 13.6°C respectively. The soil type is Pellic Vertisol with an average pH of 5.3 and soil depth ranged from 60 to 100 cm. The two stations were used as a repetition of the study in space.

2.2 Land preparation and transplanting: The land was prepared prior to sowing of the seed on 9th March 2011 at KALRO-Kabete and 15th April 2011 at PTC-Thika. Twenty nursery beds of 1 m by 2 m with an alley of 1 m maintained between plots, and a 2 m between blocks. Cabbage variety Gloria F1, a widely used variety in Kenya was sown on each plot on 21st March and 9th May, 2011, at KALRO-Kabete and PTC-Thika, respectively. One seed was placed per hole at a spacing of 15 cm between lines and 1cm between seeds within a line. Space of 20 cm was maintained from the edge of the plot to the first line. The beds were later covered with dry grass as mulch immediately after planting to prevent wind or water erosion. Nets were supported by metal rod (R6) arches measuring 0.5 m by 0.5 m by 0.5 m (two arches per plot). Nets measuring 2 m by 3 m were placed on the arches and covered on the ground with soil to prevent any opening to the crop. Watering was done manually using watering can, every day in the evening at 6.30 pm for the first two weeks, and thereafter done every three days until the seedling were ready for transplanting. A systemic fungicide, Previcur (propamocarb hydrochloride) was applied as a drench in all plots at the rate of 15 ml in 10 litres of water, for control of fungal diseases (damping off and downy mildew), which are rampant in the area. However, it was only applied after symptoms of damping off were observed on the seedlings and this was done two weeks after germination. Healthy cabbage seedlings from nursery plots protected by nets were transplanted to ploughed, fine tilled main experimental field/plots, on 30th April and 24th June

2011, at KALRO-Kabete and PTC-Thika, respectively. During transplanting, one teaspoonful (50 g) of diammonium phosphate 18:46:0 fertilizer was applied per planting hole and mixed with the soil before placement of the seedling. Watering wa

2.3 Experimental design: In the nursery experiment, treatments included; 1) insecticide (factory impregnated with apha cypermethrine 1%) Agronet with 0.9 mm mesh, 2) untreated 0.4 mm mesh Agronet, 3) grass shading as a farmer practice, and 4) control where there was no Agronet use. In the field, treatment included; 1) insecticide (factory impregnated with alpha cypermethrin 1%) Agronet with 0.9 mm mesh, 2) untreated with 0.9 mm mesh Agronets, and 3) control where there was no Agronet use. The treatments had five replications arranged in a completely randomized block design. Plots measuring 1 x 2 m for nursery and 2 x 6 m for field experiment, with a 1 m path between consecutive plots and 2 m between consecutive blocks were used.

2.4 Data recording: Sampling was done from 9.00 am for 3 hours, on 20 randomly selected plants in each plot. Sampling in the nursery experiment was done within 30 days before transplanting, at weekly interval. For each selected seedling, disease incidence and severity was recorded per plant beginning first week after germination. Disease severity were scored as; 1= up to 20% leaf area affected, 2= >20 to 40% leaf area affected, 3= >40 to 60% leaf area affected, 4= > 60 to 80% leaf area affected, and 5= >80 to 100% leaf area affected.

2.5 Statistical analysis: All the data were subjected to Analysis of variance (ANOVA) using GENSTAT Discovery 4th edition. Significance was tested at 95% confidence limit and if found significant means were separated using Students-Newman Keuls (SNK) test. The means are provided with standard error (SE). Data that were not normally distributed were transformed using square root method $(x+0.05)^{0.5}$ before performing ANOVA.

114 3.0 RESULTS

115 Diseases observed in nursery included; downy mildew, damping off and black rot. Downy
 116 mildew was most severe at the nursery compared with the field. The first symptoms observed
 117 were small, light green-yellow lesions on the upper leaf surface, then later showing on the
 118 undersurface. The spots turned yellow as they enlarged. Downy mildew severity was
 119 significantly ($P<0.05$) higher on seedlings covered by Agronets compared with those grown
 120 under grass shade and those with no cover (control) at both Kabete and Thika. However, for
 121 damping off, there was no significant ($P>0.05$) difference across treatments at Kabete (Table 1)
 122 although in Thika, the severity was significantly lower ($P<0.05$) on those seedlings covered by
 123 Agronets compared with those shaded with grass and control. There was no significant
 124 difference between Agronets despite the difference of mesh size (0.9 mm and 0.4 mm)

125 Table 1: Mean disease severity on cabbage seedlings at nursery grown under different
 126 (treatments): treated net, untreated net, grass shading and control; at KALRO Kabete and
 127 KALRO Thika, March and June 2011, respectively.

Treatment	Damping off		Downy mildew	
	Kabete	Thika	Kabete	Thika
Treated net	0.27	0.53a	2.07a	2.01a
Untreated net	0.33	0.53a	2.13a	2.07a
Grass shading	0.53	1.00b	0.60b	0.60b
No net	0.86	1.47b	0.73b	0.73b

P-Value	0.279	0.028	0.002	0.020
SED	0.142	0.349	0.259	0.249

Mean in the same column followed by the same letter are not significantly different by Students-Newman Keuls (SNK) at $P = 0.05$

The disease incidences on cabbage seedlings both at Kabete and Thika did not vary significantly ($P < 0.05$) across the treatments. However, lower incidences of damping off disease were observed on plots covered by Agronets and those grown under grass shading compared with the control. Comparably higher incidences of downy mildew were observed on seedlings covered by Agronets compared with uncovered seedlings at Kabete. In Thika downy mildew did not differ significantly ($P > 0.05$) across the treatments (Table 2).

Table 2: Mean disease incidence on cabbage seedlings at nursery grown under different (treatments): treated net, untreated net, grass shading and control; at KALRO Kabete and KALRO Thika, March and June 2011, respectively.

Treatment	Damping off		Downy mildew	
	Kabete	Thika	Kabete	Thika
Treated net	4.33	4.33	56.33	25.00
Untreated net	5.00	4.53	51.33	29.33
Grass shading	0.01	7.33	32.67	24.00
No net	8.33	8.00	20.33	26.67

P-Value	0.071	0.871	0.111	0.954
SED	0.629	0.822	1.401	1.599

At Kabete, severity of downy mildew was greater the 2nd week after germination, this was observed across the treatments. However, higher severity was observed in seedlings covered by Agronets compared with the uncovered seedlings and those grown under grass shading. The severity later reduced on the 3rd week after a control measure had been applied (Figure 1).




Figure 1 Downy mildew severity on cabbage seedlings at Kabete

At Kabete, damping off severity was significantly lower on seedlings covered by Agronets compared with those grown under grass shading and uncovered seedlings. The severity increased towards the end of the 1st week, and highest severity was recorded in the 2nd week. A control measure was applied and the severity reduced from the 2nd week subsequently until the crop was ready for transplanting (Figure 2).




Figure 2 Damping off severity on cabbage seedlings at Kabete

At Thika, damping off disease severity was significantly lower in seedlings covered by Agronets compared with uncovered seedling and those grown under grass shading. Severity was highest in the 2nd week and later decreased towards the end of the 2nd week and the 3rd week. This was due to the control measure applied (Figure 3).




Figure 3 Damping off severity on cabbage seedlings at Thika

At Thika, a similar trend was observed in downy mildew severity like the one at Kabete. The severity was greater in the 2nd week of transplanting across the treatments. A control measure was applied and the severity reduced to a lower level towards the end of 2nd week until the seedlings were ready for transplanting (Figure 3.a).




Figure 3.a Downy mildew severity on cabbage seedlings at Thika

After planting out, downy mildew severity was significantly ($P < 0.05$) higher on cabbages covered by Agronets compared with uncovered cabbages, this was observed in Thika. At Kabete downy mildew severity did not vary significantly ($P > 0.05$) across the treatments, however, severity was relatively higher in cabbages covered by nets. At Thika, damping off diseases attack on cabbages were significantly lower ($P < 0.05$) in cabbages covered by nets compared with uncovered cabbages. At Kabete, the damping off severity did not vary significantly ($P > 0.05$) across the treatments. However, general trend showed relatively lower severity on covered cabbages compared with uncovered plants. Black rot was significantly lower ($P < 0.05$) on covered plants compared with uncovered cabbages both at Kabete and Thika (Table 3).

174 Table 3: Mean diseases severity on field cabbage for the whole season at KALRO Kabete and
175 Thika, April-July and July-October 2011, respectively.

Treatment	Damping off		Downy mildew		Black rot	
	Kabete	Thika	Kabete	Thika	Kabete	Thika
Treated net	0.13	0.13	0.38	0.42a	0.13a	0.12a
Untreated net	0.12	0.15	0.38	0.42a	0.13a	0.17a
No net (control)	0.20	0.40	0.30	0.13b	0.55b	0.38b
P-Value	0.74	0.05	0.25	0.04	0.01	0.03
SED	0.08	0.06	0.09	0.09	0.08	0.07

176 Mean in the same column followed by the same letter are not significantly different by Students-
177 Newman Keuls (SNK) at P =0.05

178 Damping off disease in field cabbage at Thika was observed the 1st week after transplanting this
179 was observed across the treatment (Figure 4). Greater severity was recorded in plots with no
180 covers. However, after application of a control measure the effect was minimized.




Figure 4 Severity trend of damping off disease on cabbage at Thika

Black rot severity was observed across the treatments, with higher severity recorded on plots with no cover (Figure 5). This was in comparison with those plants covered by Agronets, which were relatively lower in severity. A control measure was applied in all the treatments before the 4th week that reduced their severity, which remained low until harvesting. However, uncovered plants were later attacked from the 6th week, and severity increased progressively throughout the remaining period of the season.





Figure 5 Black rot disease incidences on cabbage crop in field trial at Kabete

191 Black rot symptoms were observed in the 2nd week after transplanting in Thika. Greater severity
 192 was recorded on uncovered cabbages, in the same week a control measure was applied across the
 193 treatments and the infection was minimized. However uncovered plants showed higher severity
 194 compared with covered plants from the 5th week until the crop matured (Figure 6)



195
 196 Figure 6 Severity of black rot in field cabbage at Thika

197 4.0 DISCUSSION

198 Downy mildew severity was greater on seedlings covered by Agronets due to the shade effects of
 199 the net that could have increased moisture content inside the net and raised temperatures, both
 200 factors enabling disease development in a crop. This is in consistent with other findings in Kenya
 201 by Gogo *et al.* (2012), there was 15% increase in mean daily temperature and relative humidity
 202 under the netting treatment due to reduced air movements, evapotranspiration and shading effect
 203 compared with the control. In related studies (Petzoldt and Seaman, 2006), some pathogens such
 204 as apple scab, late blight, and several vegetable root pathogens were reported to be more likely to
 205 infect plants with increased moisture. Other pathogens like powdery mildew species tend to
 206 thrive in conditions with lower (but not low) moisture. Moreover, higher growth rates of leaves
 207 and stems observed for plants grown under high carbon dioxide concentrations results in greater

208 fungal spore production. The high severity was also enhanced by the delayed control measure
 209 applied on the growing seedlings; this was observed at KALRO-Kabete. This study shows that
 210 users of Agronets for cabbage production should keenly monitor for disease presence so that
 211 control options can commence before the severity is too high to cause economic damage.
 212 Damping off disease was significantly lower on seedlings covered by Agronets compared with
 213 those with no cover. This was probably due to the mode of disease spread. Being a soil borne
 214 pathogen, the Agronet could have acted by reducing water flow since irrigation was applied due
 215 to better moisture retention. In addition, there was controlled runoff on plots covered with
 216 Agronets since the rain drops took more time to sip down to the soil as a result of small mesh
 217 size. Therefore, covering of plants with Agronets can also provide an opportunity to include
 218 integrated disease management practices to protect cabbages both in nursery and field. In related
 219 studies by Singh (2002); Bal (2006), the effects of shelterbelt (hessian cloth) on the development
 220 of bacterial canker (*Xanthomonas axonopodis pv. citrii*) and fungal scab complex (*Elsinoe*
 221 *fawcetti*) in citrus nursery, reportedly provided protection by acting against wind-driven rains.
 222 Moreover, it has been reported that nylon net can be a useful cultural practice for protection of
 223 cauliflower seedlings against Alternaria blight (*Alternaria brassicicola*) and black rot at least
 224 when they are still in the nursery bed (Kashyap and Dhiman, 2010). Black rot is a seed-borne
 225 disease (Neergaard, 1945), though it attacks already growing crop. The spread of black rot in
 226 field cabbage was minimized by covering plants with Agronet, since there were reduced rain
 227 splashes from infected plants unlike uncovered plants. According to Carissie *et al.*, (1997), the
 228 dispersal of black rot pathogen is due to the combined effect of flow of contaminated water
 229 droplets, rain splashes and windy conditions. If diseases are well managed, Agronet technology
 230 could provide sustainable production of cabbages and other crops with little or no pesticide

application in a season. This would be encouraging considering the high amounts of pesticides consumed in protection of cabbages against arthropod pests in Kenya. For growers, even though the technology may be useful, it should be taken as component of integrated pest and disease management system where monitoring/scouting is critical. This could assist in identifying the best low cost ways of reducing infection at the onset of a disease and thus avoid calendar or frequent use of fungicides in cabbages.

5.0 CONCLUSION

The results from this study show that Agronets should not expected to control cabbage diseases though the problem may be minimized by use of Agronets. Thus frequent monitoring and use of monitored fungicides at onset of the diseases should form part of integrated pest and disease management on cabbages when Agronets are used. This will prevent disease development to uneconomic levels.

6.0 RECOMMENDATIONS

1. Field monitoring is important for early detection of disease incidence and for early application of control measures of the diseases especially when growing nursery cabbage seedlings under Agronets.
2. Tests should be done to determine action threshold limits of the diseases and mode of disease spread on Agronet protected cabbages.
3. Farmers should be encouraged to use systemic fungicide at transplanting as it will reduce the effects of soil borne and seedling diseases.

7.0 Disclaimer: - This manuscript was presented in the conference “34 th 13 KARI Biennial Scientific Conference BOOK OF ABSTRACTS 35 th 13 KARI

**Biennial Scientific Conference SESSION 3B: KENYA POLLINATION
INFORMATION NETWORK ...”** available link is “
[http://www.abstract.xlibx.com/a-economy/6789-20-34-kari-biennial-scientific-
conference-book-abstracts-kari-bie.php](http://www.abstract.xlibx.com/a-economy/6789-20-34-kari-biennial-scientific-conference-book-abstracts-kari-bie.php)”

8.0 REFERENCE

- Bal, R. S. and Dhiman, J.S. (2006). Suppression of development of canker scab complex in
Kinnow nursery through cultural practices. *Journal of Reskin Punjab Agriculture
University* 34: 21-24.
- Briassoulis, D., Mistriotis, A. and Eleftherakis, D. (2007). Mechanical behavior and properties of
agricultural nets-Part I: Testing methods for agricultural nets. *Polymer testing* 26: 822-
832
- Carissie, O., Wellman-Desbiens, E., Toussaint, V. and Therese, O. (1997). Preventing black rot.
Agricultural and Agri-food, Canada.
- Day, L. A., Feldman, S., Minja, R. and Wien, C. (1992). Vegetable production in Tanzania, a
case study in Arumeru District. Special Survey Report, Cornell University, United
States of America and Sokoine University of Agriculture, Tanzania. 46pp
- Gogo, O. E., Saidi, M., Itulya, M. F., Martin, T., Ngouajio, M. (2012). Microclimate
modification using eco-friendly nets for high quality tomato transplant production by
small-scale farmers in East Africa. *Horticulture Technology* 22: 292-298
- Jorgensen, M.C. and Walter J.M. (1955).The 1953 Outbreak of Black rot at Ruskin. *Proceedings
of Florida State Horticultural Society* 67: 109-111

- 275 Kashyap, P.L. and Dhiman J.S. (2010). Eco-friendly strategies to suppress the development of
276 *Alternaria* blight and black rot of cauliflower. *Academic Journal of plant Sciences* 3: 140-
277 146
- 278 Licciardi, S., Assogba-Komlan, F., Sidick, I., Chandre, F., Hougard, J.M., Martin, T. (2008). A
279 temporary tunnel screen as an eco-friendly method for small-scale growers to protect
280 cabbage crop in Benin. *International Journal of Tropical Science* 27: 152-153
- 281 Martin, T., Assogba-Komlan, F., Houndete, T., Hougard, J. M. and Chandre, F. (2006). Efficacy
282 of mosquito netting for sustainable small holders' cabbage production in Africa. *Journal*
283 *of Economic Entomology* 99: 450-454
- 284 Massomo, S.M. (2002). Black rot of Cabbage in Tanzania: Characterization of *Xanthomonas*
285 *campestris* pv. *campestris* and Disease Management Strategies. PhD thesis. The Royal
286 Veterinary and Agricultural University, Copenhagen. Denmark. 215 pp
- 287 Mguni, C.M. (1996). Bacterial black rot (*Xanthomonas campestris* pv. *campestris*) of vegetable
288 *Brassicas* in Zimbabwe. The Royal Veterinary and Agricultural University, Copenhagen.
289 Denmark 144 pp.
- 290 Ministry of Agriculture, (2006). Annual Report of year 2005, Nairobi-Kenya.
- 291 Muleke, E.M., Saidi, M., Itulya, F. M., Martin, T. and Ngouajio, M. (2013). The Assessment of
292 the Use of Eco-friendly Nets to Ensure Sustainable Cabbage Seedling Production in
293 Africa. *Agronomy*, 3: 1-12
- 294 Neergaard, P. (1945). Danish species of *Alternaria* and *Stephylium*. *Oxford University Press*.
295 London.
- 296 Nyambo, B. and Lohr, B. (2005). The Role and Significance of Farmers Participation in
297 Biocontrol-based IPM for *Brassica* crops in East Africa. In Second International

298 Symposium on Biological control of Arthropods, Davos, Switzerland (compiled by
299 Hoddle, M. S). *USDA Forest Service Publication FHTET*

300 Onsando, J. M. (1992). Black rot of crucifers. In: Chaube H. S., Sign U. S., Mukhopadyay, A.
301 N. and Kumar. Disease of Vegetable and Oil Seed Crops. *Journal of Plant Disease of*
302 *International Importance*. Prentice Hall, Englewood Cliffs, New Jersey. 243-252 pp

303 Petzoldt, C. and Seaman, A. (2006). Climate Change Effects on Insects and Pathogens, p. III 1–
304 16. In Wolfe, D. W., Grubinger, V., Burtis, B. (eds). *Climate Change and Agriculture:*
305 *Promoting Practical and Profitable Responses*. Cornell University, Ithaca, NY. 120 pp

306 Singh B.K., Sharma S.R and Singh B. (2009).Heterosis for mineral elements in single cross-
307 hybrids of cabbage. *Scientia Horticulture* 122: 32-36.

308 Singh R. (2002). Dynamics of canker-scab complex in kinnow nursery and its management.
309 M.S.c thesis, Punjab Agricultural University, Ludhiana. 68 pp

310 Swarup V. (2006). Vegetable Science and Technology in India. *Kalyani Publishers*, 350pp
311
312
313