

Original Research Article

Field performance and differential response of micro-propagated potential F₁ genotypes of *Gerbera jamesonii*

ABSTRACT

Aims: To evaluate gerbera hybrids with respect to mass propagation under *in vitro* conditions and their characterization under field conditions for floral traits for selection of promising gerbera lines for commercial utilization.

Study design: Field performance and differential response of micro-propagated gerbera were evaluated in randomized block design.

Place and Duration of Study: CSIR-IHBT, Palampur (HP), India 2012-2015.

Methodology: F₁ genotypes of gerbera were developed through controlled crossing program and subjected to *in vitro* micro-propagation to achieve quick multiplication. Mature seeds obtained from different cross combinations were used for the establishment of *in vitro* gerbera cultures. Seeds were cultured on Murashige & Skoog (MS) basal medium and the developing micro-shoots from seeds were further cultured on MS media supplemented with different doses of growth regulators to achieve shoot proliferation.

Results: Highest number of micro-shoots, number of leaves and leaf length were observed in M3 proliferation medium (MS supplemented with 1mg/l 6- Benzylaminopurine + 0.030 mg/l Indole Butyric Acid (IBA) + 0.025 mg/l Naphthalene Acetic acid) which gave best proliferation among the gerbera genotypes. The half strength MS medium supplemented with 0.4mg/l IBA was found best for *in vitro* rooting. Also, there was differential response of hybrid gerbera genotypes to micro-shoot proliferation.

Conclusion: Characterization of two hybrid F₁ genotypes (23-1 and E-3) under field conditions with respect to floral traits in comparison to parental genotypes provided information about new gerbera lines with high micro-propagation potential for commercial utilization.

Keywords: *Gerbera jamesonii*, *in vitro* proliferation, root initiation, 6- Benzylaminopurine, IBA, NAA

1. INTRODUCTION

Gerbera (*Gerbera jamesonii*) is an important commercial cut flower of the trade having extensive demand throughout the world due to its numerous colors, shapes and prolonged vase life. The plant is ideal for beds, borders, pots and rock gardens. It has a great export potential and ranks fifth among top cut flowers in the international market [1]. There is an increasing demand of quality gerbera cut flowers, planting material and new types of varieties for export as well as in domestic markets. *Gerbera* is out crossing in nature and the possibility of genotype fixation by clonal propagation unfolds a large potential for novel cultivar development. New cultivars of gerbera are developed by hybridization of diverse types to achieve genotypic recombination, which requires information about the genetic diversity among the parental genotypes, especially in order to create novel genotypes based on combination of desirable traits that can be easily fixed by vegetative propagation. In this context, with an aim of varietal development, selective breeding of gerbera (for desirable flower colors and complete double types) is being done under an ongoing gerbera breeding program at CSIR-Institute of Himalayan Bioresource Technology, Palampur, Himachal

Pradesh, India [2,3]. The parents were selected based on prior information with respect to variations for flower color [4], shape (double, semi-double and single), size, and good performance under protected cultivation.

Although conventional vegetative propagation of gerbera, by division of suckers is comparatively easy to perform, only limited plants can be developed from one plant and it takes 10–12 months to attain the next cycle of division [5]. Floriculture industry requires about nine plants of gerbera per square meter for clonal plantation under protected cultivation and a huge number of clonal plants are required every year when gerbera cultivation is to be done on commercial scale. Under such a situation it is difficult to produce large number of clonal plants of a promising genotype within a short period of time through vegetative propagation.

In order to meet the industrial demand, we need micro-propagation for rapid and mass multiplication of elite gerbera genotypes under in vitro conditions. However, response of different genotypes and the genetic control of component traits constitute crucial factors for the success of mass propagation program. At least two component traits, viz., proliferation rate and in vitro root induction possibly influence the response of a genotype to mass propagation. Therefore, the present study was undertaken to evaluate gerbera hybrids with respect to mass propagation under in vitro conditions and their characterization under field conditions for floral traits in comparison to parental genotypes for selection of promising gerbera lines for commercial utilization.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

To evaluate response of F_1 gerbera genotypes to micro-propagation, seeds obtained through the crossing program were used as explants for establishing cultures for in vitro clonal multiplication. Twelve F_1 genotypes (23-1, 24-1, 24-2, 24-3, 24-4, A-6, B-2, D-3, E-1, E-2, E-3 and Y-1) generated in this study were evaluated for in vitro response and characterized for floral characteristics under poly-house conditions in the present study (Table 1).

Table 1. List of gerbera genotypes involved in the study

S.No.	Genotypes	Remarks	Shape	Colour
1.	IHBT-Gr-01 (P1)	Parent	Single	White (RHS 155 C)
2.	IHBT-Gr-02 (P2)	Parent	Semi-Double	Red-Purple (RHS 58 D)
3.	IHBT-Gr-03 (P3)	Parent	Double	Yellow-Orange(RHS158 C)
4.	IHBT-Gr-04 (P4)	Parent	Semi-Double	Orange-Red (RHS 40 A)
5.	IHBT-Gr-05 (P5)	Parent	Single	Red-Purple (RHS 66 B)
6.	IHBT-Gr-06 (P6)	Germplasm	Single	Red-Purple (RHS 58 D)
7.	IHBT-Gr-07 (P7)	Parent	Single	Red (RHS 52 C)
8.	23-1 (P2 x P3)	Hybrid F_1	Double	Yellow-Orange (RHS 16 C)
9.	24-1 (P3 x P4)	Hybrid F_1	Semi-Double	Orange

69	10.	24-2 (P3 x P4)	Hybrid F1	Single	Orange-Red
70	11.	24-3 (P3 x P4)	Hybrid F1	Double	Orange-Red
71	12.	24-4 (P3 x P4)	Hybrid F1	Semi-Double	Yellow-Orange
72	13.	Y1 (P3 x P7)	Hybrid F1*	Double	Yellow
73	14.	E-1 (P1 x P7)	Hybrid F1*	Double	White
74	15.	E-2 (P1 x P7)	Hybrid F1	Single	White
75	16.	E-3 (P1 x P7)	Hybrid F1	Double	White (RHS 155 D)
76	17.	A-6 (E1 x Y1)	Hybrid F1	Double	Yellow
77	18.	B-2 (P1 x P5)	Hybrid F1	Single	Red
78	19.	D-3 (P4 x P2)	Hybrid F1	Semi-Double	Orange-Red

79 *used as parents in breeding program

80 Seeds were washed with few drops of Tween-20 to remove the surface dust particles with
81 constant shaking for 5-6 minutes and washed thoroughly with distilled water (6-7 times). The
82 seeds were surface sterilized with 0.1% mercuric chloride solution for 2 minutes under
83 aseptic condition in laminar air flow cabinet and rinsed 4-5 times with autoclave distilled
84 water. The seeds were cultured aseptically on basal MS medium [6]. The pH of the medium
85 was adjusted to 5.8 with 1N NaOH and agar (8g/l) was used for solidification of medium. The
86 media were dispensed into tissue culture flask and each contained 100ml of culture medium.
87 The flasks were autoclaved at 121°C at 15 psi for 20 minutes and then allowed to cool at
88 room temperature before culturing the seeds. Cultures were given a 16 h photoperiod with
89 light intensity of 37.40 mmol-2 s⁻¹ which was maintained by using photosynthetically active
90 radiation (PAR) light sources and fluorescent tubes. Temperature in the culture room was
91 maintained at 25 ± 2°C.

92 After germination the seedlings were transferred to shoot proliferation medium
93 supplemented with 6- Benzylaminopurine at higher levels (BAP - 1.0 mg/l) than Indole 3-
94 butyric acid (IBA - 0.030 mg/l) and Naphthalene acetic acid (NAA - 0.025 mg/l) along with
95 30g/l sucrose for initiating micro-shoots in hybrid gerbera genotypes. When multiple shoots
96 are formed in the cultures, individual shoots are separated using fine scalpel under laminar
97 air flow and subcultures done at regular intervals of four weeks on shoot proliferation
98 medium to attain rapid in vitro multiplication. To study the effect of different concentrations of
99 BAP (0.5 (M1), 0.75 (M2), 1.0 (M3), 1.5 (M4) mg/l) and fixed concentrations of IBA (0.030
100 mg/l) and NAA (0.025 mg/l) an experiment was laid out employing completely randomized
101 design with five explants per flask per treatment and replicated thrice (5 explants x 4
102 treatments x 3 replications = 60 explants per genotype). Data were recorded for number of
103 micro-shoots per plant, leaf number and leaf length in proliferation medium.

104 To initiate rooting, shoots were transferred to rooting medium with half strength of MS salts,
105 supplemented with varying concentration of IBA (0.1, 0.2 and 0.4 mg/l), 15 g/l sucrose and
106 6g/l agar were used. In case of rooting medium, days taken for root initiation were recorded.
107 Rooted plantlets of gerbera were removed from culture flasks and washed thoroughly with
108 running tap water to remove agar. The plantlets were then transferred to plastic trays

109 containing sand and covered with plastic bag to maintain humidity. After 3 weeks of
110 acclimatization they were transplanted in sleeves for further cultivation in soil.

111 Data were analyzed using F-test to evaluate the response of different hybrid gerbera
112 genotypes under proliferation medium. The data regarding different proliferation media and
113 rooting media were analyzed using Student's t-test. All tissue culture experiments were laid
114 out in a Randomized Block Design and the significance of differences between means was
115 estimated at 5% level of significance.

116 In the field experiment conducted during 2013-14 and 2014-15, two hybrid genotypes of
117 gerbera 23-1 and E-3 which performed well under in vitro conditions were evaluated along
118 with parental genotypes IHBT-Gr1, IHBT-Gr2, IHBT-Gr3 and IHBT-Gr7. All the six genotypes
119 were planted polyhouse in randomized block design with four replications and data were
120 recorded for different agro-morphological and floral traits viz., leaf number per plant, plant
121 spread (cm), number of suckers per plant, leaf length (cm), leaf width (cm), flower color,
122 flower shape, disc color, flower number per plant, scape length (cm) and flower diameter
123 (cm). Data were collected on five flowers of each genotype in each replication. All the data
124 were recorded at the commercial stage of flower harvest; the stage at which the stigmata of
125 the ray florets have been exerted [7]. Flower color of selected gerbera genotypes along with
126 parental lines was compared using the Royal Horticultural Society's (RHS) color chart [8].
127 Flower shape (single - with whorls of only ray florets, semi-double - with separate whorls of
128 ray florets and smaller-sized transition florets and complete double with ray florets covering
129 the entire disc) [9] and disc color (light green / dark brown) were observed on visual basis.
130 Data with respect to quantitative traits leaf number per plant, plant spread (cm), number of
131 suckers per plant and flower number per plant were recorded on yearly basis on five random
132 but competitive plants of each genotype while leaf length (cm), leaf width (cm), scape length
133 (cm) and flower diameter (cm) were recorded on five mature leaves and five flowers of each
134 genotype. The data over two years was studied separately to assess overall performance of
135 the gerbera genotypes. Analysis of variance was conducted to work out significance of
136 variability for different agro-morphological and floral traits.

137

138 3. RESULTS AND DISCUSSION

139

140 In the tissue culture experiments, initially the seedlings raised on MS basal medium of twelve
141 hybrid genotypes viz. 23-1, 24-1, 24-2, 24-3, 24-4, A-6, B-2,D-3,E-1, E-2, E-3 and Y-1 when
142 cultured on proliferation medium supplemented with BAP (1.0 mg/l) + IBA (0.030 mg/l) and
143 NAA (0.025 mg/l) gave a differential response with respect to micro-shoot formation during
144 first three sub-cultures. Of these, only five genotypes (23-1, 24-4, A-6, E-3 and D-3)
145 responded well in the proliferation medium and taken forward for experimentation to
146 evaluate their proliferation potential under varying concentrations of BAP and different
147 concentrations of IBA for rooting (Plate 1). Since sufficient micro-shoot proliferation in other
148 genotypes was not achieved to the same extent in the first three sub-cultures, they were
149 excluded from the experiment so as to homogenize the experimental data set of 60 explants
150 per genotype. The data on effect of concentrations of BAP on micro-shoot multiplication
151 recorded after one month of culture is given in Table 2. The maximum number of micro-
152 shoots (12.73), leaves (44.7) and leaf length (3.48 cm) were produced in the medium M3
153 containing 1 mg/l BAP+ 0.030 mg/l IBA+ 0.025mg/l NAA followed by M2 medium with
154 0.75mg/l BAP+ 0.030 mg/l IBA+ 0.025mg/l NAA.



155

156 Plate 1. Differential response of different gerbera genotypes to in vitro proliferation

157 Table: 2. Effect of BAP on number of micro-shoots, number of leaves and largest leaf length in five different genotypes of gerbera

158	Media	Number of micro-shoots per explant						Number of leaves per explant						Largest leaf length (cm)					
159		E-3	23-1	24-4	D-3	A-6	Mean	E-3	23-1	24-4	D-3	A-6	Mean	E-3	23-1	24-4	D-3	A-6	Mean
160	M1	9.0	10.3	5.0	5.4	4.8	6.9	31.7	31.8	21.5	23.9	18.9	25.5	2.1	2.6	2.2	2.9	2.1	2.4
161	M2	10.4	10.6	5.2	5.7	5.1	7.4	32.8	33.3	23.4	25.1	21.1	27.1	2.1	2.7	2.4	3.0	2.3	2.5
162	M3	11.6	12.7	5.6	5.9	5.4	8.2	44.7	41.5	30.0	32.7	22.8	34.4	2.2	2.9	2.3	3.4	2.2	2.6
163	M4	11.2	11.5	5.1	5.6	4.9	7.6	34.2	34.4	22.4	28.9	19.5	27.8	2.2	2.8	2.2	3.2	2.2	2.5
164	Mean	10.5	11.3	5.2	5.6	5.0		35.8	35.2	24.3	27.6	20.5		2.2	2.8	2.3	3.1	2.2	

165 Currently, *in vitro* propagation is the preferred method for quick multiplication of the planting
166 material [10, 11, 12, 13, 14]. Gerbera plants have been regenerated from calli derived from a
167 variety of explants [15, 16, 17, 18, 19]. Adventitious shoots have also been regenerated from
168 the flower buds of Gerbera plants grown in the greenhouse [20, 21]. However, in this study,
169 seeds were available which were developed through crossing under an ongoing gerbera
170 breeding program and used for *in vitro* propagation. Accordingly, standardization of culture
171 medium with optimum concentration of growth regulators is critical for evaluating micro-
172 propagation response of the different genotypes involved in the study.

173 To further understand the response of potential gerbera genotypes, data generated as per
174 randomized block design in tissue culture experiment were analyzed and maximum micro-
175 shoot formation was observed in 23-1 followed by E-3 (Table 2), whereas more leaves were
176 formed in E-3 as compared to 23-1 (Plate 2). Analysis of variance of the shoot proliferation
177 data indicates significant genotypic variations (Table 3). Overall, among the five gerbera
178 genotypes evaluated for proliferation potential, 23-1 and E-3 have significantly better
179 response than other genotypes under *in vitro* conditions. The present investigation suggests
180 that number of micro-shoots leaf number and leaf size are independent traits which are
181 affected by media conditions and contribute to overall shoot proliferation in gerberas.
182 Considerable variations were obtained for shoot proliferation parameters indicating
183 genotypic specificity of gerbera accessions to these component traits of *in vitro* propagation.
184 Genotypic response to *in vitro* propagation has also been reported in other plant species
185 such as sugarcane [22] and *Phoenix dactylifera* [23].



186
187 Plate 2. Growth of gerbera genotypes 23-1 and E-3 (left and right, respectively) after one
188 month in proliferation medium

189 Table 3 ANOVA test for the differences in micro-shoot formation per explant, number of
190 leaves per explant and leaf size among five different genotypes of gerbera

191

192	S. No.	Source of variaiton	Variance	F- value
193	1.	micro-shoot formation per explant	38.22	136.12*
194	2.	number of leaves per explant	180.27	53.06*
195	3.	leaf size (cm)	0.70	58.13*

Based on Student's t-test, media M3 and RM3 were significantly better from the other media for shoot proliferation and root induction in gerbera. Different combinations of cytokinin and auxins have been used to achieve multiple shoot induction in Gerbera [24, 25, 26]. In the present studies, among the different concentrations of BAP in combination with IBA and NAA, the MS medium supplemented with 1 mg/l BAP was found to be best medium as compared to others because it results in maximum number of shoots and shoot length, whereas high concentration of BAP (1.5 mg/l) induced negative effect on plant growth (Figure 1). These results are in conformity with that of [27] who obtained highest shoot number in gerbera with 1mg/l BAP. Highest multiplication rates [28] were observed in gerbera with 3 mg/l BAP which is contrary to the present finding. Genotypic specificity to growth regulators may possibly be the reason for such large variations in response to concentrations of growth regulators. The gerbera genotypes which did not respond well under initial tissue culture experiment need to be evaluated for higher doses of BAP so as to ascertain their growth regulator requirements for effective micro-propagation. There are reports [29, 30] that the cultivars varied widely in their growth regulator requirements at multiplication phase. In the present study, among different genotypes evaluated for micro-propagation, genotypes 23-1 and E-3 were observed best because micro-propagation rate was found to be high in these two genotypes.





Figure 1 Comparison of proliferation media (M1-M4) for micro-shoot formation per explant, number of leaves per explant and leaf size (cm) in gerbera

Rooting was obtained in all the root induction media, but the main difference was time taken for root initiation. Earliest response for root initiation was observed in RM3 medium (Figure 2). IBA plays important role in root induction in tissue culture plants. IBA has been reported as the best medium for root induction in Gerbera [19]. The root initiation and root development were better in MS medium with higher concentration of IBA as compared to lower concentration in all the five varieties used in the present study. 100% rooting in red yellow and white varieties was obtained the cultured on half MS containing 0.2 mg/l IBA within 15-20 days of root initiation [31]. Successful rooting was obtained from *in vitro* cultured gerbera shoots in MS medium containing 1.5mg/l IBA [27]. The results obtained in our study confirm the earlier reports. However, response to root initiation in rooting media was observed to be at par among the gerbera genotypes. A large number of rooted plantlets of gerbera were successfully hardened by removing them from culture flasks, washed thoroughly under running tap water to remove agar and then transferred to plastic trays containing sand and covered with plastic bag to maintain humidity. After 2-3 weeks of acclimatization they were transplanted in sleeves for further cultivation in soil (Plate 3). The new gerbera genotypes were characterized with respect to floral traits and showed high to moderate growth response to *in vitro* proliferation as compared to other gerbera genotypes and have good potential for commercialization (Plate 4).



235

236 Figure 2 Mean differences of rooting media (RM1-RM3) for days to root initiation in gerbera



237

238 Plate 3. Root initiation in gerbera cultures, hardening of rooted plantlets in trays and transfer
239 of plantlets in sleeves in hardening chamber



240

241 Plate 4. Potential gerbera genotypes 23-1 (left) and E-3 (right) with good field performance
242 and high micro-propagation rate

243 The floral description and agro-morphological performance of genotypes is mentioned in
244 Tables 1 and 4. Significant variability was observed among the F1genotypes 23-1 and E-3
245 as compared to parental lines Gr1, Gr2, Gr3 and Gr7 for the agro-morphological and floral
246 traits. The extent of variability based on analysis of variance estimates for different traits
247 (Table 5) suggests that there is wide scope for improvement of agro-morphological as well
248 as floral traits in gerbera through a breeding program. Data generated for quantitative traits
249 over two years were analyzed separately and not pooled because *Gerbera jamesonii* is a

perennial plant species and the observations made were not repetition but continuation of the same experiment. Significant genotypic variations were recorded for the agromorphological and floral traits although variations among the replicates were not significant which is expected considering that the plants of each genotype have clonal origin and the experiment was undertaken under controlled poly-house conditions. Based on mean performance of hybrid gerbera genotypes with the respective parents, E-3 was superior to at least one parental genotype for traits plant spread in a year, leaf length (cm) and leaf width (cm), while it was at par to parental genotypes for the other traits. In case of 23-1, the genotype was superior to at least one parental genotype for leaf width (cm), number of flowers per plant in a year and scape length (cm). Comparison of data for the two years suggests that some of the traits tend to stabilize over the years (leaf number per plant/year, leaf length, leaf width, flower number per plant/year and flower diameter), while others become more variable (plant spread, number of suckers per plant/year and scape length). As is evident from the data, characters such as plant spread, number of suckers per plant/year and scape length are associated with overall growth of the plants and may not completely express during first year of plant growth.

Table 4. Mean performance of gerbera genotypes over two years

S. No.	Source of variation	IHBT-Gr-01		IHBT-Gr-07		E-3	
		2013-14	2014-15	2013-14	2014-15	2013-14	2014-15
1.	Leaf number per plant/year	31.25	33.15	32.05	33.05	29.3	31.85
2.	Plant spread (cm)/year	57.75	58.45	48.95	49.45	63.25*	63.8*
3.	No. of suckers / plant/year	2.45	4.45	2.7	6.25	2.75	5.7
4.	Leaf length (cm)	32.55	32.60	23.0	23.75	34.25*	33.425*
5.	Leaf width (cm)	11.55	11.3	7.17	7.075	12.87*	11.975*
6.	Flower no. per plant/year	21.15	22.95	17.6	19.55	18.8	21.8
7.	Scape length (cm)	34.6	33.55	50.05	49.05	39.3	40.75
8.	Flower diameter (cm)	8.22	8.25	7.55	7.775	8.05	8.225

*Significant at $P = .05$

S. No.	Source of variation	IHBT-Gr-02		IHBT-Gr-03		23-1	
		2013-14	2014-15	2013-14	2014-15	2013-14	2014-15
1.	Leaf number per plant/year	20.85	23.7	22.75	24.45	25.4	28.65
2.	Plant spread (cm)/year	66.05	66.55	66.25	66.8	65.85	66.55
3.	No. of suckers / plant/year	2.3	4.5	2.45	5.15	2.8	5.75
4.	Leaf length (cm)	37.2	37.31	36.35	35.68	38	37.46
5.	Leaf width (cm)	12.17	11.67	14.67	14.6	16.2*	15.22*

287	6.	Flower no. per plant/year	13.95	16.35	12.75	14.75	23.5*	23.4*
288	7.	Scape length (cm)	32.8	31.95	32.6	31.5	45.4*	44.75*
289	8.	Flower diameter (cm)	9.65	9.72	9.35	9.55	10.85	10.75

290 *Significant at $P = .05$

291

292 Table5. Analysis of variance for quantitative traits among new gerbera hybrid genotypes and
293 their parental lines during 2013-14 and 2014-15 under poly-house conditions

294	S. No.	Source of variation	2013-14		2014-15	
295			Variance	F- value	Variance	F- value
296	1.	Leaf no. per plant/year	85.81	367.09*	72.43	185.25*
297	2.	Plant spread (cm)/year	189.34	472.31*	190.21	498.80*
298	3.	No. of suckers/plant/year	0.16	5.66*	2.12	21.68*
299	4.	Leaf length (cm)	123.00	421.41*	104.66	471.24*
300	5.	Leaf width (cm)	38.32	388.85*	33.60	306.44*
301	6.	Flower no. /plant/year	67.94	323.36*	51.48	137.89*
302	7.	Scape length (cm)	209.45	750.73*	217.42	497.28*
303	8.	Flower diameter (cm)	6.04	97.88*	5.23	62.73*

304 *Significant at $P = .05$

305

306 4. CONCLUSION

307

308 In conclusion, gerbera is one of the important cut flowers having commercial significance.
309 Vegetative propagation is possible through divisions, but rate of multiplication is too slow for
310 commercial purposes. *In vitro* culturing is an important tool for acceleration of plant
311 multiplication and the plants produced through tissue culture are true and are free from
312 diseases. On the basis of the present study a suitable protocol for shoot regeneration,
313 multiplication and root induction has been identified for mass multiplication of gerbera plants.
314 The MS medium supplemented with 1mg/l BAP + 0.030mg/l IBA + 0.025mg/l NAA produced
315 maximum micro-shoots, leaf number and leaf length. Half strength MS medium with 0.4mg/l
316 IBA was observed to be the best medium for early root initiation of *in vitro* raised micro-
317 shoots of gerbera. The protocol can be utilized for mass multiplication of new gerbera
318 genotypes on a commercial scale. Characterization of new hybrid F1 genotypes of gerbera
319 in comparison to parental genotypes under field conditions suggests the scope of
320 improvement of gerbera for desirable floral traits such as double flower shape, standard size
321 of flowers and various colours through breeding program which will result in new varietal
322 types for cultivation in future.

323 REFERENCES

324

325 1. Tjia B. Cooperative extension services. 2001. International Food and Agricultural Science,
326 University of Florida, p. 1-5.

327 2. Singh S, Dhyani D and Karthigeyan S. Characterization of variability in F1 progeny of
328 inter-varietal cross of gerbera (*Gerbera jamesonii* Bolus ex. Hooker F). Journal of
329 Ornamental Horticulture 2009: 12(4): 225-231.

330 3. Singh S, Dhyani D and Kumar Ashok. Flower colour variations in gerbera (*Gerbera*
331 *jamesonii*) population using image analysis. Indian Journal of Agricultural Sciences 2011: 81
332 (12): 1130-1136.

333 4. Singh S, Kumar A and Dhyani D. Inheritance of pigment markers associated with
334 flavonoid and carotenoid pigmentation in flowers of *Gerbera jamesonii*. Canadian Journal of
335 Plant Breeding 2013: 1(3): 110-114.

336 5. Peper H, Brandis AV, Dopke H. Clonal propagation of gerberas can be profitable. Result
337 from Ahlem on the culture and clonal propagation of gerberas. Taspo 1971: 105, 7.

338 6. Murashige I, Skoog F. A revised medium for rapid growth and bioassay with tobacco
339 tissue cultures. Physiol. Plant 1962:15, 473-487.

340 7. Meeteren U Van. Water relations and keeping-quality of cut gerbera flowers. I. The cause
341 of stem break. Scientia Horticulturae 1978: 65-74.

342 8. RHS colour chart. Published by the Royal Horticultural Society, 80 Vincent Square,
343 London, SW1P 2PE. 1995.

344 9. Drennan D, Harding J, Byrne TG. Heritability of inflorescence and floret traits in gerbera.
345 Euphytica 1986: 35: 319-330.

346 10. Reynoird JP, Chriqui D, Noin M, Brown Sand Marie D. Plant propagation from in vitro
347 leaf culture of several gerbera species. Plant Cell Tissue and Organ Culture 1993: 33: 203-
348 210.

349 11. Aswath C, Choudhary ML. Rapid plant regeneration from *Gerbera jamesonii* Bolus callus
350 cultures. Acta Botanica Croatica 2002: 61:125-134.

351 12. Xi M, Shi JS. Tissue culture and rapid propagation of *Gerbera jamesonii*. Journal
352 of Nanjing Forestry University 2003: 27: 33-36.

353 13. Prasanth M, Sekar K. Studies on age of explant on callus induction in gerbera cv.
354 Mammut. Scientia Horticulturae 2004: 9: 207-211.

355 14. Kumar S, Kanwar JK, Sharma DR. In vitro regeneration of *Gerbera jamesonii* Bolus from
356 leaf and petiole explants. Journal of Plant Biochemistry and Biotechnology 2004: 13: 73-75.

357 15. Zheng XF, Wang JH, Li MY. Factors affecting organogenesis in *Gerbera jamesonii* Bolus
358 cultures in vitro. Journal of Jiangsu Forestry Science and Technology 2002: 29: 29-31.

- 359 16. Modh FK, Dhaduk BK, Shah RR. Factors affecting micropropagation of gerbera from
360 capitulum explants. *Journal of Ornamental Horticulture* 2002; 5: 4-6.
- 361 17. Tyagi P, Kothari SL. In vitro regeneration of *Gerbera jamesonii* (H. Bolus ex Hook f.)
362 from different explants. *Indian Journal of Biotechnology* 2004; 3: 584-586.
- 363 18. Ray T, Saha P, Roy SC. In vitro plant regeneration from young capitulum explants of
364 *Gerbera jamesonii*. *Plant Cell Biotechnology and Molecular Biology* 2005; 6: 35-40.
- 365 19. Kumar S, Kanwar JK. Regeneration ability of petiole, leaf and petal explants in gerbera
366 cut flower cultures in vitro. *Folia Horticulturae* 2006; 18: 57-64.
- 367 20. Pierik RLM, Jansen JLM, Maasdam A, Binnendijk CM. Optimization of gerbera plantlet
368 production from excised capitulum explants. *Scientia Horticulturae* 1975; 3: 351-357.
- 369 21. Laliberte S, Chretien I, Veith J. In vitro plantlet production from young capitulum explants
370 of *Gerbera jamesonii*. *Horticulture Science* 1985; 20: 137-139.
- 371 22. Cheema K L, Hussain M. Micropropagation of sugarcane through apical bud and axillary
372 bud. *International Journal of Agriculture and Biology* 2004; 2: 257-9.
- 373 23. Al-Khayri J M, Al-Bahrany A M. Genotype dependent in vitro response of date palm
374 (*Phoenix dactylifera* L.) cultivars to silver nitrate. *Scientia Horticulturae* 2004; 99: 153-62.
- 375 24. Murashige T, Sepra M, Jones JB. Clonal multiplication of gerbera through tissue culture.
376 *Hort. Sci.* 1974; 9: 175-180.
- 377 25. Schiva T, Lercari B, Giusta R. Micropropagation of *Gerbera*: variable response to in vitro
378 culture. *Annali-dell'Istituto-Sperimentale-per-la-Floricoltura* 1982; 13: 56-61.
- 379 26. Barbosa MHP, Pasqual M, Pinto JEBP, Arello EF, Barros IDe, Barros I. Effects of benzyl
380 amino purine (BAP) and 3-indole acetic acid (IAA) on in vitro propagation of *Gerbera*
381 (*Gerbera jamesonii* Bolus ex Hook) cv. Appelbloesem. *Pesquisa Agropecuaria Brasileira*
382 1993; 28: 15-19.
- 383 27. Mariam A, Mahmed A, Igbal A. Evaluation of micro and macro propagation techniques of
384 *Gerbera* (*Gerbera Jamesonii*) under different conditions. Second RUFORUM Biennial
385 Meeting 2010: 20 - 24 September.
- 386 28. Shailaja VP. Studies on in vitro propagation of *Gerbera jamesonii* Bolus. M.Sc. Thesis,
387 University of Agricultural Sciences, Dharwad 2002. (India).
- 388 29. Pierik RLM, Steegmans HHM, Verhaegh JAM, Wouters AN. Effect of cytokinin and
389 cultivar on shoot formation of *Gerbera jamesonii* in vitro. *Netherlands J. Agric. Sci.* 1982; 30:
390 341-346.
- 391 30. Harel D, Kuzmicic I, Jug-Dujakovic M, Jelaska S. The effect of genotype on *Gerbera*
392 shoots multiplication in vitro. *Acta Bot. Croatia* 1993; 52: 25-32.
- 393 31. Akter N, Hoque MI and Sarker RH. In vitro propagation in three varieties of gerbera
394 (*Gerbera jamesonii*Bolus.) from flower bud and flower stalk explants *Plant Tissue Culture*
395 and *Biotechnology* 2012; 22(2): 143-152.