

1
2
3
4
5
6
7
8
9

Short Research Article A Rapid Morphological Screening Procedure for Pea Under Drought Stress in Greenhouse Settings

Aims: This work proposes a rapid morphological screening procedure for peas grown under drought stress aiming to provide phenotypic evaluations for further research. **Material and methods:** the plant yield and morphological changes of two widely grown pea cultivars to drought stress was evaluated in this work. A total of eighty seeds of each pea variety were grown in a randomized completed block with four replicated in a greenhouse located in Moscow, ID. Plants were subjected to optimal irrigation and two treatments that reduced optimal irrigation rates by 40% and 60% to induce drought stress. The test was repeated. **Results:** Varieties differed in their response to water deficiency but emergence and flowering dates were not affected by the treatments. Treatments were also significantly different for all measurements (root pod numbers, width, length and number of viable seeds, shoot height, root length, and shoot dry weight). Root weight was not significantly different between treatments. **Conclusions:** It was possible to observe the effect of drought in each treatment with significant differences between varieties and treatment ($p=0.00$). The variety 'Banner' appeared to be the most drought tolerant, which was reflected by both growth and yield parameters. 'Aragorn' had shown many drought stress symptoms even in the moderate drought treatment, such as very low yield. Our work also identified potential parental material to be used in breeding for drought tolerance.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Keywords: drought, greenhouse plants, agronomic methods, stress tolerance, peas.

1. INTRODUCTION

Over the past several decades, drought stress has become one of the biggest concerns in agriculture worldwide. Adaptation to abiotic stresses is an important acquirement of agriculturally relevant crops and development of enhanced drought tolerance plays an essential role in plant breeding [1].

The development of drought tolerant plants is of increasing priority and relevance as the area affected by significant drought is expected to markedly increase in the future due to global climate change [2]. While progressive water withdrawal has been studied in the greenhouse for beans, it generally does not mimic actual field conditions [3].

While limited studies and techniques for simulating field drought stress in a greenhouse environment have been conducted [4], more studies are needed to adequately formulate a reliable protocol for legumes, especially peas. It is the aim of this research to provide a rapid morphological screening procedure for pea under drought stress in greenhouse settings.

The use of peas as a model crop for genetic studies has a very rich history, beginning with Mendel in the 1860s. *Pisum* spp. is adapted to cool, semi-arid to sub-humid growing conditions [5]. However, pea has been grown in diverse environments across the globe and the demand for production continues to increase [6]. Drought is of great concern in pea and

it has been reported to result in a loss of between 38.5 and 43% production under different stress levels [7].

2. MATERIAL AND METHODS

Two dry pea cultivars, 'Aragorn' and 'Banner,' commonly grown in the U.S. and New Zealand and developed by ProGene Plant Research LLC in conjunction with NZ Plant Research LTD were utilized for this experiment. 80 seeds of each genotype were prepared by sterilization utilizing a 10% bleach aqueous solution, rinsed in sterile, distilled H₂O and air dried for a period of 24 hours.

Five seeds of each variety were planted in a 3.75 liter high density polyethylene/polypropylene polycan container (Anderson Pots, Oregon, USA) filled with 690 g of Sunshine Mix #1 growing media (Sun Gro Horticulture, Agawam, MA) and arranged in a complete randomized design with four replicates per cultivar. The experiment was repeated.

The peas were grown under controlled greenhouse conditions with temperatures ranging from 12 to 18°C at night and 21 to 27°C during daylight hours (MicroGrow Greenhouse Systems Growmate (Temecula, CA) utilizing 400 Philips HID Ceramalux High Pressure Sodium bulb). The plants were subjected to a 12-hour photoperiod per 24 hours.

Three different irrigation regimes were tested: control, moderate drought and severe drought. Field capacity of the growing media served as the experimental control. Field capacity was determined by wetting 690 g of media until saturation was reached. The media was then allowed to drain for a period of 24 hours and oven dried to determine maximum moisture holding capacity (field capacity).

Previous field trials conducted by ProGene Plant Research [8] have shown that approximate reductions in irrigation of 40% and 60% caused visual drought stress symptoms on plots of field peas. Control plants were irrigated to the full field capacity of the growing media (1400 ml) and irrigated once every 7 days until flowering and then once every 3.5 days from flowering until maturity. Drought stress test pots received amounts of water decreased by 40% (840 mL) and 60% (560 mL) of field capacity, respectively, at the same watering interval.


Phenotyping methods consisted of the collection of the following agronomic notes: emergence and flowering dates, plant survival, shoot height and root length, shoot and root dry weight, number of pods per plant, pod length and width, and number of viable seeds produced per plant. Statistical analysis of the data was performed using Minitab 17 Statistical Software (Minitab, State College, PA) and Agrobase Generation II® (Agronomix Software Inc., Winnipeg, Manitoba).

3. RESULTS AND DISCUSSION

As shown in previous research, emergence and flowering dates were not affected by drought stress treatments [9]. However, varieties differed in their response to water deficiency and plant survival was jeopardized. The 40% reduction from field capacity trial yielded a 20% rate of mortality for both varieties tested. The 60% reduction from field capacity trial resulted in 20% mortality among 'Aragorn' and 27% among 'Banner.'

A statistically significant difference in shoot height ($P=0.0$) was observed in both variety and treatment ($P=0.0$) comparisons. 'Banner' was taller than 'Aragorn' in all treatments and both

77 varieties suffered from increasing stunt with relation to decreasing irrigation (Fig.1).



78

79 Fig.1 Shoot height (cm) of 'Banner' and 'Aragorn' across water treatments - control (1400
80 ml), moderate drought (840 ml) and severe drought (560 ml) stress.

81

82 'Banner' and 'Aragorn' exhibited a 6.3% and 9.8% reduction in plant shoot height,
83 respectively, in the moderate drought treatment. Both had the least shoot height, at only
84 70% of their potential, when compared to the control plants, in the severe drought treatment.
85 Shoot dry weight was also impacted by drought, being significantly different per treatment
86 ($P=0.0$) and per variety ($P=0.0$). For the moderate and severe drought stress treatments
87 'Banner' shoot dry weight (Figure 2) was reduced by 44% and 70%, respectively, while
88 'Aragorn' was reduced by 55% and 69%, respectively.

89 Growth reduction measured by height and dry weight was observed in the drought stress
90 treatments. It was also observed that leaf area was visually reduced. In a similar study
91 involving wheat cultivars, leaf area was also reduced but root development was retarded in
92 only one cultivar [10]. It was also showed a reduction in plant height, lower leaf and stem dry
93 weights, and less overall aerial biomass among drought-stressed plants grown in a
94 greenhouse environment [4].

95




Fig.2 Shoot weight (g) of 'Banner' and 'Aragorn' across water treatments: control (1400 ml), moderate drought (840 ml) and severe drought (560 ml) stress.

Root length differed statistically per treatment ($P=0.1$) but not per variety ($P=0.86$). 'Aragorn' roots were not consistently affected on the moderate drought treatment but had an important reduction of 32% on root length on the severe drought; while 'Banner' roots were not consistently affected among the treatments. Root dry weight did not differ significantly per treatment ($P=0.99$) but did among varieties ($P=0.3$). Overall, 'Aragorn' produced more roots than 'Banner' in all trials.

The number of pods per plant was statistically different per treatment ($P=0.0$) as well as per variety ($P=0.03$). In all treatments, 'Banner' had more pods than 'Aragorn' but both varieties exhibited a reduction in the number of pods with increasing drought. 'Banner' displayed a 38.5% and 41.5% reduction in the overall number of pods as drought increased, respectively; while 'Aragorn' had a reduction of 42.7% and 56%, respectively.

In addition, pods also became shorter and thinner with decreasing irrigation when compared to the control. 'Aragorn' did not produce any viable seeds in either drought treatment while 'Banner' was able to produce viable seed throughout all treatments. 'Banner' suffered a reduction in the number of viable seeds by 19% in the moderate drought treatment and 27% in the severe drought treatment.

Yield measured by yield components (number of pods, pod length, width, and number of viable seeds per pod) was also reduced under simulated drought conditions, with simulated severe drought underperforming all other treatments. In a similar study, analysis of variance has indicated that there were significant differences among genotypes regarding agronomic and morphological traits of wheat under drought [11].

Plant characteristics can influence drought tolerance. Comparing the two varieties used in our study, 'Banner', the variety that performed better under drought stress, has good early vigor being five to six days earlier than 'Aragorn' [8].

There are many metabolic changes that can enable a plant to withstand drought, including specific proteins such as chaperonins and dehydrins that are induced in certain plants under drought [12]. Water stress also induced the accumulation of soluble sugars in epicotyls between 2.8- and 5.1-fold in another study and the osmotic adjustment varied greatly depending on cultivar [13]. Moreover, temperature can contribute to drought stress and can induce significant changes in plant cell gene expression [14].

4. CONCLUSION

Our study has proposed a simple, rapid morphological screening procedure for pea under drought stress in greenhouse settings and provided evaluation of two commercial pea cultivars. 'Banner' appeared to be the most drought tolerant, which was reflected by both growth and yield parameters. 'Aragorn' had shown many drought stress symptoms even in the moderate drought treatment, such as very low yield. Our work also identified potential parental material to be used in breeding. As drought and stress conditions are only expected to increase as global climate change progresses, breeding for drought is a promising area.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Witt S, Galicia L, Lisec J, Cairns J, Tiessen A, Araus JL, Fernie AR. Metabolic and phenotypic responses of greenhouse-grown maize hybrids to experimentally controlled drought stress. *Molecular plant*. 2011; ssr102.
2. de Paiva Rolla AA, Carvalho JD, Fuganti-Pagliarini R, Engels C, do Rio A, Marin SRR, Oliveira MCN, Beneventi MA, Corrêa F, Guimarães M, Farias JBC, Neumaier N, Nakashima K, Yamaguchi-Shinozaki K, Nepomuceno AL. Phenotyping soybean plants transformed with rd29A: AtDREB1A for drought tolerance in the greenhouse and field. *Transgenic research*. 2014; 23(1), 75-87.
3. Kavar T, Maras M, Kidric M, Sustar-Vozlic J, Meglic V. Identification of genes involved in the response of leaves of *Phaseolus vulgaris* to drought stress. *Mol Breeding*. 2008; 1(21), 159-172.
4. Pennypacker BW, Leath KT, Stout WL, Hill RR. Technique for simulating field drought stress in the greenhouse. *Agronomy journal*. 1990; 82(5), 951-957.
5. Coyne C J, McGee RJ, Redden RJ, Ambrose MJ, Furman BJ, Miles CA. Genetic Adjustment to Changing Climates: Pea (pp. 238-250). Wiley Blackwell: Chichester, UK. 2011.
6. EMBRAPA – Empresa Brasileira de Pesquisa Agropecuária. Centro Nacional de Pesquisa de Hortaliças. Ervilha para produção de grãos verdes. 2 Nov. 2008. <www.cnph.embrapa.br/cultivares/ervilha>.

- 168 7. Carvalho JA, Rezende, FC, Aquino RF, Freitas WA, Oliveira EC. Produção da ervilha
169 cultivada em ambiente protegido sob diferentes tensões de água no solo. Revista Brasileira
170 de Engenharia Agrícola e Ambiental, Campina Grande. 2012; 1(16), 44-50.
- 171 8. Braunwart K, Wood M. Drought Stress on Pea Crops. ProGene Plant Research; 2014.
- 172 9. Muchow RC. Phenology, Seed yield and WaterUse of Grain Legumes Grown under
173 Different Water Regimes in a Semi-arid Tropical Environment. FieldCrops Research. 1985;
174 11, 81-87
- 175 10. Klamkowski K, Treder ww. Response to Drought Stress of Three Strawberry Cultivars
176 Grown Under Greenhouse Conditions. J. Fruit Ornam. Plant Res. 2008; 180 (16):179-188.
- 177 11. Ahmadizadeh M, Nori A, Shahbazi H, Aharizad S. Correlated response of morpho-
178 physiological traits of grain yield in durum wheat under normal irrigation and drought stress
179 conditions in greenhouse. African Journal of Biotechnology. 2014; 10(85), 19771-19779.
- 180 12. Vaseca I, Akiscan Y, Demirevska K, Anders I, Feller U. Drought stress tolerance of red
181 and white clover: comparative analysis of some chaperonins and dehydrins. Scientia
182 Horticulturae, Canada. 2011; 4(130), 653-659.
- 183 13. Sánchez FJ, Andrés EF, Tenorio J, Ayerbe L. Growth of epicotyls, turgor maintenance
184 and osmotic adjustment in pea plants (*Pisum sativum* L.) subjected to water stress. Field
185 Crops Research. 2014; 2(86), 81-90.
- 186 14. Talalaiev O, Korduym E. Expression of small heat shock protein (sHSP) genes in the
187 garden pea (*Pisum sativum*) under slow horizontal clinorotation. Plant signaling & behavior.
188 2014; 9(4), e29035.