

Original Research Article

BLOOD PROFILE OF WEST AFRICAN DWARF GOATS FED COMPLETE DIETS CONTAINING GRADED LEVELS OF SWEET ORANGE PEEL MEAL

ABSTRACT

Sixteen West African dwarf (WAD) goats of about 6 - 8 months of age, having an average weight of 6.79 kg were used in a completely randomized design to evaluate the haematology and serum biochemistry of WAD goats fed diets containing graded levels of sweet orange peel meal (SOPM). Four dietary treatments were formulated and compounded to contain 0%, 12.5%, 25%, and 50% of SOPM, and these were designated T₁, T₂, T₃ and T₄ respectively. Results showed that the values of total protein (TP), haemoglobin (Hb), red blood cells (RBC), white blood cells (WBC), mean corpuscular haemoglobin (MCH), mean corpuscular volume (MCV), and mean corpuscular haemoglobin concentration (MCHC), did not show significant difference ($p>0.05$) among the treatments means. Serum biochemical components such as urea, total protein, albumin, serum glutamic oxaloacetic transaminase (SGOT), serum glutamic pyruvic transaminase (SGPT) and creatinine also did not show any significant variation ($p>0.05$) among the treatments. The study showed that replacement of maize offal with SOPM up to 50% level in the complete diets of WAD goats did not adversely affect their haematology and serum biochemistry.

Key words : Haematology, serum biochemistry, sweet orange peel meal, WAD goats.

Introduction

Ruminants constitute a very important part in the livestock subsector of the Nigerian agricultural economy. The potential of small ruminant production in alleviating the low animal protein intake by man in developing nations such as in Nigeria needs no emphasis (Fajemisin *et al.*, (2010). Notwithstanding, the high cost of formulating livestock feed has been a major constraint militating against the increased production of valuable sources of animal protein (Fajemisin and Adeleye, 2005) in Nigeria. This has necessitated the use of agricultural by products, crop residues and agro industrial by products. A good number of by products is derived from sweet orange fruit, these include citrus pulp, citrus molasses, citrus peel oil and citrus peel. Sweet orange peels in Nigeria can be easily collected at no cost from sweet orange fruit retailers who peel and sell sweet orange fruits to consumers for direct consumption. The peels are available all through the year and can constitute environmental hazard when poorly disposed, but if properly harnessed it can be a potential feed ingredient for ruminant feeding. Bawala *et al.* (2008) reported that nutritional studies should not be limited to performance, carcass quality and protein intake alone, as the effect of feed materials on blood constituents is also very relevant and should be evaluated. Laboratory study and tests on blood profile are important tools that help detect any deviation from normal animal or human body (Benjamin, 1981). Blood examination is a good way of assessing the health status of animals as it plays a vital role in the physiological, nutritional and pathological status of animals (Aderinboye *et al.*, 2009). Haematology and serum

biochemistry facilitates diagnosis of a disease condition, enhances prognosis and assessment of efficacy of therapy and toxicity of drugs and substances (Fajemisin *et al.*, (2010). Thus, the study was designed to investigate the effect of diets containing sweet orange peels on the haematology and serum biochemistry of West African Dwarf (WAD) goats.

Materials and methods

Sweet orange peels were collected from retailers of sweet orange fruits around Makurdi metropolis. It was sundried on concrete floors for 48 hours, when it became crispy, it was packed, milled and stored in synthetic bags for use. The SOPM was use to replace maize offal in the diets of WAD goats at 0%, 12.5%, 25% and 50% constituting four dietary treatment groups (T₁, T₂, T₃ and T₄ respectively). Sixteen grower WAD goats, aged between 5-8 months and having an average weight of 6.79kg were used. The animals were obtained from medium scale goat producers within Makurdi metropolis and its environs, and were vaccinated against PPR, using PPR vaccine at source and dewormed using ivomectin. Two weeks to the arrival of the animals to the experimental site, the pens were thoroughly washed using a strong disinfectant (moriguard) and allowed to dry. On arrival, the animals were weighed and randomly distributed into four treatment groups of four animals each, in a completely randomized design. Each animal was kept in separate pens which were bedded with wood shaving to serve as both litter materials and beddings. Animals were fed between 08:00hr and 09:00 hr daily and also served with clean water *ad libitum* and for 84 days. On the last day of the experiment, blood was collected from each animal via the jugular vein before morning feeding to analyse for haematological and serum biochemical components. Blood samples for haematology was collected into sterile sample tubes containing EDTA, while sample tubes for serum biochemistry were without EDTA in order to allow the blood to clot for the serum to be decanted for analysis. PCV, Hb, RBC and WBC were determined as reported by Schalm *et al.* (1975). MCV, MCH, and MCHC were calculated from the PCV, Hb, and RBC as established by Jain (1986). Serum total protein was obtained by the biuret method as described by Kohn and Allen (1995), while the albumin was by the bromocresol green method (Doumas and Biggs, 1972). The SGPT and SGOT was determined calorimetrically. Serum urea and creatinine was by the modified method of valley *et al.* (1980) while cholesterol was determined by the method of Abel *et al.* (1952).

Results and discussion

The effect of SOPM on the haematological indices is presented in Table 2. There were no significant differences ($p>0.05$) among the treatments in PCV, Hb and RBC, this implies that the animals were not anaemic (Fajemisin *et al.*, 2010). The PCV values of 25.50 - 27.75% were similar with 25.50 - 28.00% reported by Aderinboye *et al.* (2009) when he evaluated the effect of dietary monensin inclusion on selected haematological parameters of WAD goats. The Hb values of 8.50 - 9.25 g/dl were similar with 8.10 - 9.40 g/dl reported by (Fajemisin *et al.*, 2010) and 8.65 - 9.30 g/dl reported by Aderingboye *et al.* (2009). Since the Hb values were similar to the control, it means that all the treatments were capable of supporting high oxygen carrying capacity in the experimental animals. The WBC values for treatments containing the test

ingredient were similar to the control indicating that the animals were not militating against any disease condition. MCV, MCH, and MCHCH were also not significantly different ($p < 0.05$) among the treatments, implying that the experimental animals were not suffering from any type of anaemia. Result of the effect of SOPM on serum biochemistry is present in Table 3. Urea, total protein, albumin, globulin, SGOT, SGPT, cholesterol and creatinine did not show any significant variation among the treatments. Similarity of the values of urea in treatments having the test ingredient with the control implies that, replacing maize offal with SOPM did not cause any undue elevation in the urea level of the animals as a result of amino acid imbalance. Blood urea level is also commonly considered in ruminants to reflect the protein quality of the diet (Wikipedia, 2009), thus, it also implies that the protein quality of all the diets were adequate. Albumin which is a very strong predictor of health was observed to be normal in this study. The albumin range of 28.47g/l - 33g/l was similar to 29.30 - 32.70g/l reported by Babeyemi *et al.* (2003) when he evaluated the haematological and biochemical components of WAD goats fed *Tephrosia bracteolata*-based forage. SGOT and SGPT values were 189.20 - 198.33 iμ/l and 37.45 - 48.60mg/dl respectively. SGOT and SGPT are liver enzymes, and similarity among all the treatments implies that, treatments containing SOPM were safe and did not hamper with the levels of these enzymes as to tamper with their normal functioning to cause deleterious effects on the liver. Cholesterol levels were normal, meaning that the meat from the experimental animals was safe, and would not lead to cholesterol elevation in consumers. Creatinine values were between 6.93 - 7.81 mg/dl and this was normal, suggesting that, increasing the levels of SOPM from 0 - 50% in the diets of the experimental goats did not have deleterious effects on the lean tissue mass of the animals as to cause emaciation.

Conclusion

The results from this study showed that replacing maize offal with SOPM up to 50% level did not cause any negative effect on the haematological and serum biochemistry of the WAD goats as to jeopardize their health. Therefore farmers can incorporate SOPM in the diets of WAD goats up to 50% level, particularly during the dry period for economic goat production without any adverse effect.

REFERENCES

- Abel, L. C., Levy, B. B., Brodie, B. B and Kendall, F. E. (1952). A Simplified method for estimation of total cholesterol in serum, and demonstration of its specificity. *J. Biol. chem.* 195: 357 – 366.
- Aderinboye, R. Y., Onwuka, C. F. I., Aina, A. B. J. and Oduguwa, O. O. (2009). Effect of dietary monensin inclusion on selected haematological parameters in West African Dwarf goats. Proc. 14th Ann. Conf. Anim. Sc. Ass. Of Nig. (ASAN). Sept. 14th – 17th 2009. LAUTECH Ogbomosho. Nigeria. pp. 619 – 621.
- Babayemi, O.J., Bamikole, M.A and Odunguwa B.O. (2003). Haematological and biochemical components of West African dwarf goats fed *Tephrosia bracteolata* – based forage. *Trop. Anim. Prod. Invest.* 6:31-38.
- Bawala, T. O., Aina, A. B. J., Onwuka, C. F. I., Oni, A. O., Adebisi, Y. and Alli, M. A. (2008). Studies on performance of sheep fed sugar Cane (*saccharum officinarum*) tops supplemented with varying Levels of *Leucaena leucocephala* foliage. Proc. 34th Ann.

- Conf. Nig. Soc. for Anim. Prod. Held on 15th-18th March, 2008 at the University of Uyo, Uyo, Nigeria. pp. 575 – 578.
- Benjamin, M. M. (1981). Outline of veterinary clinical pathology (3rd Edition). The IOWA state University Press. Ames, IOWA, U.S.A. Pp 5 – 162.
- Doumas, B. T and Biggs, H. G. (1972). Simple method of albumin determination. *Clin. Chem.* Vol. 7: 175
- Fajemisin, A.N. and Adeleye, I.O.A. (2005). Time related qualitative changes in the haematological values of rat blood kept at room temperature. Proc. 30th Ann. Conf. Nig. Soc. for Anim. Prod. 30:41 – 44.
- Fajemisin, A. N., Fadiyimu, A. A. and Alokun, J. A. (2010). Nitrogen retention and haematological indices of West African Dwarf rams fed sun dried and fermented rumen digesta and cage hen dropping diets. Proc. 35th Conf. Nig. Soc. for Anim. Prod. 14th – 17th March, 2010. Univ. of Ibadan. Nigeria. pp. 604 – 607.
- Kohn, R. A and Allen, M. S. (1995). Enrichment of proteolytic activity relative to nitrogen in preparation from the rumen for *in vitro* studies. *Anim. feed Sci. and Technol.* 52 (1 & 2): 1- 14.
- Schalm, N., Jain, N. C. and Carroll, E. J. (1975). Veterinary Haematology, 3rd edition. Lea and Febiger, Philadelphia. U.S.A.
- Valley, H., Goweloek, A. H and Bell, M. (1980). Determination of serum urea using Theiacetyl monoxine method. Practical Clinical biochemistry. 5th ed. William Heinemann Medical Books Ltd London.
- Wikipedia (2009). Urea. Free encyclopedia. <http://en.wikipedia.org/wiki/urea>.

Table 1: Dietary composition of experimental diets fed to the WAD goats (%)

Ingredients	T ₁	Experimental diets		
		T ₂	T ₃	T ₄
Rice offal	20.00	20.00	20.00	20.00
Maize offal	48.81	42.71	36.61	24.40
SOPM	0	6.10	12.20	24.41
Full-fat soybean meal	28.19	28.19	28.19	28.19
Bone ash	2.00	2.00	2.00	2.00
Salt	1.00	1.00	1.00	1.00
Total	100.00	100.00	100.00	100.00
Calculated values				
Crude protein	17.00	16.92	16.96	16.92
Crude fibre	15.27	15.30	15.33	15.42
Ether extract	7.95	8.50	9.05	10.15
Nitrogen free extract	59.78	59.22	58.20	57.50
ME (kcal/kg)	3395.14	3419.09	3427.04	3489.81

SOPM = Sweet orange peel meal, ME = Metabolizable energy

Table 2: Effect of sweet orange peel meal on haematological indices of the experimental goats

Parameters	Experimental diets				SEM
	T ₁	T ₂	T ₃	T ₄	
PCV (%)	23.50	27.75	26.75	26.25	1.69 ^{ns}
Hb (g/dl)	8.50	9.25	8.75	8.65	0.55 ^{ns}
RBC (x10 ⁶ /μl)	12.01	13.79	12.38	12.24	1.99 ^{ns}
WBC (x10 ³ /μl)	6.80	7.40	5.30	7.20	1.13 ^{ns}
MCV (fl)	21.71	20.75	22.52	21.08	2.48 ^{ns}
MCH (pg)	7.24	6.91	7.22	7.36	0.79 ^{ns}
MCHC (g/dl)	33.20	33.33	32.81	33.14	0.25 ^{ns}

168 SOPM = Sweet orange peel meal, SEM= Standard error of mean, Hb = haemoglobin

169 RBC = red blood cells, WBC = white blood cells, PCV = packed cell volume

170 MCV= mean corpuscular volume, MCH = mean corpuscular haemoglobin

171 MCHC = mean corpuscular haemoglobin concentration

172 T₁ = 0%SOPM, T₂ = 12.5%SOPM T₃ = 25%SOPM, T₄ = 50%SOPM

173 fl = fentolitre, pg = picogram

174

175 Table 3: Effect of sweet orange peel meal on the serum biochemistry of the experimental goats

Parameters	Experimental diets				SEM
	T ₁	T ₂	T ₃	T ₄	
Urea mmol/l	16.28	13.53	14.86	15.83	2.17 ^{ns}
Total protein g/l	80.06	79.51	82.41	75.78	3.87 ^{ns}
Albumin g/l	30.53	28.47	32.13	33.33	2.85 ^{ns}
Globulin g/l	48.75	50.47	50.29	42.45	3.04 ^{ns}
SGOT iμ/l	198.33	180.20	195.73	193.10	24.11 ^{ns}
SGPT mg/dl	48.60	41.33	44.83	37.45	8.06 ^{ns}
Cholesterol mg/dl	61.50	76.25	84.25	92.00	10.62 ^{ns}
Creatinine μmol/l	7.23	6.93	7.81	7.11	5.45 ^{ns}

176 SGOT = Serum glutamic oxaloacetic transaminase, SGPT = Serum glutamic pyruvic
 177 transaminase, SEM = Standard error of mean, ^{ns} = Not significant

178 T₁ = 0%SOPM, T₂ = 12.5%SOPM, T₃ = 25%SOPM, T₄ = 50%SOPM

179