

Application of Different Reclamation Methods on Salt Affected Soils for Crop Production

ABSTRACT

Effects of salic conditions on soil properties under a given land use type and the methods for reclamation has not received the desired research attention in Niger. Understanding of how soil properties and crop yield respond to the influence of salic conditions is required for application of location specific management strategies for the economic agricultural production. This study was conducted on salt affected soils during 2011/2012 and 2012/2013 crop years to evaluate the soil properties, determine their effects on two test crop performances, and its reclaimability under three different approaches. Reclamation approaches were employed not only to increase efficiency but also to reduce the time of reclamation. Soil samples were collected from the surface (0-15 cm depth) and subsurface (15-30 cm) of soils for laboratory determinations. The soils were treated with gypsum (CaSO_4) only at 100% Gypsum requirement (GR); gypsum plus farm yard manure (FYM) at 25 Mg ha^{-1} and chiseling. Leaching with irrigation water over a period of 2-4 hours per week was done. Rice and millet crops were grown for the two crop years. Result of pre-cultivation soils showed that the physical properties of bulk density had a mean value of 2.37 g cm^{-3} and a low total porosity value of 17.46. Chemically, the soils are high in acidity ($>\text{pH } 7.5$) and then large quantities of exchangeable bases and high base saturation. Post-cultivation results show that gypsum application proved to be the best treatment giving highest grain yield of rice and millet. However, this treatment followed statistically similar results with gypsum + FYM. No application (control) remained low. Chemically, the pHs was reduced after harvesting of second crop (millet 2005-06). Electrical conductivity (ECe) was reduced after growing of first millet crop except control. The reduction of sodium adsorption ration (SAR) was higher after the 2nd crop. The soil parameters in control treatment did not improve. The study concluded that continuation of gypsum + FYM + Chiseling was more effective in improving the soil condition for sustainable land use.

Keywords: Saline soil, management, gypsum, farm yard manure, cereal crops

1.0 INTRODUCTION

Arid and semi-arid regions have always been characterized by permanent features of salinity and sodicity due to fluctuations in climatic conditions. The earth contains a

considerable amount of sodium salts, which mere existence in the soil does not create a problem. However, the excess of salts dissolved in water accumulate slowly and gradually on the surface of the soil as the water evaporates; causing damages to soil properties and crop yield. (Ref)

Soil salinity increases due to capillary action from the saline water table and concentration of salt water in the field. ~~Instances bound in Pakistan, (12) reported that~~ about 6.68 million hectare (M ha) land areas ~~the salt affected and out of these 3.77 M~~ ha ~~is~~ saline and 2.91 M ha ~~is~~ saline-sodic. This area is expected to increase with spread of waterlogging and salinity due to an increment in channel irrigation. In India, ~~(22) reported that~~ about 6.73 M ha area ~~is~~ lying barren or produced very low and uneconomically yields of various crops due to excessive accumulation of salts. (22).

~~Report has it that~~ cultivation of crops especially maize in alkaline soils ultimately resulted in low crop yields(8). Earlier, ~~(9) and (7)~~ observed that concentrated sodium ion caused damage to plant tissue thus reduced plant growth and sometimes plant death(7,9). Similarly, ~~(4) noted that~~ nutritional disorders (~~Ca~~, Ca^{2+} , and Zn^{2+} deficiency and Na^+ toxicity) resulted in poor crop growth. ~~Arising from their studies, the dominant cation in~~ the exchange complex was Na^+ and may deteriorate the soil's physical properties, thus, soluble source of Ca^+ was essential for reclamation of such soils. ~~(10) noted that~~ gypsum was the source of Ca^{2+} most commonly used to reclaim sodic soils and improve soil water infiltration. (10).

The appropriate management of the constrained soil resources for the economic agricultural production is the main emphasis in agriculture. There are different approaches for reclamation of salt affected soils. The prominent ones are chemical,

biological and agronomic. The combination of these approaches not only increases the efficiency but also reduce the time of reclamation. The crop production and fertilizer efficiency of these soils can be increased by integrated approaches i.e. use of amendment preferably gypsum and organic manure which helps in maximizing and sustaining yields, improving soil health and input use efficiency (25).

The physical methods of soil reclamation include deep ploughing, sub soiling, sanding, flushing salt out through the soil by applying water periodically and use of acid forming fertilizers to raise the acid status of the soils.  reported that subsoiling (50+5cm crosswise furrows (20-150cm apart) and rice-wheat crop rotation successfully reclaimed two calcareous saline-sodic soils within a period of three years. (15).

In Keana, a town in north central Nigeria, the community has long history of salt mining and processing but due to crude methods employed; the soils might have been affected. This would have consequent effect on cultivated crops and overall agricultural productivity of the area .

The objectives of the study were to: examine the extent of salinity of the soils; determine the efficiency of the applied reclamation approaches on the improvement of salt-affected soils productivity, over two crop production cycles.

2.0 MATERIALS AND METHODS

2.1 Description of the study location

The study was conducted in 2012 and 2013 crop years at the salt mining village in Keana town, which serves as the Headquarter of Keana Local Government Area (LGA) of Nasarawas State, North Central Nigeria. Keana L.G.A has a population of 253, 186 (16). It is located Latitude 8⁰05' 00" E; Longitude 8⁰45' 27" N and altitude of 600m above

sea level, masl (Ministry of Land and Survey, 2006). The weather is that of tropical humid type with distinct rainy and dry seasons. Keana has a mean annual rainfall of 1553.28mm; mean annual maximum temperature of 34.12⁰C, minimum of 22.60⁰C (16).

The area is characterized by a gently undulating topography with a soil type of mostly loam (1). Cereal cropping system of maize, millet, sorghum and rice is the main agricultural use of the land.

2.2 Field and Laboratory Techniques

The field study site was identified after a reconnaissance visit to know nature of salt affected soils and type of reclamation measures to take; drainage characteristics, topography and presence of hard pan. The first site was one hectare of land under no cultivation of crops and quite adjacent to the salt water pond. Soil samples were randomly taken from the fields at the depth of 0-15 and 15-30cm following difference in vegetation pattern. These depths represented the depth of tillage where most nutrients and organic matter are found (6). Samples for the field were composited and bulked, taken to the laboratory, air dried for physical and chemical determinations. Water samples were taken from the salt pond for determination of the Na⁺, Ca²⁺ and Mg²⁺ load.

Second field study was cultivation of the selected field. The land was cleared, leveled and plots prepared. Leveling was to ensure uniform application of water and prevention of accumulation of water in the field. Treatments were arranged in randomized complete block design (RCBD) with three replications. The following treatments were used for the experiment: T₁ = Control; T₂ = Gypsum @100%GR; T₃= T₂+ Light cultivation (chiseling); T₄= T₂+ farm yard manure (FYM) @ 25tha⁻¹; T₅= T₂+ chiseling + FYM@ 25tha⁻¹. The soils in T₃and T₅were tilled with ox driven chisel plow whereas other

treatments were prepared manually with hoe. The gypsum and FYM were applied with subsequent leaching with drainage water.

Rice and millet were grown in sequence for two years. The yield data was recorded at maturity and analyzed statistically using critical difference (CD) test (24). Post harvest soil samples were collected from 0-30cm soil depth after each harvest. This depth is reasoned to provide favorable environment for feeder crops that are not deep rooted. (6) earlier noted that soil physical and chemical characteristics at the two depths (0-15 and 15-30cm) are usually related to cereal crop yield.

Other field studies: Twelve core samplers were used to collect undisturbed soil samples. They were properly labeled for easy identification in the laboratory. They were weighed; oven-dried for 24 hours at 105⁰C and re-weighed. The weights obtained thereafter were used to determine bulk density, porosity and moisture content parameters. These physical characteristic determinations were done at Agronomy laboratory of College of Agriculture, Lafia, Nasarawa State.

2.3 Laboratory Determinations

Soil samples from the field were air-dried, gently crushed and sieved through a 2mm mesh and analyzed in the laboratory for the following properties: Soil particle size distribution, soil p^H, total N, Organic carbon, available P, exchangeable bases (Ca, Mg, Na and K), and cation exchange capacity (CEC). Total acidity, basic saturation and sodium adsorption ratio were also determined.

Particle size distribution (textures) was obtained by the hydrometer method (Day, 1965). Soil pH was determined using Beekman Zeromatic p^H meter after equilibrating for 30 minutes (McLean, 1982). Organic carbon, total N and available phosphorus were

measured by wet-oxidation method (Nelson and Sommers, 1982; Olsen and Sommers, 1982), Micro kjeldahl method (Bremmer and Mulvaney, 1982), respectively. N_a and K were obtained by using the flame photometer, while soluble Ca^{2+} and Mg^{2+} were determined by atomic absorption spectrophotometer (AAS) (Odiete *et al.*, 2006). These determinations including soluble salt content (TSS) were done at the standard laboratory of Federal College of Land Resources Technology, Kuru, Jos, Nigeria.

In terms of physical determinations, Bulk density was obtained by the method of Blake and Hartge (1986), while percentage moisture content was calculate as:

$$\frac{\text{Weight of wet soil} - \text{weight of oven dry soil}}{\text{Weight of wet soil}} \times 100;$$

Percentage porosity = $\frac{\text{wt. of wet soil} - \text{oven dry wt.}}{\text{Volume of sampler}} \times 100;$

Exchangeable sodium percentage (ESP) = $\frac{\text{Exch. Sodium ions}}{\text{Soil CEC}} \times 100$

Sodium adsorption ratio (SAR) = $\frac{Na}{(Ca^{2+} + Mg^{++})^{1/2}}$

Gypsum requirement according to Schoonover's method (26) was used.

Statistical analysis

Statistical differences were tested using two-way analysis of variance (ANOVA) following the General Linear Model (GLM) procedure of Statsoff version 5.5 for Windows (Statsoff, 1998). The least significant difference ($LSD_{0.05}$) was used for mean separation when the analysis of variance showed statistically significant differences ($P < 0.05$) between the investigated properties.

3.0 RESULTS AND DISCUSSION

3.1 Effects of Treatment on Soil Physical Properties

160 The results of the pre-treatment soil analysis showed that the soil has loamy sand texture
 161 with bulk density of the undisturbed soil (mean value of 2.37 g cm^{-3}); an indication of hard
 162 structure (Table 1).

163 Table 1: Soil Physical properties of Kean saline soils

Core	Sand	Silt	Clay	Textural class	Bulk density (g/cm)	Moisture content (%)	Porosity (%)
A	54	36	10	Loamy sand (LS)	2.33	5	12.21
B	57	34	9	LS	2.44	8	21.36
C	55	36	9	LS	2.30	6	16.27
D	56	36	8	LS	2.37	8	19.33
E	58	36	6	LS	2.52	11	26.45
F	50	39	11	LS	2.30	4	9.16
Mean	55	36.2	8.8	LS	2.37	7	17.46

164

165 The implication to agriculture is that enough water will not be retained. This trend was
 166 substantiated by low values of infiltration rates and percentage porosity (19.46%)
 167 obtained. This may be associated to alkaline nature of soils which usually make soils
 168 impermeable to water and air. However, soil analysis of cultivated soils had better
 169 physical properties that will favor crops production in terms of soils texture, porosity and
 170 water retention.

171 In terms of pre-cultivation soil chemical analysis, the soil had ~~high~~ p^{H} range of 7.40 to
 172 7.90 indicating alkalinity (Table 2) 

173 Table 2: Properties of keana saline soils.

Particle size analysis					Exchangeable bases Cmol/kg											
Depth (cm)	Sand %	Silt %	Clay %	Textl. class	Ca	Mg	K	Na	OC %	TN %	P ppm	CEC Cmol/kg	Exch. Acidity	BS %	SAR	Esp %
0-15	80	18	2	LS	18.54	1.60	5.37	3.83	0.74	0.070	43.75	36.0	0.10	81.50	1.21	10.64
15-30	78	18	4	„	18.74	1.47	4.48	4.78	0.74	0.053	43.75	32.0	1.10	92.09	2.18	14.94
Mean	79	18	3	„	18.64	1.54	4.93	4.31	0.74	0.062	43.75	34.0	0.60	86.79	1.68	12.75

174 NB: LS = loamy sand, Ca = calcium, Mg = magnesium, k = potassium, Na = sodium,
 175 OC= organic carbon, TN = Total nitrogen, P = phosphorus, BS = base saturation, SAR =
 176 Sodium adsorption ratio, ESP = Exchangeable sodium percentage.

177
178
179 Further presence of large quantities of exchangeable bases ($\text{Ca} = 18.64 \text{Cmolkg}^{-1}$; $\text{Mg} =$
180 1.535Cmolkg^{-1} ; $\text{K} = 4.925 \text{Cmolkg}^{-1}$ and $\text{Na} = 4.305 \text{Cmolkg}^{-1}$) and high base saturation
181 (86.79%) confirmed the alkalinity nature of the soils. Low sodium adsorption ratio and
182 exchangeable sodium percentage (1.675 and 12.75%, respectively) also confirmed that
183 the soil was alkaline (Table 2).

184 The result of the water analysis showed that the degree of impurities in the water sample
185 was very negligible. The only elements identified were Ca ($43.0664 \text{ mg L}^{-1}$) and Mg
186 (3.9647 mg L^{-1}) (Table 3).

187 Table 3: Chemical properties of the saline water 

Element	mg L^{-1}
Magnesium (Mg)	3.9647
Calcium (Ca)	43.0664

188
189 This suggested that the water was alkaline and may serve as the major sources of high
190 basic cations found in the soil.

191 3.2 Effects of Treatment on Soil Chemical Properties

192 The results showed that the soils were high in soil pH. High soil p^{H} denoted the
193 dominance of sodium among the cations and carbonate /bicarbonates among the anions.
194 These important chemical parameters decreased in all the treatments except the control
195 (Figures 1 and 2).

196

197

198




Fig 1: Original and Post crop soil analysis for pHs (0-15cm).Key: □ Original analysis; ○ Post rice 2012; Δ Post millet 2011-2012;— Post rice 2013; + Post millet 2012-13

significantly increased when different amendments and cultural practices were applied before transplanting of rice in 2012 with subsequent leaching than control. This increase was higher in gypsum (T₂) alone or T₄ as compared to T₃ (gypsum + chiseling) or T₅ (gypsum + fym + chiseling). Similar results by (15) were also reported on two calcareous sodic soils in 4 years of cropping. The average rice paddy yield from both the soils was in the order: gypsum (1.99 Mgha⁻¹) >gypsum + subsoiling (1.84 Mgha⁻¹) > subsoiling (1.41Mgha⁻¹) > bioremediation (1.02 Mgha⁻¹). Gypsum + subsoiling treatments had similar values for wheat grain yield (2.72Mgha⁻¹) followed by subsoling (1.79Mgha⁻¹) and bioremediation (1.46Mgha⁻¹).




Fig 2: Original and Post crop soil analysis for pHs (15-30cm). Key: □ Original analysis; ○ Post rice 2012; Δ Post millet 2011-2012; – Post rice 2013; + Post millet 2012-13

Crop cultivation and application of gypsum alone or in combination with FYM reduced these parameters. However, pH value was greater than 8.5 in control, gypsum + chiseling, and gypsum + FYM + chiseling treatments.

The gradual decrease in p^H values was observed after harvesting of each crop. The p^H values reduced to <8.0 in all the treatments as compared to the control. As far as the lower depth (15-30cm) was concerned, the p^H also reduced in all treatments after harvesting millet in the 2012-13 crop years. This may be attributed to the removal of carbonates and bicarbonates of Na to a greater extent during reclamation. Similar results were obtained by (14).





Fig 3: Original and Post crop soil analysis for ECe (0-15cm). Key: □ Original analysis; ○ Post rice 2012; Δ Post millet 2011-2012;— Post rice 2013; + Post millet 2012-13



273 Fig 4: Original and Post crop soil analysis for ECe (15-30cm). Key: □ Original analysis;
 274 ○ Post rice 2012; Δ Post millet 2011-2012; – Post rice 2013; + Post millet 2012-13
 275

276 A significant decrease in electrical conductivity (ECe) (Figures 3 and 4) was observed
 277 when gypsum at 100%GR (T₂) was applied alone or gypsum + FYM (T₄). The decrease in
 278 ECe might be due to light texture of the soil. However, a little decrease was observed in
 279 the control plots. The best treatment appeared to be the combination of all treatments (T₅)
 280 i.e. gypsum + FYM + Chisel. Possible reason may be the improvement in porosity and
 281 hydraulic conductivity which resulted in enhanced leaching of salts.

282 Decrease in ECe was also reported (21). The application of gypsum at 12 t ha⁻¹ and other
 283 cultural practices during reclamation of dense sodic soil decreased p^H values from 10.2 to
 284 9.1. ECe decreased from 2.1 to 0.8 d.Sm⁻¹ more rapidly during first year of reclamation
 285 but later on, the effect of amendments was still evident but the rate of amelioration was
 286 slow (20). The ECe was higher in the lower depth than the upper depth indicating the
 287 downward movement of salt due to reclamation process after harvesting of first rice crop.

288 It decreased
 289 gradually and reached a tolerable level in the lower depth also after 2012-2013.

290 Sodium adsorption ratio (SAR); the SAR decreased significantly with the different
 291 treatments as can be seen in Figures 5 and 6.




Fig. 5: Original and Post crop soil analysis for SAR (0-15cm). Key: □ Original analysis; ○ Post rice 2012; Δ Post millet 2011-2012; – Post rice 2013; + Post millet 2012-13




Fig 6: Original and Post crop soil analysis for SAR (15-30cm). Key: □ Original analysis; ○ Post rice 2012; Δ Post millet 2011-2012; – Post rice 2013; + Post millet 2012-13

However, the decrease in SAR was more with treatments 5 (gypsum + fym +chiseling) than T₄ (gypsum + fym) and T₂ (gypsum) alone.

The less reduction is SAR in only T₂ (gypsum @ 100% GR) treated plots might be due to slow reaction of gypsum. The most effective treatments were the combination of all the three practices and T₄ (gypsum + fym @ 25tha⁻¹) for reduction of SAR after harvesting the first rice crop. The soil was reclaimed and SAR decreased to safe limits in all treatments except control after harvesting of 3rd crop (rice) in 2013. The decrease in SAR was essentially due to removal of exchangeable Na from the soil complex. The results are in agreement with those of (11). The rate of decrease in SAR was greater in upper soil layer than in lower depth. This pattern was attributed to the decreasing Ca²⁺: Na – ratio in the soil solution as it moved down the profile displacing exchangeable sodium (Na⁺).

3.3 Effect of Treatments on Crop Yields.

Data in Table 4 indicate that biomass of rice and millet was significantly increased when different amendments and cultural practices were applied before transplanting of rice in 2012 with subsequent leaching than control. This increase was higher in gypsum (T₂) alone or T₄ as compared to T₃ (gypsum + chiseling) or T₅ (gypsum + fym + chiseling). Similar results (15, 14) were also reported on two calcareous sodic soils in 4 years of cropping. The average rice paddy yield from both the soils was in the order: gypsum (1.99 Mgha⁻¹) > gypsum + subsoiling (1.84 Mgha⁻¹) > subsoiling (1.41Mgha⁻¹) > bioremediation (1.02 Mgha⁻¹). Gypsum + subsoiling treatments had similar values for wheat grain yield (2.72Mgha⁻¹) followed by subsoling (1.79Mgha⁻¹) and bioremediation (1.46Mgha⁻¹).

Table 4: Biomass and grain yield of Paddy and Millet (Mgha⁻¹) during 2012-13 crop years.

Treatment	Rice 2012		Millet 2012-13		Rice 2012		Millet 2012-13	
	Biomass Paddy		Biomass Grain		Biomass Paddy		Biomass Grain	
T ₁	6.84 ^C	1.04 ^c	8.22 ^c	2.13 ^c	10.33 ^c	1.62 ^c	6.97 ^c	2.06 ^c
T ₂	13.30 ^{ab}	2.70 ^a	8.40 ^{bc}	2.61 ^a	18.35 ^a	3.00 ¹¹	10.66 ^a	3.08 ^a
T ₃	11.13 ^b	2.05 ^b	7.82 ^c	1.96 ^d	14.41 ^b	2.38 ^b	8.54 ^b	2.76 ^b
T ₄	13.96 ^a	2.72 ^a	10.21 ^a	2.60 ^a	17.24 ^a	3.10 ^a	10.36 ^a	2.96 ^{ab}
T ₅	12.81 ^{ab}	2.39 ^{ab}	9.10 ⁶	2.41 ^b	17.91 ^a	2.94 ^a	8.86 ^b	2.83 ^{ab}
LSD	2.476	0.5485	0.6.948	0.1096	1.740	0.3857	0.3471	0.2146

Means sharing same letters are statistically at par at 5% level of probability.

The millet grain yield reduced during 2012-13 in FYM treatments as compared to gypsum alone. The reduction in yield might be due to fading effect of farm yard manure (fym) with passage of time. Gypsum + chiseling remained inferior in production of biomass, paddy and grain yield of millet in all the years than gypsum + farm yard manure + chiseling. However significant increase was found in gypsum + chiseling than control (T₁). The gradual increase in biomass as well as in paddy and millet grain yield in other treatments may be the result of improved soil properties. The results were in line with those of (11) who concluded that most superior combination was gypsum + H₂S04 + Farm yard manure. The improvement in physical and chemical properties of salt affected soil was the major reason for enhancement of crop yield.

4.0 CONCLUSION

The study revealed that gypsum application alone proved the best treatment, followed by gypsum + farm yard manure in increasing rice and millet yield, while gypsum + farm yard manure + chiseling performed better on improving the soil properties.

358 .

359 REFERENCES

- 360 1. Ahmadu Bellow University, ABU. Soils of Assakio, Awe, Doma, Keana, Lafia and
361 Obi. Ahamdu Bellow University Press. 1983; pp: 16-39.
- 362
- 363 2. BlakeGR, Hartge KH. Bulk density. In Methods of Soil Analysis, part 1, A. Klute (ed)
364 Agron. No 9. USA Madison W.I. 1986; pp 370-373.
- 365
- 366 3. Bremmer YM, Mulvaney CS. Total N. *In: Page et al* (Ed.)Methods of Soil Analysis,
367 Part 1. Agron. Monogr. 9 ASA and SSSA. Madison, W.I. 1982; P 595-626.
- 368
- 369 4. Dara SS. Environmental Chemistry and Pollution Control. Int. Science Publication.
370 John Wiley and Sons. 2004
- 371
- 372 5. Day PR. Particle Fractionation and Particle Size Analysis. 1965; P 545-567. In C.A.
373 Black (ed). Methods of Soil Analysis. Part 1. Agron. Monogr. 9 ASA. Madison, W.I.
- 374
- 375 6. Ezeaku PI, Akamigbo FOR, Asadu CLA. Maize yield predictions based on soil fertility
376 parameters in Southeastern Nigeria. Journal of Agro technology and extension, 2002; 2
377 (2):30-38.
- 378
- 379 7. Food and Agricultural Organization, FAO. Irrigation Water Management. Corporate
380 Document Repository. Agricultural Department, FAO, Rome. 1985.
- 381
- 382 8. Frazen D. Managing Saline Soils in North Dakota. *In: NDSU Extension service. SL-*
383 *1087 (revised). 2003; Pp.108*
- 384
- 385 9. Gary EV, Richard TK, Ulery A. Frequently asked questions; saline and/or sodic water
386 and soil. M.S.U. The Department of Land and Environmental Sciences. Extension
387 Service. 1980.
- 388
- 389 10. Gupta RK, Singh CP, Abrol IP. Disolution of gypsum in alkali solution. *Soil*
390 *Science*,1985; 140:382-6.
- 391
- 392 11. Hussain N, Hassan N, Irshadular G, Mojeed F. Evaluation of amendments for the
393 improvement of physical properties of sodic soil. Int. J. Agric. Bio; 2001; 3: 319-22.
- 394
- 395 12. Khan GS. Soil Salinity/Sodality Status in Pakistan. Soil Survey of Pakistan, 1998; pp
396 20-5.
- 397
- 398 13. Mclean EO. Soil pH and Lime Requirements. In: A.L. Pape (ed) Methods of Soil
399 Analysis. Part 2. 2nd ed. Agron. Monogr. 9 ASA and SSSA, Madison W.I. 1982; p. 199-
400 224
- 401

14. Mohammed S, Kaliq A. Comparisons of Various Chemicals and Organic Amendments for Acclaiming Saline/sodic soils. Bull. Irrig. Rain. Flood Control Res. Council, 1975; 5:50-4.
15. Mohammed S, Ghafor A. Reclamation of two saline sodic soils series through subsoiling and gypsum application using marginal water for leaching. *In*: Ahmad R and S. Pietro (eds): prospects for Biosaline Research. Pp 221-33. Proc. Workshop US-Pak Biosaline Resources. Sept. 22-26, 1985. Karachi, Pakistan. 1986; Pp 230
16. Nasarawa Agricultural Development Program (NADP). Weather Records. P4-16 National Population Commission, 2006. Bulletin 13. 2005; p 1-6
17. Nelson DW, Sommers CE. Total Carbon, Organic Carbon and Organic Matters,. *In*: A.C. Page and C.A. Black (eds). Methods f Soil Analysis Part 2, 2nd ed. Agron. Monogr. 9 ASA and SSA, Madison, W.I. 1982; P. 539-579
18. Olsen SA, Sommers CE. Phosphorus. *In*: A.L. Page, C.A. Black (eds) Methods of Soil Analysis. Part 2. 2nd ed. Agron. Monegr. ASA and SSSA, Madison, W.I. 1982; pp. 403-430
19. Odiete .I, Ejekwolu CA, Aadamu JU, Adegboye MA. Different ratios of P-fertilization on some yield indices of Irish potato (Nicola spp) under a typic Haplustalfs at kuru, Jos. Production Agriculture & Technology, 2006; 2 (1): 14-20
20. Rao DIN, Singh NF, Gupta RK, Tyagi NK. Salinity Management for Sustainable agriculture. Control Sol. Salinity Res. Inst., Kamal, India. 1994; 314 p.
21. Sharma DP, Melita KK, Yador JSP. Effect of reclamation practices on soil properties and Crop growth on Farmers field. J. Indian Soc. Soil Sci. 1982; 29:356-60.
22. Sharma RK, Sharma TK. Irrigation Engineering (including Hydrology). S. Chand and co. Ltd. Ram Nagar, New Delhi-110055. 2004; Pp 210
23. Statsoff. Statistical Statsoft Co., 2300 East, 14th Street, Tulasa, Ok 74104, USA. 1998.
24. Steel RGD, Torrie JH. Principles and Procedures of statistics. A Biometrical Approach. 2nd ed. Mc Graw Hill Book. Coy, Inc. N.Y. USA, 1980; 663ppt xxi
25. Swarp .A. Chemistry of Sodic Soils and Management. *In*: Advances in Sodic Land Reclamation. INT. Conf. on Sustainable Management of Sodic Lands, Lucknow, Feb. 9-14.2004; pp 27-52.
26. US Salinity Laboratory Staff, USSLS. Diagnosis and Improvement of Saline and Alkali Soils. USDA Handbook No 60. Washington D.C. USA 1954.