

Effect of dietary inclusion of probiotics and prebiotics on haematological indices, serum biochemistry and gut salmonella count in broilers sourced from Salmonella infected hatcheries within south-west zone of Nigeria.

Comment [FB1]: This is not originally study

Abstract

This study investigated the effect of dietary inclusion of probiotics and prebiotics against gut salmonella, haematological indices and serum biochemistry of broilers sourced from Salmonella infected hatcheries within south-west zone of Nigeria.

In this experiment, 480 male Anak 2000 day-old broiler chickens (160 birds from each state) were collected from hatcheries that were positive to *Salmonella* organisms and used for performance testing for 8 weeks. The birds were laid out in a 3x5 factorial arrangement comprising of 5 dietary treatments including a control and 4 different feed additives. Data were collected on blood, and gut salmonella count. Birds sourced from location 3 showed highest haemoglobin (Hb), white blood cell, serum total protein, uric acid and serum albumin. Among the broiler included in the study, haemoglobin, RBC, total serum protein, serum albumin, uric acid and creatinine were significantly influenced ($P < 0.05$) with highest values recorded for source 3. The WBC, Glucose, serum globulin, uric acid and cholesterol were significantly ($P < 0.05$) reduced by the various additives included. In all the phases of the study, gut salmonella reduced drastically with inclusion of various additives. Birds fed diet supplemented with MOS showed the least gut salmonella count. MOS and AXOS could be used as replacement for antibiotics to improve performance and control the prevalence of *Salmonella* organism in broiler chickens.

Comment [FB2]: Ethical approval for the research was not informed in the manuscript. The authors must include the Protocol number of the Ethics Committee of Animal Use or explain why this information wasn't include in this paper, since the authors used broiler for this experiment

Keywords: Oxytetracycline, Mannose oligosaccharide, Arabiloxylans oligosaccharide and *Pediccocusacidilactici*.

Comment [FB3]: Key words (give 4-5 key words and arrange a-z), separate by, and each entry starts with first capital Word

INTRODUCTION: Poultry production is an attractive enterprise because the production of poultry birds yields high economic value within a very short generation interval. Poultry is an important source of animal protein, income, employment, industrial raw materials, and manure (Rosen, 1995). Therefore, importance of broiler production cannot be over emphasized in the face of the rising demand for animal protein within developing countries like Nigeria. To maximize the genetic potential of broiler for production, these birds must be free from diseases as well as fed with appropriate diets that will meet their requirement for optimal production (Tannock, 1998). Enteric diseases are an important concern to the poultry industry because it reduced productivity, increased mortality and could be linked with contamination of poultry products leading to food poisoning in humans (Tannock, 1998). In the healthy animal, balance in floral ecology in the gastrointestinal tract helps in efficient

Comment [FB4]: Does the manuscript contain new and significant information to justify publication?: No

35 digestion, optimal absorption of nutrients, and increases the body's resistance to infectious
36 diseases (Rosen, 1995).

37 Increase in Salmonella contamination of poultry product which enter the human food chain as
38 a result of infected droppings material, poor hygiene or from infected eggs were reported as
39 major aetiological source for wide prevalence of Salmonella typhoid and paratyphoid cases in
40 approximately 13 million cases of paratyphoid infections worldwide (Murugkaret *al.*, 2005).
41 Unfortunately, poultry meat is the major source of food borne Salmonella paratyphoid
42 infection (Mayrhofer *et al.*, 2003) which had caused severe economic losses, high mortality
43 and medication cost but with attendant reduction in egg production in breeder flocks, poor
44 chicks quality and high cost for eradication and control measures.

45 Salmonella infection or salmonellosis, is major cause of high mortality of wild birds,
46 affecting many bird species worldwide (Hafez, 2001). Several strains of Salmonella bacteria
47 are more closely associated with poultry morbidity than their ubiquitous distribution of
48 various pathogenic serotypes (serovars) like Salmonella pullorum and S. gallinarum to
49 zoonotic serotypes such as S. enteritidis and S. typhimurium (Hafez and Jodas, 2000). These
50 pathogenic serovars constantly passed down the food chain through poultry products to infect
51 numerous consumers. In the poultry industry, antibiotics are used worldwide to prevent
52 poultry pathogens and disease so as to improve meat and egg production. However, the use of
53 dietary antibiotics or feed additives has resulted in high level antimicrobial resistance (Sorum
54 and Sunde, 2001), while its residues in poultry products significantly enhance imbalance of
55 normal intestinal microflora and mucosa fluid and electrolyte. Therefore, alternative and safe
56 intervention to overcome these important problems is the use of probiotics and prebiotics
57 bacteria which have a wide range of nutritional benefit, support physiological absorption
58 activity of intestinal mucosa and effectively combat diarrheal syndromes (Huyghebaert *et al.*,
59 2011). Competition for nutrients, production of antimicrobial compounds (such as low pH
60 volatile fatty acids and bacteriocins), binding sites on the intestinal epithelium and stimulation
61 of immune system (Rolfe, 2000) are proposed mechanisms of pathogen inhibition by
62 probiotics organisms and prebiotics (Koenen *et al.*, 2004).

63 The use of alternative in-feed growth stimulant like probiotics and prebiotics beneficially
64 affect the host microbial flora (Huyghebaert *et al.*, 2011), while probiotics which are direct-fed
65 microbial feed supplements modulate the gut microflora by successfully competing with
66 pathogens through a competitive exclusion process (Mountzouris *et al.*, 2007; [Ayasan, 2013](#)). Beneficial
67 effects of probiotics were observed in toxin neutralization, inhibition of microbial metabolism
68 and immunity stimulation (Sleator, 2008, Sherman *et al.*, 2009) conferring healthy
69 gastrointestinal microbiota on host (Gibson *et al.*, 2004).

70 One of such prebiotic is mannose oligosaccharide (MOS) which is derived from the cell wall
71 of the yeast *Saccharomyces cerevisiae* and effectively influences intestinal activities and
72 animal performance. MOS has the capacity to modulate the immune system and the gut
73 microflora, agglutinate a wide range of pathogenic bacteria and as a result preserve the
74 integrity of the intestinal absorptive surface (Sevage *et al.*; 1996). Reducing intestinal
75 colonization with *S. Enteritidis* using competitive exclusion compounds (probiotics)

Comment [FB5]: The below mentioned literature is related to this paper, but authors did not use them in the text. It must be added these references.

AYASAN T, 2013. Effects of dietary inclusion of protexin (probiotic) on hatchability of Japanese quails. *Indian J Anim Sci*, 83(1): 78-81

during the grow-out period is crucial to improve the microbiological quality of poultry products as well as for consumer health (Vugia 1993). Probiotics are usually live or attenuated organisms and spores (Patterson and Burkholder 2003), non-traditional chemicals (Moore *et al.*, 2006), bacteriophages (Higgins *et al.*, 2005), organic acids (Jarquin *et al.*, 2007; Wolfenden *et al.*, 2007) and several others that have emerged in the last decades which are potentially useful as prebiotic for pathogen control and poultry performance especially with *S. Enteritidis* in newly hatched chicks with successful results (Revolledo *et al.*, 2009).

This study therefore seeks to investigate the effect of dietary inclusion of MOS probiotics on health status, haematological indices and serum biochemistry of broilers sourced from Salmonella infected hatcheries within south-west zone of Nigeria, West Africa.

MATERIALS AND METHODS

Experimental Location: The *in vivo* studies were carried out at the Teaching and Research Farm Development (TREFAD) of the University of Agriculture, Abeokuta, Nigeria. Abeokuta ($7^{\circ} 10' \text{N}$ and $3^{\circ} 2' \text{E}$) area which is 76m above sea level, humid and located in the tropical rain forest vegetation zone with an average temperature of 34.7°C (ref8)

Experimental birds and management: Preliminary screening of all hatcheries located within south-west Nigeria were done by collecting hatching eggs (50 eggs/hatchery) on weekly basis over a period of 8 weeks. Three potentially salmonella infected hatcheries identified in previous findings by the author (Olorunsola *et al.*, 2012) were used in this study. A total number of four hundred and eighty (480) male, Anak 2000 day-old broiler birds sourced from the three major hatcheries located in Oyo state (Location 1), Ogun state (Location 2) and Lagos state (Location 3), all in southwest Nigeria with incidence of salmonellosis were selected for this study. One hundred and sixty (160) day old broiler chicks were randomly sourced from each hatchery making a total of four hundred and eighty (480) broilers used for the study. Birds from each location were allotted into 5 treatment groups of 32 birds each. Each treatment was replicated four times with 8 birds each. Birds contained in each replicate were housed in individual pen each measuring 2.7mx 0.9m with a total floor area of 2.43m^2 . Birds were brooded for 21 days on a deep litter system littered with wood shavings. Water and feed were supplied *ad libitum*. No medication was administered throughout the experimental period which lasted for 56 days.

Experimental Diets: A basal diet containing no additive (control diet) was formulated while additional four diets were formulated to include antibiotics (oxyteracycline at 1g/kg), MOS (1 g/kg), arabinoxylose oligosaccharide (1g/kg) and Sim[®]-Lac (1g/kg). SimLac (a product containing 1×10^{10} cfu viable strain of *Pediococcus acidilactis per gram*) was supplied by a commercial company (Simbiyotec Biological Product Inc. Tuzla [İstanbul-TURKEY](#)). Mannose oligosaccharides (MOS) and arabinoxylans oligosaccharides (AXOS) (Alltech Inc. Kentucky, USA) were used as prebiotics. The inclusion level of Sim[®]-Lac used was according to manufacturer's specifications. Inclusion level of prebiotics (AXOS and MOS) used in the current study was based on previous studies (Bovera *et al.*, 2010) while therapeutic dosage of

oxytetracycline was used as a positive control. The additives were added to the basal diet and thoroughly mixed to formulate the experimental diets.

117

Table 1 Percentage Composition of Broiler Starter diets and essential proximate in Broiler Starter diets (0 -4 weeks)

Ingredients %	Experimental		Diets		
	1	2	3	4	5
Maize	46.00	46.00	46.00	46.00	46.00
SBM	19.00	19.00	19.00	19.00	19.00
GNC	15.00	15.00	15.00	15.00	15.00
Fish meal (72%)	2.00	2.00	2.00	2.00	2.00
Wheat offal	11.95	11.95	11.95	11.95	11.95
Bone meal	2.00	2.00	2.00	2.00	2.00
Oyster shell	3.00	3.00	3.00	3.00	3.00
Salt	0.25	0.25	0.25	0.25	0.25
*Premix	0.25	0.25	0.25	0.25	0.25
Methionine	0.30	0.30	0.30	0.30	0.30
Lysine	0.25	0.25	0.25	0.25	0.25
¹ .oxytetracycline	-	+	-	-	-
² MOS	-	-	+	-	-
³ .AXOS	-	-	-	+	-
⁴ . Sim [®] lac	-	-	-	-	+
Total	100.00	100.00	100.00	100.00	100.00
Determined Analysis					
Metabolizable energy	11.83	11.83	11.83	11.83	11.83
Crude protein %	23.22	23.22	23.22	23.22	23.22
Crude fibre %	4.88	4.88	4.88	4.88	4.88
Ether extract %	5.63	5.63	5.63	5.63	5.63
Available Ca %	1.97	1.97	1.97	1.97	1.97
Available P %	0.44	0.44	0.44	0.44	0.44
Lysine %	1.34	1.34	1.34	1.34	1.34
Methionine %	0.65	0.65	0.65	0.65	0.65

*Vitamin and mineral premix based on 2.5kg/ ton [Vit A; 4000000 Iu, Vit D:800000, Vit B12: 25mg, Niacin:60000mg, Vit E 40000, Viut k3 800mg, Vit B3 1000mg, Vit B26000mg, Vit B6 5000mg, Panthotenic Acid: 20000, Folic Acid: 200mg, Biotine 8mg, Maganese:300000mg, Iron 80000mg, Zinc: 20000mg, Copper: nill, Cobalt: 80mg, Iodine: 400mg, Selenium:40mg,Choline:800000mg]

124

Table 2 Percentage Composition of Broiler Finisher Diets (4 -8 weeks)

Ingredients %	Experimental		Diets		
	1	2	3	4	5
Maize	50.00	50.00	50.00	50.00	50.00
SBM	12.00	12.00	12.00	12.00	12.00
GNC	11.00	11.00	11.00	11.00	11.00
Fish meal (72%)	2.00	2.00	2.00	2.00	2.00
Wheat offal	19.00	19.00	19.00	19.00	19.00
Bone meal	2.00	2.00	2.00	2.00	2.00

Comment [FB6]: With which nutrient recommendation guide and with which diets formulation software program?

Oyster shell	3.00	3.00	3.00	3.00	3.00
Methionine	0.25	0.25	0.25	0.25	0.25
Lysine	0.25	0.25	0.25	0.25	0.25
*Premix	0.25	0.25	0.25	0.25	0.25
Salt	0.25	0.25	0.25	0.25	0.25
¹ .Oxytetracycline	-	+	-	-	-
² .MOS	-	-	+	-	-
³ .AXOS	-	-	-	+	-
⁴ .Sim® lac	-	-	-	-	+
Total	100.00	100.00	100.00	100.00	100.00
Determined					
Metabolizable	12.43	12.43	12.43	12.43	12.43
Crude protein %	19.90	19.90	19.90	19.90	19.90
Crude fibre %	5.66	5.66	5.66	5.66	5.66
Ether extract %	4.81	4.81	4.81	4.81	4.81
Available Ca %	1.97	1.97	1.97	1.97	1.97
Available P %	0.44	0.44	0.44	0.44	0.44
Lysine %	1.13	1.13	1.13	1.13	1.13
Methionine %	0.56	0.56	0.56	0.56	0.56

- Vitamin and mineral premix based on 2.5kg/ ton

Vit A: 4000000 Iu, Vit D:800000, Vit B12: 25mg, Niacin:60000mg, Vit E 40000, Viut k3 800mg, Vit B3 1000mg, Vit B2 6000mg, Vit B6 5000mg, Panthotenic Acid: 20000, Folic Acid: 200mg, Biotine 8mg, Maganese:300000mg, Iron 80000mg, Zinc: 20000mg, Copper: nill, Cobalt: 80mg, Iodine: 400mg, Selenium: 40mg, Choline: 800000mg

Determination of haematological Parameters: At 28 days, blood samples (2.5 ml each) were collected from the brachial wing vein of four birds per pen (n = 16 per treatment) into vials containing Ethylene diamine tetra-acetate (EDTA) for the determination of haematological indices while another set of blood was collected into heparinised tubes for serum biochemical determination. Haemoglobin concentration (Hb) was estimated using the cyanmethaemoglobin method (Cannan 1958). Packed cell volume (PCV) of blood samples were determined in a Wintrobehaematocrit tube according to the method of Schalm et al. (1975) while total red blood cell and white blood cell count (WBC) were determined using standard method. Differential leucocyte counts (heterophils, lymphocytes, eosinophils and monocytes) were carried out on blood smears stained with May-Grunwald-Giemsa stain and further calculated.

Estimation of serum biochemistry parameters: Total serum protein, albumin and globulin were determined using bromocresol purple method (Varley et al., 1980). Serum creatinine and uric acid concentration was determined according to standard procedures. Serum enzymes (alanine transaminase (ALT) and aspartate transaminase (AST) were analysed spectrophotometrically using Randox^R diagnostic kit. The serum cholesterol was estimated (Qualigens India. Pvt. Ltd., Catalogue number 72201-04)

Salmonella bacteria estimation and identification: Intestinal content of the birds gut was aseptically collected into sterile plain bottle and total salmonella load was estimated using pour plate method described by Hedges (2012). Salmonella colonies were counted and the

Comment [FB7]: Ethical approval for the research was not informed in the manuscript. The authors must include the Protocol number of the Ethics Committee of Animal Use or explain why this information wasn't include in this paper, since the authors used broiler for this experiment

results were expressed according to plate count proceedings established in Normative Instruction (Brazil, 2003). Salmonella bacteria isolation was carried out according to modified method of from Desmidt *et al* (1998). The intestinal sample, 1 gramme of different probiotic, prebiotic fed bird and control was pre-enriched at 37°C for 1 hour in 2 ml BPW (Buffer Peptone Water) and later directly inoculated onto XLD agar (Xylose Lysine Deoxycholate agar). Bacteria colonies found were examined for their respective colonial morphology and biochemically characterized for Salmonella identification while antigenic serological assay was also performed.

Statistical Analysis: The experimental layout was 3 x 5 factorial arrangements within a randomized complete design. Data obtained were subjected to analysis of variance using SAS (1998). Significant means among variables were separated using Duncan Multiple Range Test as outlined by Steel and Torrie (1990). The colony count data were transformed into log base 10 and were subjected to one-way analysis of variance and those with statistical differences were submitted to the Tukey's test at the 0.05% level using Stat view for windows 5.0 (SAS Inc, Cary, NC, USA)

RESULTS

Comment [FB8]: Enough

The result of the main effect of hatchery source and inclusion of various feed additives on the haematological indices and serum biochemistry of starting broiler is presented in Table 3. Result revealed that Hb, RBC, total serum protein, serum albumin, uric acid and creatinine was significantly influenced ($P<0.05$) with highest values recorded for hatchery in Location 3. The WBC, Glucose, serum globulin, uric acid and cholesterol were significantly ($P<0.05$) reduced by the various additives included. However, dietary inclusion of various additives showed no significant effect ($P>0.05$) on the PCV, Hb, RBC, total serum protein, serum albumin and creatinine, respectively. The WBC also reduced significantly ($P<0.05$) across the treatments all the hatcheries with the control diet having higher ($P<0.05$) value of $31,456 \times 10^3/\text{mm}^3$. Serum cholesterol reduced ($P<0.05$) with birds fed diet supplemented with MOS and Sim Lac having the least ($P<0.05$) values of 212.7 and 214.2 mg/dl respectively.

The interaction effect of hatchery source and feed additives on the haematological indices and serum biochemistry of starting broiler is compared and shown in Table 4. The results revealed a significant ($P<0.05$) interaction effect on PCV, Hb, WBC count, red blood cells, total serum protein, serum albumin and uric acid. Total serum protein showed higher ($P<0.05$) values for birds sourced from location 1 hatchery and fed diet supplemented with Sim[®] Lac. However, there was a significant ($P<0.05$) increase in uric acid of birds sourced from hatcheries in location 3 and fed with the control diet. Glucose, serum globulin, cholesterol and creatinine showed no significant ($P>0.05$) interaction effect.

The result of the main effect of hatchery sources and dietary inclusion of feed additives on haematological indices and serum biochemistry of finisher broilers is presented in Table 5. Hb and RBC counts were significantly ($P<0.05$) increased with birds sourced from hatcheries in location 1 and 2 while birds from hatcheries in location 3 had the highest values. The

191 serum glucose, serum total protein, serum albumin, serum globulin uric acid and creatinine
192 levels of the broilers significantly ($P<0.05$) affected by the location of the hatchery. Birds fed
193 with the control showed higher ($P<0.05$) values of serum albumin and uric acid. The control
194 diet and antibiotics had statistically similar values for white blood cell, serum albumin and
195 serum cholesterol. Other parameters measured, PCV, Hb, Red blood Cell, total protein and
196 creatinine across the treatment were not significantly ($P>0.05$) affected. This result followed
197 the same trend with the result of starting broilers.

198 The interaction effect of hatchery source and additives on the haematological indices and
199 serum biochemistry of finishing broiler is presented in table 5. The result showed that packed
200 cell volume, haemoglobin and white blood cells were significantly ($P<0.05$) increased in the
201 control diet across the locations compared to other dietary treatments applied. The control
202 diets had 50.5% PCV and 33,743 of white blood cells. Higher significant ($P<0.05$) value for
203 red blood cells was recorded for antibiotic and AXOS at hatchery location 3. Total serum
204 protein recorded a significant ($P<0.05$) increase with increasing AXOS, Sim lac[®] which were
205 statistically similar to antibiotic. The uric acid recorded significant ($P<0.05$) reduction across
206 the location having the control diet with higher values of 18.0 and 14 mg/dl respectively. The
207 serum globulin, cholesterol and creatinine were not significantly ($P>0.05$) affected by
208 interaction of location of hatchery and feed additives.

209 Table 7 showed a significant ($P<0.05$) reduction in Salmonella count at day 1 of the study
210 when various additives were added. Main effect of location showed no effect on gut
211 salmonella count of the birds. At day 28 and 56, inclusion of MOS showed the least
212 salmonella count. Similar salmonella counts were obtained at day 28 for birds fed diet
213 supplemented with antibiotics and AXOS. In all the phases of the study, gut salmonella
214 reduced drastically with inclusion of various additives. Birds fed diet supplemented with
215 MOS showed the least gut salmonella count.

216 **Table 3 Main effect of location of hatchery and feed additives on blood parameters of broiler chickens (0-**
217 **4weeks)**
218

Parameters	Location of Hatchery				Additives					
	1	2	3	SEM	Control	AB	MOS	AXOS	SIML	SEM
PCV (%)	31.9	32.7	35.9	1.55	34.8	31.1	31.1	33.0	33.8	1.99
Hb(g/dl)	10.6 ^b	11.0 ^b	14.6 ^a	0.50	12.6	12.2	11.3	12.4	12.0	0.65
WBC ($10^3/\text{mm}^3$)	30.69	30.27	30.69	27.01	31.46 ^a	30.53 ^{bc}	31.48 ^{ab}	30.25 ^c	29.37 ^d	3.49
RBC ($\times 10^{12}/\text{L}$)	2.7 ^b	2.7 ^b	2.8 ^a	0.11	4.1	4.3	3.8	4.1	3.9	0.15
Glucose (mg/dl)	137.1 ^c	154.4 ^a	141.1 ^b	3.84	123.5 ^d	151.2 ^b	141.2 ^{bc}	168.6 ^a	136.4 ^c	4.96
T. Protein(g/l)	41.7 ^b	39.2 ^c	43.7 ^a	1.18	38.4	43.0	41.4	43.5	41.5	1.5
Serum Globulin(g/l)	14.3 ^b	17.6 ^a	18.3 ^a	0.56	18.3	18.6	14.1	16.9	15.8	0.73
Serum Albumin (g/l)	27.4 ^a	21.6 ^c	25.4 ^b	1.02	20.1 ^c	24.4 ^b	27.3 ^a	26.6 ^a	25.7 ^{ab}	1.31

Uric Acid (mg/dl)	9.9 ^b	7.9 ^c	13.9 ^a	0.21	11.8 ^a	10.5 ^b	10.5 ^b	9.6 ^c	10.1 ^{bc}	0.27
Cholesterol (mg/dl)	240.7	232.0	244.9	11.61	252.8 ^b	287.2 ^a	212.7 ^c	228.8 ^{bc}	214.2 ^c	14.99
Creatinine(mg/dl)	0.9 ^c	1.5 ^b	4.9 ^a	0.30	2.2	2.1	2.3	2.1	3.2	0.4
219	abc means on the same row having different superscript are significantly different (P<0.05); Control=No Additives; AB=									
220	Antibiotics (oxytetracycline); MOS=Mannose oligosaccharide; AXOS=Arabinoxylans Oligosaccharide; Sim									
221	lac= <i>Pediococcus acidilactici</i>									
222										

223 Table 4 : Interaction effect of location of hatchery and additives on blood parameter of starting broiler

224

Parameters	Location 1					Location 2					Location 3					
	Control	AB	MOS	AXOS	SIML	Control	AB	MOS	AXOS	SIML	Control	AB	MOS	AXOS	SIML	SEM
PCV (%)	27.5 ^g	36.5 ^c	28.5 ^{lg}	36.0 ^c	31.0 ^{ef}	45.5 ^a	27.5 ^g	32.3 ^{de}	23.0 ^h	35.3 ^c	31.5 ^d	40.5 ^b	32.5 ^{bc}	40.0 ^b	35.0 ^{cd}	1.46
Hb(g/dl)	9.1 ⁱ	11.8 ^e	9.5 ⁱ	12.7 ^{cd}	10.1 ^{ef}	15.5 ^b	9.0 ⁱ	11.0 ^{ef}	7.7 ^j	12.0 ^d	13.1 ^c	15.8 ^b	13.5 ^c	16.7 ^a	14.1 ^{bc}	0.68
WBC (10 ³ /mm ³)	33.733 ^a	30.200 ^e	30.867 ^c	28.578 ^g	30.075 ^f	26.900 ^h	31.175 ^d	31.708 ^{bc}	33.60 ^b	27.98 ^{gh}	33.737 ^a	30.204 ^e	30.870 ^c	28.582 ^g	30.079 ^f	5.1649
RBC (x 10 ¹² /L)	2.4 ⁱ	3.2 ^d	2.3 ⁱ	3.1 ^d	2.6 ^{ef}	3.6 ^c	2.4 ⁱ	2.8 ^c	2.2 ⁱ	2.7 ^e	4.4 ^b	5.3 ^a	4.3 ^b	5.1 ^a	4.6 ^{ab}	0.27
Glucose (mg /dl)	115.3	140.3	127.1	171.7	130.9	136.0	169.1	165.3	158.3	143.3	119.3	144.3	131.1	175.7	134.9	4.85
T. Protein(g/l)	38.7 ⁱ	45 ^{cd} .0	42.5 ^e	46 ^c .o	36.5 ^j	35.8 ^k	37.0 ^{is}	37.1 ^{is}	36.4 ^j	49.4 ^h	40.7 ⁱ	47.0 ^b	44.5 ^d	48.0 ^{ab}	38.5 ^j	1.19
Globulin(g/l)	17.4	17.9	9.2	15.5	11.5	16.0	16.1	19.8	15.7	20.3	21.4	21.9	13.2	19.5	15.5	0.90
Albumin (g/l)	21.3 ^{gh}	27.1 ^e	33.3 ^a	30.5 ^b	24.9 ^j	19.8 ⁱ	20.9 ^h	17.3 ^j	20.6 ^h	29.2 ^{cd}	19.3 ⁱ	25.1 ^f	31.3 ^c	28.5 ^d	22.9 ^g	1.24
Uric Acid (mg/dl)	12.0 ^d	9.8 ^e	9.4 ^c	8.7 ^f	9.4 ^e	7.2 ^j	8.3 ^h	8.7 ^f	7.6 ⁱ	7.5 ⁱ	16.0 ^a	13.4 ^b	13.4 ^b	12.7 ^c	13.4 ^b	5.68
Cholesterol (mg/dl)	267.8	319.9	203.0	218.5	194.4	218.9	217.7	228.1	245.5	249.8	271.8	323.9	207.0	222.5	198.4	10.22
Creatinine(mg/dl)	0.9	0.8	1.0	0.8	0.9	1.0	0.9	1.0	0.8	1.7	2.8	2.8	3.0	2.8	2.9	0.24

225

226 ^{abc} means on the same row having different superscript are significantly different (P<0.05); Control=No Additives; AB= Antibiotics (oxytetracycline); MOS=Mannose

227 oligosaccharide; AXOS=ArabinoxylansOligoscharide; Sim lac=*Pediococcusacidilactici*

228 **Table 5 Main effect of Location and feed additives in blood parameters of broiler(4-8weeks)**

Parameters	Location of hatchery				Additives					
	1	2	3	SEM	Control	AB	MOS	AXOS	SIML	SEM
PCV (%)	36.9	37.7	40.9	1.00	39.8	36.1	36.1	38.0	38.8	0.66
Hb(g/dl)	12.6 ^c	14.0 ^b	16.6 ^a	1.04	14.6	14.2	13.3	14.4	14.0	0.20
WBC (10 ³ /mm ³)	30.700	30.281	30.704	2.70	31.466 ^a	30.5367 ^{ab}	31.159 ^{ab}	30.264 ^{bc}	29.386 ^c	3.25
RBC (x 10 ¹² /L)	3.7 ^b	3.7 ^b	5.7 ^a	0.54	5.1	5.3	4.8	5.1	4.9	0.08
Glucose (mg /dl)	140.1 ^c	154.4 ^a	141.1 ^b	3.77	123.5 ^d	161.2 ^b	141.2 ^{bc}	168.6 _a	136.4 ^c	7.37
T. Protein(g/l)	36.7 ^{ab}	34.2 ^b	38.7 ^a	1.06	33.4	38.0	36.4	38.5	36.5	0.79
Serum Globulin(g/l)	10.3 ^b	13.6 ^a	14.3 ^a	1.01	14.3 ^a	14.6 ^a	10.1 ^c	12.9 ^{ab}	11.8 ^{bc}	0.74
Serum Albumin (g/l)	26.4 ^a	20.6 ^b	24.4 ^a	1.39	19.1 ^b	23.4 ^a	26.3 ^a	25.6 ^a	24.7 ^a	1.14
Uric Acid (mg/dl)	11.9 ^b	9.9 ^c	12.9 ^a	0.72	13.8 ^a	12.6 ^b	12.5 ^b	11.6 ^c	12.1 ^{bc}	0.33
Cholesterol (mg/dl)	230.7	222.0	234.7	3.06	242.8 ^{ab}	277.2 ^a	202.7 ^b	218.8 ^b	204.2 ^b	12.51
Creatinine(mg/dl)	1.7 ^b	1.3 ^b	2.7 ^a	0.34	2.0	2.9	2.1	1.9	2.9	0.19

229 ^{abc} means on the same row having different superscript are significantly different (P<0.05); **Control=No**
230 **Additives; AB= Antibiotics (oxytetracycline); MOS=Mannose oligosaccharide; AXOS=Arabinoxylans**
231 **Oligosaccharide; Sim lac=*Pediococcus acidilactici***

232

233

234

Table 6: Interaction effect of Location of hatchery and additive on blood parameter of finishing broiler.

	Location 1					Location 2					Location 3					
Parameter	Control	AB	MOS	AXO S	SIML	Contr ol	AB	MOS	AXO S	SIML	Contr ol	AB	MOS	AXO S	SIML	SEM
PCV (%)	32.5 ^l	41.5 ^c	33.5 ^f	41.0 ^c	36.0 ^e	50.5 ^a	32.5 ^l	37.3 ^d	28.0 ^j	40.3 ^c	36.5 ^e	45.5 ^b	37.5 ^d	45.0 ^b	40.0 ^c	1.46
Hb(g/dl)	11.1 ⁱ	13.8 ^e	11.5 ⁱ	14.7 ^{cd}	12.1 ^{ef}	17.5 ^b	11.0 ⁱ	13.0 ^{ef}	9.7 ^j	14.0 ^d	15.1 ^c	17.8 ^b	15.5 ^c	18.7 ^a	16.1 ^{bc}	0.68
WBC (103/mm3)	33.743 ^a	30.210 ^c	30.876 ^e	28.58 ^g	30.08 ^{5^f}	26.91 ^{0^h}	31.18 ^{5^d}	31.71 ^{8^{bc}}	33.61 ^{0^b}	27.98 ^{5^{gh}}	33.74 ^{7^a}	30.21 ^{4^c}	3.088 ^{0^c}	28.59 ^{2^g}	30.08 ^{9^f}	5.17.
RBC (x 1012/L)	3.4 ^f	4.3 ^d	3.3 ^l	4.1 ^d	3.6 ^{ef}	4.6 ^c	3.4 ^f	3.8 ^e	3.2 ⁱ	3.7 ^e	7.4 ^b	8.3 ^a	7.3 ^b	8.1 ^a	7.6a ^b	2.49
Glucose (mg /dl)	115.3	140.3	127.1	171.7	630.9	136.	169.1	165.3	158.3	143.3	119.3	144.3	131.1	175.7	134.9	31.67
T. Protein(g/l)	33.7 ^f	40 ^{bc}	37.5 ^d	41.0 ^b	31.5 ^j	30.8 ^{ij}	32.0 ⁱ	32.1 ⁱ	31.4 ^j	44.4 ^a	35.7 ^e	42.0 ^{ab}	40.5 ^c	43.0 ^a	33.5 ^f	1.20
Globulin(g/l)	13.4	13.9	52	11.5	9.5	12.0	12.1	15.8	11.7	16.2	17.4	17.9	10.2	15.5	11.6	8.17
Albumin (g/l)	20.3 ^e	26.1 ^c	32.3 ^a	29.5 ^b	22.0 ^d	18.8 ⁱ	19.9 ^f	16.3 ^j	19.6 ^f	28.2 ^{bc}	18.3 ⁱ	24.1 ^d	30.3 ^{ab}	27.5 ^{bc}	21.9 ^{dc}	1.39
Uric Acid (mg/dl)	14.0 ^d	11.8 ^e	11.4 ^e	10.7 ^f	11.4 ^e	9.2 ⁱ	10.3 ^{fi}	10.7 ^f	9.6 ⁱ	9.5 ⁱ	18.0 ^a	15.8 ^b	15.4 ^b	14.7 ^c	15.4 ^b	0.69
Cholesterol (mg/dl)	257.8	309.9	193.0	208.5	184.4	208.9	207.7	218.1	235.5	239	261	313.9	197.0	212.5	188.4	10.20
Creatinine(mg/dl)	0.6	0.6	0.8	0.6	0.7	0.8	0.7	0.8	0.6	0.5	0.6	0.6	0.8	0.6	0.7	0.02

^{abc} means on the same row having different superscript are significantly different (P<0.05); **Control=No Additives; AB= Antibiotics (oxytetracycline); MOS=Mannose oligosaccharide; AXOS=Arabinoxylans Oligosaccharide; Sim lac=Pediococcus acidilactici**

Table 7 Main effect of feed additives on Salmonella count in the gut of broiler chicken CFUX10⁶

Main effect of Location					Main effect of additives					
Days	1	2	3	SEM	Control	AB	MOS	AXOS	SIM L	SEM
Day 1	0.24	0.31	0.28	0.02	0.43 ^a	0.31 ^b	0.30 ^b	0.29 ^b	0.31 ^b	0.10
Day 28	0.21	0.20	0.12	0.04	0.59 ^a	0.11 ^c	0.04 ^d	0.13 ^c	0.44 ^b	0.15
Day 56	0.15	0.17	0.07	0.02	2.81 ^a	0.18 ^b	0.02 ^d	0.11 ^c	0.16 ^b	0.09

^{abc} means on the same row having different superscript are significantly different (P<0.05); **Control=No Additives; AB= Antibiotics (oxytetracycline); MOS=Mannose oligosaccharide; AXOS=Arabinosylans Oligosaccharide; Sim lac=*Pediococcus acidilactici***

Discussion

Several species of *Salmonellae* infections are now recognized as important food borne diseases contracted majorly from Salmonella contaminated or pre-infected poultry product. In many cases, salmonella infection in broilers or layers sometimes remain sub-clinical, but poultry products are considered as important sources for human food borne outbreaks (Bielke *et al.*, 2003). In the present study, the packed cell volume (PCV) value obtained lied within the normal range and similar to other findings for avians (Oyewole and Ajibade, 1990; Banerjee *et al.*, 2002). The red blood cell counts (RBC) and PCV also lied within the normal range reported for chickens while highest hemoglobin level (Hb), RBC, total serum protein and serum globulin obtained for broiler chicks sourced from hatcheries in location 3 showed improved health status of the birds. Low Hb concentrations values have been reported to be indications of poor protein intake and severe parasitic infection and damage to liver (Lindsay, 1997, Corcionivoschi *et al* 2010). Dietary treatment have been reported to influence the concentration of both red blood cell and packed cell volume (Banerjee *et al.*, 2002). This implied that birds sourced from hatcheries in location 3 efficiently utilize the feed offered. The high serum glucose obtained for birds fed with various additives showed improved energy utilization and increased serum glucose which implies high or increased energy availability. All birds at the starter phase fed diet supplemented with various additives showed reduced serum uric acid concentration which usually indicate inefficient protein utilization, resulting in wastages and excess protein in form of uric acid. This agreed with the findings of Szabo *et al.* (2005) who reported high levels of serum uric acid due to poor protein utilization. There is significant reduction in serum cholesterol in broiler chicks fed diet supplemented with different additives. Prebiotic and probiotic supplement used in this study has been shown to reduce serum cholesterol concentration in chicken (Kanna *et*

Comment [FB9]: More in-depth discussion is required.
Discussion section is not enough to properly explain reasons of the results obtained

263 *al.*, 2005). Reduction in serum cholesterol observed in this study for broiler fed with probiotic well corroborate the findings of Pereira
 264 and Gibson (2002) who reported that oral administration of probiotic significantly reduced serum cholesterol level by as much as 22 to
 265 33%. The ability of probiotic to lower serum cholesterol is usually due to inhibition of intestinal cholesterol absorption and
 266 suppression of bile acid re—absorption mechanism (Guo *et al.*; 2012)

267 At the finisher phase, birds sourced from hatchery 3 recorded the highest Hb, RBC and total serum protein. This followed similar
 268 trend with what was observed at the starter phase. This is suggestive of the fact that birds sourced from hatchery 3 showed a better
 269 healthy status than other birds sourced from the other hatcheries. Low total protein levels could suggest a liver or kidney
 270 disorder which could be caused by poor protein metabolism due to poor digestion or absorption. It was observed in the present study
 271 that dietary feed additives supplementation influenced serum protein and albumin concentrations, with highest serum protein and
 272 albumin concentrations in the additives compare with the control. Several authors have identified that probiotic supplementation in
 273 diets increased serum protein and albumin concentrations, in broilers (Agawane and Lonkar, 2004, Capcarova *et al.*, 2011), piglets
 274 (Harding *et al.*, 2008) and fish (Zhou *et al.*, 2010). Harding *et al.*, (2008) reported that probiotics did not stimulate gastrointestinal
 275 protein synthesis or reduce the severity of intestinal colitis, but reported an unidentified signaling mechanism between the gut and
 276 liver which could be responsible for the increase in liver protein and plasma protein synthesis induced by probiotics. Furthermore,
 277 dietary probiotics and prebiotics supplementation may increase the serum protein and albumin concentrations by increasing iron
 278 absorption in the lower intestine (Eizaguirre *et al.*, 2002). The increased serum protein concentration in broilers sourced from hatchery
 279 3 could be due to improvement in protein synthesis and carbohydrate and lipid metabolism (Rosa *et al.*, 2001). Reduced WBC noticed
 280 with finishing broilers fed diet supplemented with AXOS and SIML is suggestive of absence/low population of foreign bodies or
 281 pathogens introduced in the system of the birds resulting in reduced production of antibodies to fight against such. White blood cells
 282 (WBC) played prominent role in disease resistance especially with respect to the generations of antibodies and the process of
 283 phagocytosis. An elevated value of WBC is an indication that the birds were reacting to one or more factors in the feed (Oduguwa,
 284 2006).

285 Dietary inclusion of various additives supplemented in the diets showed reduced serum uric acid at both the starting and finishing
 286 phases of the birds. This is implicative of good health status. There exist an indirect relationship between serum uric acid
 287 concentration and efficiency of utilization of ingested protein (Szabo *et al.* 2005). Although birds fed with various additives showed
 288 reduced gut salmonella counts, dietary supplementation of MOS recorded the least gut salmonella count at both the starter and grower
 289 phases of the study. This could be attributed to binding effect of mannose oligosaccharides on the intestinal epithelium of the birds.
 290 Most pathogenic microbes including salmonella attach to mannose –containing cells in the intestinal tract. Studies on MOS pointed

out that the oligosaccharides is able to bind to mannose-specific lectin of gram-negative pathogens that express Type 1- fimbriae such as Salmonella, preventing salmonella organisms from proliferation hence resulting in their excretion from the intestine (Baurhoo *et al.*, 2007). This resulted in the reduction noticed in the gut salmonella count of broilers fed with MOS in this study.

Conclusion

Although broiler chicks obtained from source 3 showed prospects for improved haematological indices, dietary inclusion of MOS and AXOS in ration for broilers sourced from salmonella infected hatcheries showed improved haematological indices, reduced serum cholesterol and gut salmonella count. Therefore, AXOS and MOS are beneficial for improved healthy status of broiler and prevention of Salmonella morbidity and mortality in poultry.

References

- Agawane, S.B., Lonkar, P.S., 2004. Effect of probiotic containing *Saccharomyces boulardii* on experimental ochratoxicosis in broilers: Hematobiochemical studies. *J. Vet. Sci.* 5:359-367.
- Andremont, A 2000. Consequences of antibiotic therapy to the intestinal ecosystem, *Ann. Fr. Anesth.Reanim.* 19,395-402.
- Baurhoo, B., Letellier, A., Zhao, X., Ruiz-Feria, C.A., 2007. Cecal populations of *Lactobacilli* and *Bifidobacteria* and *Escherichia coli* after in vivo *Escherichia coli* challenge in birds fed diets with purified lignin or mannanoligo-saccharides. *Poultry.Science*. 86, 2509–2516.
- Banerjee, S. K., Patra, B. C. and Bandoypathay. P, A. (2002). Change of blood parameters in Carp, *Catlacatta*. *J. Aquatic Biology*. 17(11): 79-84
- Bovera, F., Nizza, S., Marono, S., Maliardo, K., Piccolo, G., Tudisco, R., de Martino, L, Nizza, A., 2010. Effect of mannan-oligosaccharides on rabbit performance, digestibility and rectal bacterial anaerobic populations during an episode of epizootic rabbit enteropathy. *World Rabbit Science*., 18, 9–16.
- Brasil.Ministerio da Agricultura, Abastecimento e Pecuaria. Instrucao normative, SDA N 62, de 26 agosto de 2003.Diario Oficial da Uniao,18 set. 2003; Secao 1 P.14
- Capcarova, M., Hascik, P., Kolesarova, A., Kacaniova, M., Mihok, M., Pal, G., 2011. The effect of selected microbial strains on internal milieu of broiler chickens after peroral administration. *Res. Vet. Sci.* 91(????):132-137

Comment [FB10]: The suggested reference must be supplemented to the reference section

And
References have not been written according to Guide for Authors of The ARRB

Comment [FB11]: Uniformity in references must be seen.
Poult. Sci

Comment [FB12]: Uniformity in references must be seen.

- 315 Corcionivoschi, K., Dan, D., Ioan, M., Deirdre, S., Lavinia, S., Calm, J. and Bairke 2010. Effect of probiotic on animal health, Scientific
316 Papers Animal Science and Biotechnologies, 43(1), 35-41.
- 317 Desmidt, M., Ducatelle R., Haesebrouck, F. 1998. Serological and bacteriological observations on experimental infection with
318 Salmonella Hadar in chickens. Veterinary Microbiology 1998; 60:259-269.
- 319 Eizaguirre, I., Urkia, N.G., Asensio, A.B., Zubillaga, I., Zubillaga, P., Vidales, C., Garcia-renzana, J.M., Aldazabal, P., 2002.
320 Probiotic supplementation reduces the risk of bacterial translocation in experimental short bowel syndrome. J. Pediatr. Surg. 37:699-
321 702.
- 322 Fuller, R. (1989) Probiotics in man and animals. Journal of Applied Bacteriology 66(????):
323 365-378
- 324 Gibson, G. R., Probert, H. M., Van Loo, J., Rastall, R. A., Roberfroid, M. B., 2004. Dietary modulation of the human colonic
325 microbiota: Updating the concept of prebiotics. Nutrition Research Review. 17, 259–275.
- 326 Hafez, H. M. and Jodas, S. (2000). Salmonella Infection in Turkey. In: Salmonella in domestic animals, Ed.
- 327 Guo Z, Xiao ML, Qiu-Xiang Z, Feng-Wel T, Hao Z, He-Ping Z, Wei C (2012). research advances with regards to clinical outcome
and potential mechanisms of the cholesterol-lowering effects of probiotics. Clin. Lipidology col, 5: 501-507
- 328 Harding, S.V., Fraser, K.G., Wykes, L.J., 2008. Probiotics stimulate liver and plasma protein synthesis in piglets with dextran sulfate-
329 induced colitis and macronutrient restriction. J. Nutr. 138:2129-2135
- 330 Kanna, M., Karamakaron, R., Balakrishnan, V., Prabhakar T.G 2005. Influence of prebiotics supplementation on Lipid profile of broilers
331 Int. J. Poultry Sci. 4(12): 994-997.
- 332 Lindsay, D. B. (1997). The effect of feeding patterns and sampling on By D. Lister. PP. 99-120.
- 333 Huyghebaert, G., Ducatelle, R., Van Immerseel, 2011. An update on alternatives to antimicrobial growth promoters for
334 broilers. Veterinary Journal. 187, 182–188
- 335
- 336 Lindsay, D. B. (1997). The effect of feeding patterns and sampling on By D. Lister. PP. 99-120.
- 337 Mountzouris, K. C., Tsirtsikos, P., Kalamara, E., Nitsch, S., Schatzmayr, G., Fegeros, K., 2007. Evaluation of the efficacy of a
338 probiotic containing lactobacillus, bifidobacterium, enterococcus, and pediococcus strains in promoting broiler performance and
339 modulating cecal microflora composition and metabolic activities. Poultry Science. 86, 309-317.

Comment [FB13]: Uniformity in references must be seen.

43(1): 35-41.

Comment [FB14]: Uniformity in references must be seen.

Comment [FB15]: Uniformity in references must be seen.

Comment [FB16]: Uniformity in references must be seen.

- 340 Mohan, B., Kadirve, R., Natrajan, A. and Bhaskaran, M. (1996).Effect of probiotic supplementation on growth, nitrogen utilization
341 and serum cholesterol in broilers.*British.Poultry.Science.*, 37: 395-401.
- 342 Mayrhofer, S., Paulsen, P., Smulders, F. J. M. & Hilbert, F. (2003). Antimicrobial resistance profile of five major food-borne
343 pathogens isolated from beef, *pork and poultry*. *Inter. J. Food Microbiol.*, 97 (1), 23–29.
- 344 Murugkar, H.V., Rahman, H. & Dutta, P. K. (2005). Distribution of virulence genes in *Salmonella* serovars isolated from man and
345 animals. *Indian Med. Res.*, 177, 66–70.
- 346 Oduguwa, O. O. (2006). Utilization of whole pods of Albiziasamon in diets of growing rabbits. *Nigerian Journal Animal Production*
347 33 (2): 197-202.
- 348 Ojewola, J. O. and Ajibade, H. A. (1990). The osmotic fragility of erythrocyte of turkey of two age group. *Veterinary.Arhives*. 60: 91-
349 100.
- 350 Olorunsola, R.A Erubetine, D Oyekunle, A.M Jegede, A. V and Ogunade, I M. 2012. Salmonella organisms transmission in
351 hatching broiler eggs, *journal of Biology, Agriculture and Healthcare*. The international institute for science, technology and
352 education ISSN 2224-330 (Paper), ISSN 2225-093X (online) Vol. 2 No 10
- 353 Pereira, D. I., Gibson, G. R. (2002).Effect of consumption of probiotics and prebiotics on serum lipid levels in humans.*Crit. Rev.*
354 *Biochem. Mol. Bio*. 37: 259-281.
- 355 Rolfe, R.D 2000. The role of probiotics cultures in the control of gastrointestinal health, *The journal of Nutrition*, 130, 396-402
- 356 Rosa, C. A., Maizzo, R., Magnoli, C., Salvano, M., Chiac, S. M., Ferrero, S., Saenz, M., Carvalho, E. C. and Dalcero, A. 2001.
357 Evaluation of the efficacy of bentonite from the south of Argentina to ameliorate the toxic effect of aflatoxin in
358 broilers.*Poultry.Science.*, 80: 139-144.
- 359 Rosen, G. (1995). Antibacteria in poultry and pig nutrition. In *Biotechnology in Animal Feeds and Animal Feeding*. R. J. Wallace,
360 and A. Chesson, ed. VCH, New York. Pp. 143 – 172.
- 361 Revolledo, L., Ferreira, C. S. A. & Ferreira, A. J. P. (2009).Prevention of *Salmonella* Typhimurium colonization and organ invasion by
362 combination treatment in broiler chicks. *Poult. Sci.*, 88, 734–743.
- 363 SAS (1998) SAS/STAT® User's Guide (Released 6.08) SAS institute inc.Cary, NorthCaroline,U.S.A.

Comment [FB17]: Why italic????

Comment [FB18]: Why italic????

Comment [FB19]: Uniformity in references must be seen.

Comment [FB20]: Uniformity in references must be seen.

Comment [FB21]: Uniformity in references must be seen.

Comment [FB22]: Uniformity in references must be seen.

Comment [FB23]: Why italic????

- 364 Sherman, P.M., Ossa, J.C., Johnson-Henry, K., 2009. Unravelling mechanisms of action of probiotics. *Nutrition.Clinical.Practice*. 21,
365 10–14.
366
- 367 Silva, P. R., Freitas, O. C., Laventiz, A. C., Junqueira, O. M. and Fagliari, J. J. (2007). Blood serum components and serum protein
368 test of hybro – Pa broiler of different ages. *Rev Bras. Cienc.Avic*. Vol 9 No. 4 Campinas Oct/Dec. (~~2007~~).
- 369 Sleator, R.D and Hill, C 2008. New frontiers in Probiotic research. *Lett_Appl_Microbio* 46:143-
370 147.
- 371 Szabo A. M, Horn, P., Suto Z., Bazara, and Romvari. R. (2005). Developmental dynamics of some blood biochemical parameters in
372 the growing turkey (*meleagrisGallus*). *Acta vet. Hungary* 53 (4): 397-409.
- 373 Tannock, G. W. (1998). The normal microflora; new concepts in health promotion. *Microbiological Science* 5, 4 -8.
- 374 Tannock, G. W. (1998). The normal microflora; new concepts in health promotion. *Microbiological Science* 5, 4 -8. |
- 375 Wray, C. and Wray. A CAB. Publishing ISBW085199261. Pp.133 -155.
- 376 Zhou, X., Wang, Y., Yao, J., Li, W., 2010. Inhibition ability of probiotic, *Lactococcus lactis*, against *A. hydrophila* and study of its
377 immunostimulatory effect in tilapia (*Oreochromis niloticus*). *Int. J. Eng. Sci. Technol.* 2(??):73-80.
- 378

Comment [FB24]: same