

Original Research Article

Effect of *Xylopi aethiopica* Dried Fruits (Grains Of Selim) As Additive on Performance, Microbial Response and Blood Parameters in Finisher Broilers

ABSTRACT

The effect of grains of selim (*Xylopi aethiopica*) as an additive on the performance, gut microbes and blood parameters of finisher broilers was investigated. One hundred and ninety five (195) 28days old broilers (Arbor acres strain) were randomly assigned to five treatment groups with each treatment having three replicates in a completely randomized design. Each treatment group had 39chicks and 13chicks per replicate.. Finely blended grains of selim was administered through drinking water on treatments 2, 3, 4 and 5 at concentrations of 0.4, 0.6, 0.8 and 1.0 g per litre while chicks on treatment 1 (control) received 0.3g/litre of antibiotics (doxygen). Chicks were fed with isoproteinous and isocaloric diets containing 20% crude protein and 2905.95 Kcal/kg metabolizable energy ad libitum for four weeks. Results showed no significant ($P<0.05$) differences among treatments in the final body weight, total weight gain, daily weight gain, total feed intake, daily feed intake and feed conversion ratio. There were no significant ($P>0.05$) differences in the microbial population count of the faeces before and after the birds received treatment of antibiotics and grains of selim in drinking water. There were no significant ($P>0.05$) differences in haematology parameters. However there were differences in serological parameters with creatinine, cholesterol, glucose and urea significantly ($P<0.05$) different while albumin, total protein and globulin were similar ($P>0.05$). Results indicate that grains of selim has antimicrobial and anthelmintic properties, and promotes growth in broiler chickens with no adverse health implications.

Keywords: *anthelmintic; antibiotics; antimicrobial; biochemistry; Xylopi aethiopica dried fruits (grains of selim); growth promoter; serum.*

1.0 INTRODUCTION

Xylopi aethiopica is an aromatic tree growing up to a height of 20m and of the order Magnoliales and family Annonaceae. It grows widely in the tropical zones of west, central and southern Africa [1]. The dried fruits of *Xylopi aethiopica* (Grains of Selim) are used as a spice and an herbal medicine. There is more concern on the residual effect of synthetic antibiotic use in broiler production. The consumers of poultry products are now paying more attention to quality and safety of poultry products that they consume [2]. Reports on residual effects of antibiotics have generated renewed interest in the none use of antibiotics and other drugs in poultry production [3]. It has become imperative that in raising animals for food, a more nutrition-based health strategy must be adopted in the development of animal agriculture [3]. It has been found that some herbs, spices and their extracts stimulate feed intake and endogenous secretions or possess antimicrobial, coccidiostatic or anthelmintic activities [3,4]. Many of these herbal products and their extracts have improved broiler performance, indicating

that they have growth-promoting potentials [4]. It is the opinion of [5] that herbal products could serve as an environmental friendly alternatives to the antibiotic growth promoters still being used in some developing countries. The high cost of antibiotics and other drugs in addition to their residual effects has necessitated the need to research into natural herbal plants that could serve as cheap and good alternative in place of commercial (synthetic) antibiotics. This study is designed to investigate the effects of grains of selim (*Xylopia aethiopica* dried fruits) on performance, microbial response and blood parameters in finisher broilers.

2.0 MATERIALS AND METHODS

2.1 Experimental Site: The experiment was carried out at the Poultry Research Unit of the Department of Animal Science, Delta State University, Asaba Campus, Nigeria.

2.2 Experimental Birds and Management: One hundred and ninety five (195) finisher broilers (Arbor acres strain) weighing from 1377- 1454.67g were used in this study. The birds were managed in a deep litter house of fifteen pens with dimensions of 2.04m x 2.03m and given feed and water *ad libitum* for four weeks (29-56 days).

2.3 Experimental Design: The finisher broilers were randomly assigned into five treatment groups with three replicates on equal weight basis in a completely randomized design. Each treatment group consists of 39 broiler chicks and 13 in each replicate.

2.4 Experimental Diets: Birds were fed the same isoproteinous and isocaloric diet containing 20% crude protein and 2905.95 Kcal/kg metabolizable energy. A well homogenized powdered grains of selim was obtained by pulverizing with mortar and pestle, milled and passed through a standard sieve (2mm) to remove all fibre. Weights of 0.4, 0.6, 0.8 and 1.0g of grains of selim were weighed and dissolved in a litre of water each for more than 10 hours and administered to broilers in treatments 2, 3, 4 and 5 in their drinking water, while broilers in treatment 1 (Control) were given antibiotics in their water (0.3g per litre of water). Well blended powdered grains of selim dissolved in water without sediments. Grains of selim were used to replace antibiotics in the drinking water of finisher broilers in treatments 2, 3, 4 and 5 whenever birds in treatment 1 were given antibiotics. Grains of selim and antibiotics were given in drinking water for five days. The proximate and phytochemical compositions of *Xylopia aethiopica* dried fruits are presented in Table 1. The composition of the finisher broilers diet are presented in Table 2.

2.5 Measurements: The measurements of body weight development were done weekly on replicate basis to obtain weekly body weight and body weight gain. Feed intake and mortality were also recorded on replicate basis weekly while feed conversion ratio which is a ratio of feed consumed and the weight gained over a specific period was calculated. Faecal samples were collected from each replicate for microbial identification and population count before and after administering grains of selim and antibiotics. Fresh faecal droppings were totally collected from each replicate pen before dawn and thoroughly mixed wet on the 6th day after 5 days of giving grains of selim and antibiotics. One gram of the wet faecal samples from each replicates were emulsified in a drop of normal saline on a microscopic slide and covered with a cover slip. The samples were then examined with X10 objective lens and the organisms found were identified with X40 objective lens. 1.0 g of the wet faecal samples were made into liquid suspension with 10ml of normal saline. Serial dilutions ranging from 10^{-1} to 10^{-5} were made for each faecal sample. From these serial diluents, 0.03ml each were inoculated into the media plates of the

Blood Agar (BA) and Nutrient Agar (NA) using the spreader method [6]. Both aerobic and anaerobic culturing was respectively carried out. Inoculated plates were incubated at 37°C for 24-48 hours. The growth of microbial colonies on the plates were counted and the average count from each sample was divided by 0.03ml (the volume of suspension dispensed into the plates for culturing) and multiplied by the dilution factor to obtain the total count per ml. This value was further multiplied by 10 to give the total count of microbes per gram of the faecal sample. Blood samples approximately 10ml per bird were collected from replicates in each treatment into specimen bottles with and without ethylene diamine tetra-acetic acid (EDTA). The haematological parameters such as pack cell volume (PCV) was determined using Winthrobes micro-haematocrit technique, haemoglobin (Hb) was determined with a digital photo colorimeter (Model 312E by Digital Photo Instruments, Germany) at a wavelength of 540nm, after blood had been mixed with Drabkin's solution in a ratio of 1:200 and expressed in g/dl unit. Red blood cell count (RBC) was determined using a coulter electronic counter (Model ZF by Coulter Electronics Ltd, London) and white blood cell count (WBC) were evaluated. Other haematological indices such as mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) and mean cell hemoglobin concentration (MCHC) were computed as outlined by [7].

The serum obtained was analyzed for creatinine and urea by deproteinization and Urease-Berthelot Colorimetric methods using Randox test kit (Randox Laboratories Ltd., UK.). Total protein was determined by the Biurette method using Randox test kit while albumin value was obtained by Bromocresol green method. The globulin was obtained by the method of [8]. Cholesterol was determined by Enzymatic endpoint and glucose by Colorimetric method.

2.6 Chemical Analysis: The proximate composition of the experimental spice was determined according to the procedure [9] and the phytochemical analysis was carried out by the methods of [10, 11] for the presence of saponins, steroids, tannins, flavonoids and alkaloids (Table1).

2.7 Statistical Analysis: All data generated were subjected to one way analysis of variance (ANOVA) for completely randomized design using [12] statistical package and significant differences found among the treatments means were separated using Duncan Multiple Range Test [13].

Table 1. Proximate and phytochemical Compositions of *Xylopia aethiopica* Dried fruits (Grains of Selim)

Parameters %		Phytochemical Test	Results
Dry matter	87.95	Flavonoid	+++
Ash(%)	5.84	Tannin	++++
Crude fibre	10.51	Alkaloid	++
Crude protein	2.73	Steroid	++
Ether extract	9.9	Saponin	+
Nitrogen free extract	58.97	Carbohydrate	+

+ = slightly present, ++= moderately present, +++= present, ++++= strongly present

3.0 RESULTS

The results of the phytochemical analysis showed that *Xylopi aethiopica* dried fruits contain the following phytochemicals: alkaloids, tannins, flavonoids, steroids, saponin and carbohydrates.

The results of the performance characteristics of the finisher broilers given different concentrations of *Xylopi aethiopica* dried fruits in drinking water are presented in Table 3. There were no significant ($P>0.05$) differences in final body weight, total weight gain, daily weight gain, total feed intake, daily feed intake and food conversion ratio in all the treatment groups. However, broiler finishers in treatments given *Xylopi aethiopica* dried fruits were generally better than broiler chicks in treatment 1 given antibiotic in final body weight, total weight gain, daily weight gain and feed conversion ratio.

Results of the microscopy of finisher broiler droppings before and after treatment are presented in Table 4. The results showed that before treatment all treatment groups were highly infected with *Uric acid crystals*, *Staphylococcus spp*, *E.coli*, *Candida albicans*, hookworm, round worm and their ova while treatments 2 and 4 had *Aspergillus flavus*. The results of microscopy after treatment showed a total absence or highly reduced presence of microbes in all treatment groups.

Results of the microbial population count of finisher broilers droppings before and after treatment are presented in Table 5. The colony count of microbes showed that all treatment groups were highly infected with the colony count ranging from 23.33 to 41.00 $\times 10^5$ cfu/ml before treatment. The values dropped significantly in all treatment groups after the administration of the antibiotics and *Xylopi aethiopica* dried fruits. There were no significant ($P>0.05$) differences in colony count before or after treatment in all treatment groups.

The results of haematology of finisher broilers given different concentrations of *Xylopi aethiopica* dried fruits in drinking water are presented in Table 6. Results showed no significant ($P>0.05$) differences in Haemoglobin (HB), Packed cell volume (PCV), Red blood cell count (RBC), White blood cell count (WBC), Mean corpuscular volume (MCV), Mean corpuscular hemoglobin (MCH) and Mean corpuscular hemoglobin concentration (MCHC) in all treatment groups.

Table 2. Composition of Experimental Finisher Broilers Diet

Ingredients	Treatments				
	T1* 0.3g/litre (antibiotic)	T2** 0.4g/litre	T3** 0.6g/litre	T4** 0.8g/litre	T5** 1.0g/litre
Maize	58.00	58.00	58.00	58.00	58.00
Soyabean cake	17.00	17.00	17.00	17.00	17.00
Groundnut cake	9.00	9.00	9.00	9.00	9.00
Fish meal	4.00	4.00	4.00	4.00	4.00
Weath offal	5.00	5.00	5.00	5.00	5.00

156	Palm oil	1.20	1.20	1.20	1.20	1.20
157	Bone meal	3.20	3.20	3.20	3.20	3.20
158	Limestone	1.55	1.55	1.55	1.55	1.55
159	Salt	0.30	0.30	0.30	0.30	0.30
160	Premix	0.50	0.50	0.50	0.50	0.50
161	Methionine	0.15	0.15	0.15	0.15	0.15
162	Lysine	0.10	0.10	0.10	0.10	0.10
163	Total	100.00	100.00	100.00	100.00	100.00
164	Calculated analysis:					
165	Crude protein (%)	20.07	20.07	20.07	20.07	20.07
166	Crude fibre (%)	3.57	3.57	3.57	3.57	3.57
167	M E Kcal/kg	2905.95	2905.95	2905.95	2905.95	2905.95
168	Determined analysis:					
169	Dry matter (%)	87.84	87.84	87.84	87.84	87.84
170	Ash (%)	9.22	9.22	9.22	9.22	9.22
171	Crude fibre (%)	11.33	11.33	11.33	11.33	11.33
172	Crude protein (%)	20.86	20.86	20.86	20.86	20.86
173	Ether extract (%)	3.09	3.09	3.09	3.09	3.09
174	Nitrogen free extract (%)	43.34	43.34	43.34	43.34	43.34

175 Each 2.5kg of finisher vitamin-mineral premix (Bio Organic nutrient systems LTD Nigeria) provided the following vitamins and minerals. Vit
176 A, 8,500,000.00 IU; Vit D3,1,500,000.00 IU; Vit E, 10,000.00 IU; Vit K3,1,500.00mg; Vit B1,1,600.00mg; Vit B2,4000.00mg; Niacin,
177 20,000.00mg; Pantothenic acid; 5000.00mg; Vit. B6, 1,500.00mg; Vit B12, 10.00mg; Folic acid,500.00mg; Biotin 750.00mg, Choline
178 chloride175,000.00mg, Cobalt 200.00mg; Copper 3,000.00mg;Iodine 1,000.00mg; Iron, 20mg,Manganise,40,000.00; Selenium200.00;
179 Zinc30.000.00; Antioxidant, 1,250.00 *Antibiotics (Doxy-gen 20/20 WSP : Doxycycline d hyclate 200mg and Gentamicine de sulfate 200mg)
180 **Grain of selim. ME: Metabolizable energy.

181 The results of serum biochemistry of finisher broilers given different concentrations of *Xylopi*
182 *aethiopioca* dried fruits in drinking water are presented in Table 7. There were no significant
183 ($P>0.05$) differences in total protein, albumin and globulin of their serum. Significant ($P<0.05$)
184 differences were found in creatinine and glucose while highly significant ($P<0.01$) differences
185 occurred in cholesterol and urea content of the treatment groups. The creatinine content of
186 treatment 1 (control) was significantly ($P<0.05$) higher than that of treatment 5 but similar
187 ($P>0.05$) to those of treatments 2, 3 and 4 while treatments 2, 3, 4 and 5 given *Xylopi*
188 *aethiopic* dried fruits were similar ($P>0.05$). The cholesterol values of treatments 1 and 2 were
189 highly significantly ($P<0.01$) higher than those of treatments 3, 4 and 5. Treatment 1 was highly
190 significantly ($P<0.01$) higher than treatments 4 and 5 but similar ($P>0.01$) to treatments 2 and 3
191 in urea, while treatments 2, 3 and 4 were also similar ($P>0.01$). The glucose content of treatment
192 2 was significantly ($P<0.05$) different from all other treatments but others were similar ($P>0.05$).

193

Table 3.: Performance Characteristics of Finisher Broilers Given Different Concentrations of *Xylopi aethiopica* Dried Fruits (grains of selim) in Drinking Water.

Parameters	Treatments					SEM	SIG
	T1*	T2**	T3**	T4**	T5**		
	0.3g/litre	0.4g/litre	0.6g/litre	0.8g/litr	1g/litre		
Initial weight (g)	1431.00	1454.67	1421.00	1377.00	1431.33	16.85	NS
Final body weight (g)	3186.00	3274.33	3181.33	3415.33	3394.00	45.12	NS
Total weight gain (g)	1754.00	1867.67	1789.33	2009.33	1891.33	47.72	NS
Daily weight gain (g)	62.64	66.70	63.90	71.76	67.55	170.46	NS
Total feed intake (g)	4786.00	4888.67	4751.67	4727.67	4543.00	76.20	NS
Daily feed intake (g)	170.93	174.60	169.70	168.84	162.25	27.22	NS
Feed conversion ratio	2.74	2.65	2.67	2.34	2.42	0.071	NS

*Antibiotics (Doxy-gen 20/20 WSP : Doxycycline d'hyclate 200mg and Gentamicine de sulfate 200mg)

**Grains of selim

NS: Not significant

4.0 DISCUSSION

The results of the phytochemical analysis showed that *Xylopi aethiopica* dried fruits has alkaloids, flavonoids, tannins, steroids and carbohydrates present in it while saponin is marginally present. Alkaloids have been documented to possess analgesic, antispasmodic and bactericidal effects [14]. Flavonoids are hydroxylated phenolic substances synthesized by plants in response to microbial infection and they have been known to be antimicrobial substances against a wide range of microorganisms *in vitro*. Their activity is probably due to their capability to complex with extracellular and soluble proteins and to complex with bacterial cell wall [15]. They are also good antioxidant and show strong anticancer activities. They lower the risk of heart diseases [10]. Tannins bind to proline rich protein and interfere with protein synthesis [15]

Table 4.: Microscopic Analysis of Finisher Broiler Droppings Before and After Giving Different Concentrations of *Xylopi aethiopica* Dried Fruits (grains of selim) in Drinking Water.

Treatments	Appearance	Observations Before Treatment	Observations After Treatment
Treatment 1*	Normal formed sample	Hookworm ++, Uric acid crystals +++, growth of <i>Staphylococcus spp</i> , Ova of round worms and <i>E.coli</i>	Crystals ++ and <i>Candida albicans</i> and <i>E.coli</i> but no growth
Treatment 2**	Normal formed sample	Cyst of <i>E.coli</i> +, mature worms, growth of <i>E.coli</i> , <i>Candida albicans</i> , growth of <i>Staphylococcus spp</i> and <i>Aspergillus flavus</i>	Ova of round worm, growth of <i>E.coli</i> and <i>Staphylococcus spp</i>
Treatment 3**	Normal formed sample	Ova of <i>Isosprabelli</i> ++ and growth of <i>Staphylococcus spp</i> ,	Crystals +, and growth of <i>E.coli</i> and <i>Isosporabelli</i>

Treatment 4**	Normal formed sample	Crystals ++, growth of <i>E.coli</i> and <i>Candida albicans</i> and Ova of round worm + Cryst of <i>E.coli</i> +, <i>Candida albicans</i> and growth of <i>Staphylococcus spp</i> , <i>Aspegillus flavus</i> , Mature round worm + and Crystals +++,	<i>E.coli</i> and <i>Staphylococcus spp</i>
Treatment 5**	Normal formed sample	Crystals +, <i>E.coli</i> , growth of <i>Staphylococcus spp</i> and <i>Candida albicans</i>	Crystal ++, growth of <i>Candida albicans</i> and <i>E.coli</i>

Table 5.: Microbial Population Count of Finisher Broiler Droppings Given Different Concentrations of *Xylopiia aethiopia* Dried Fruits (grains of selim) in Drinking Water

	Treatments					SEM	SIG
	T1*	T2**	T3**	T4**	T5**		
Microbial counts	0.3g/litre	0.4g/litre	0.6g/litre	0.8g/litre	1g/litre		
POPULATION BEFORE TREATMENT ($\times 10^5$)	37.00	33.33	41.00	35.33	23.33	2.53	NS
POPULATION AFTER TREATMENT ($\times 10^5$)	0.00	11.33	3.33	12.67	4.67	1.71	NS

*Antibiotics (Doxy-gen 20/20 WSP : Doxycycline d'hyclate 200mg and Gentamicine de sulfate 200mg)

**Grains of selim

NS: Not significant

and this make tannin an anti-nutritional substance. Steroids have been found to have antibacterial properties and are very important compounds in relationship with sex hormones [15]. The saponins found in grains of selim are also known to produce inhibitory effect on inflammation and has the property of precipitating and coagulating red blood cells. Some of the characteristics of saponins include formation of foams in aqueous solutions, hemolytic activity, cholesterol binding properties and bitterness. The results obtained in this study thus suggest the identified phytochemical compounds may be the bioactive constituents in grains of selim, and this could

Table 6.A: Haematology Parameters of Broilers Given Different Concentrations of *Xylopiia aethiopia* Dried Fruits (grains of selim) in Drinking Water.

	Treatment					SEM	SIG
	T1*	T2**	T3**	T4**	T5**		
Parameters	0.3g/litre	0.4g/litre	0.6g/litre	0.8g/litre	1g/litre		
PCV%	27.67	24.67	26.00	27.33	28.67	0.78	NS
HB(g/dl)	9.23	7.57	8.10	8.57	8.87	0.36	NS
RBC($\times 10^6/\mu\text{l}$)	4.33	3.67	4.00	4.33	4.67	0.20	NS
WBC	5.40	6.53	6.33	6.17	5.67	0.30	NS
MCV(FL)	643.33	694.44	700.00	662.22	623.33	42.68	NS

MCH(pg)	214.67	215.00	220.11	208.22	172.83	16.90	NS
MCHC(g/dl)	33.37	29.26	30.99	31.35	30.92	0.65	NS

*Antibiotics (Doxy-gen 20/20 WSP : Doxycycline d'hyclate 200mg and Gentamicine de sulfate 200mg)

**Grains of selim

Table 6.B: Haematology Parameters of Broilers Given Different Concentrations of *Xylopia aethiopica* Dried Fruits (grains of selim) in Drinking Water.

Parameters	Treatment					SEM	SIG
	T1*	T2**	T3**	T4**	T5**		
	0.3g/litre	0.4g/litre	0.6g/litre	0.8g/litre	1g/litre		
PCV%	27.67	24.67	26.00	27.33	28.67	0.78	NS
HB(g/dl)	9.23	7.57	8.10	8.57	8.87	0.36	NS
RBC(X10 ⁶ /μl)	4.33	3.67	4.00	4.33	4.67	0.20	NS
WBC	5.40	6.53	6.33	6.17	5.67	0.30	NS
MCV(FL)	643.33	694.44	700.00	662.22	623.33	42.68	NS
MCH(pg)	214.67	215.00	220.11	208.22	172.83	16.90	NS
MCHC(g/dl)	33.37	29.26	30.99	31.35	30.92	0.65	NS

*Antibiotics (Doxy-gen 20/20 WSP : Doxycycline d'hyclate 200mg and Gentamicine de sulfate 200mg)

**Grains of selim

make this spice to be an increasingly valuable reservoir of bioactive compounds of substantial medicinal merit [15]. This result is similar to the work done by [16, 17, 18, 19] who carried out Phytochemical screening of *Xylopia aethiopica* oil and reported the presence of plant sterols and phenolic compounds such as flavonoids, tannins and the saponins.

The similarities in the final body weights, total weight gain, daily weight gain, total feed intake, daily feed intake and feed conversion ratio in Table 3. is an indication that finisher broilers in all treatment groups responded similarly and positively in weight development to the diet and additives provided. The weight improvement pattern in treatment 1 given antibiotic (Doxy-gen 20/20 WSP : Doxycycline d'hyclate 200mg and Gentamicine de sulfate 200mg) and that of other treatments given *Xylopia aethiopica* dried fruits indicate that *Xylopia aethiopica* dried fruits has growth promoting potential just like the antibiotic and, therefore, *Xylopia aethiopica* dried fruits can be used to replace antibiotic as additive in broiler finisher production. These results are similar to what was reported by [5] who gave different concentrations of *Xylopia aethiopica*

Table 7.: Serology Parameter of Broiler Given Different Concentrations of *Xylopia aethiopica* Dried Fruits (grains of selim) in Drinking Water.

Parameters	Treatments					SEM	SIG
	T1*	T2**	T3**	T4**	T5**		
	0.3g/litre	0.4g/litre	0.6g/litre	0.8g/litre	1g/litre		
Creatinine(mg/dl)	0.77 ^a	0.57 ^{ab}	0.53 ^{ab}	0.50 ^{ab}	0.43 ^b	0.04	*
Cholesterol(mg/dl)	82.00 ^a	72.33 ^a	48.33 ^b	44.67 ^b	35.33 ^b	5.4	**
Glucose(mg/dl)	98.00 ^b	110.00 ^a	101.67 ^b	101.67 ^b	101.67 ^b	1.26	*
Urea(mg/dl)	9.67 ^a	9.33 ^{ab}	8.00 ^{abc}	7.67 ^{bc}	7.33 ^c	0.32	**

Albumin(g/dl)	2.10	2.37	2.23	2.13	2.1	0.07	NS
Total Protein(g/dl)	5.13	5.1	5.13	5.17	5.00	0.17	NS
Globulin(g/dl)	3.03	2.73	2.90	3.03	2.90	0.15	NS

*Antibiotics (Doxy-gen 20/20 WSP : Doxycycline d'hyclate 200mg and Gentamicine de sulfate 200mg)

**Grains of selim

a, b, c means with different superscripts in the same row are significantly (P<0.05)*, (P<0.01)** different. NS: Not significant.

dried fruits in drinking water to broiler chicks and obtained improved performance just like that of antibiotic growth promoter. [20, 21, 22] also reported that garlic (powder or aqueous extract) as an additive in broiler production produced improved body weight gain, daily feed intake and feed conversion ratio. The result is also in agreement with the results of [23, 24, 25, 26, 27] who reported improved body weight gain in broilers when ginger (*Zingiber officinale*, a spice) was used at different rates in feed or drinking water as an additive. [28, 21, 29, 30, 31, 32] also used thyme as an additive in broiler production and had improved body weight performance and concluded that thyme oil or powder may be a promising alternative to antibiotic growth promoters. The similarities in feed intake and feed conversion ratio which conform to the growth pattern in this study also agree with the reports of these authors. The improved performance of *Xylopiya aethiopica* dried fruits may be attributed to the presence of several phytochemical compounds in the fruits such as alkaloids, flavonoids, sterols, saponins and tannins which have biological activities as antioxidant, antimicrobial and pharmacological effects [5]. However the final weight in the finisher phase of this experiment of broiler birds given different concentration of *Xylopiya aethiopica* dried fruits in their drinking water which are treatments 2 to 5 compared to the control treatment indicate that broiler birds can be given all concentrations of *Xylopiya aethiopica* dried fruits in drinking water from 0.4g/litre to 1.0g/litre with positive effect on body weight performance.

The elimination of mature worms in the treatments 2, 3, 4 and 5 given grains of selim shows that the spice possesses some anthelmintic activities. The decrease of microbial population count of treatments given grains of selim also indicate that grains of selim has antimicrobial properties. The dominant microorganisms identified in the population were the bacteria *Candida albicans*, *E.coli* and *staphylococcus spp*. The decrease in the population of *Candida albicans*, *E.coli* and *staphylococcus spp* in all treatments given grains of selim in drinking water showed that grains of selim has components that possess antibacterial activities and establish the fact that grains of selim is an antimicrobial spice for finisher broilers and could effectively replace antibiotics in their production. This result is similar to the findings of [33] who reported a continuous reduction in the population of *E.coli* with increased concentrations of grains of selim. The result is also similar to the findings of [34] who obtained significant reduction in the bacterial counts of *Staphylococcus spp*, *E. coli* and *Salmonella spp* when essential oil isolated from ginger was used as a decontaminating agent in chicken meat.

The haematological parameters investigated were not significantly affected by giving different concentrations of grains of selim in drinking water of broilers compared to those given antibiotic. Red blood cells (RBC) are responsible for the transportation of oxygen and carbon dioxide in the blood as well as manufacture of haemoglobin hence higher value indicate a greater potential for this function and a better state of health. White blood cells are involve in protecting the body from infection and consist of lymphocytes, monocytes, Neutrophils, Eosinophils and Basophils. They help to eliminate virus-infected cells, enhance the production of antibodies and engulf

foreign materials (antigens) that enter the body. The general none significance of the white blood cell count (WBC) shows that whatever antibiotic can do to impair or enhance the broiler ability to wade off infection, grain of selim can equally do the same at the concentrations of 0.4g/litre to 1.0g/litre of water. These results are in agreement with that reported by [34] that garlic mobilized immune system and empowers the defense ability of the body against infectious organisms. These results are also in agreement with the studies of [35], who reported that these plants improve the health status and performance of broilers, in addition to improve antibody titre against Newcastle Disease virus.

The none significant differences in total protein, albumin and globulin among treatments given grains of selim compared to control treatment indicate that grains of selim have the same effect as antibiotic on albumin, total protein and globulin content in the birds. The total protein values in this study agree with what was reported by [37] who found that ginger supplementation didn't affect the total blood protein and the none significant differences in albumin and globulin shows that grains of selim is as good as antibiotics and can effectively replace antibiotics. Result showed that grains of selim is anti-cholesterol and the higher the concentrations of grains of selim given to broilers the lesser their cholesterol levels. This result is similar to what was reported by [21, 38] who reported that feeding garlic decreased blood cholesterol significantly in broiler chicks. [21, 38, 30, 40] reported significant decrease in blood cholesterol in response to thyme oil administration in broiler birds. These findings are similar to those of [41], who found a significant decrease in blood cholesterol when feeding chicks up to 6% ginger powder. However the results on serum glucose is in agreement with what was reported by [35] who stated in their study that thyme supplementation in broiler diets significantly increased glucose level compared to those of the control group. The possible reason for increase serum glucose may be due to the abdominal lipids catabolism of gluconeogenesis process in the birds with the administration of grains of selim, just as crushed thyme consumption by broiler chickens was reported to have increased the serum glucose [42]. Results showed significance decrease in blood serum creatinine and urea in treatments 4 and 5 compared to control. These results showed that the higher the concentrations of grains of selim given to broilers the lesser their blood serum creatinine and urea. This implies that the kidney of broilers given grains of selim functioned better than that of the control, since elevated creatinine level signifies impaired kidney function or kidney disease. As the kidneys become impaired for any reason, the creatinine level in the blood will rise due to poor clearance of creatinine by the kidney. Blood urea level is another indicator of kidney function, urea is also a metabolic by-product which can build up if kidney functioning is impaired. It can, therefore, be concluded that grains of selim administration in broilers has no adverse health implications on the birds.

5.0 CONCLUSION

The similarity of the weight performance, the reduction effect on microbial population, the similarity in haematological parameters, the better serum chemistry values of broilers given grains of selim in this study indicate that *Xylopi aethiopica* dried fruits (grains of selim) has growth promoting potentials and can serve as a natural alternative to antibiotics in broiler production with no adverse health implications on birds or consumers of the products. The elimination of mature worms also indicate that grains of selim possess some anthelmintic activities.

REFERENCES

1. Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A. (2009). Agroforestry Database: a tree reference and selection guide version 4.0 World Agroforestry Centre, Kenya.
2. Yegani M, Smith TK, Leeson S, Boermans HJ. Effects of feeding grains naturally contaminated with *Fusarium* mycotoxins on performance and metabolism of broiler breeders. *Poult. Sci.* 2006; 85: 1541-1549.
3. Hossain, M.M. Herbs versus drugs: Aspect World Poultry. 2009. Accessed February 14, 2015. Engormix.com/165/po.htm
4. Karimi A, Yan F, Coto C, Park JH, Min Y, Lu C, Gidden JA, Lay JO, Waldroup PW. Effect of level and source of oregano leaf in starter diet for broiler chickens. *J. Appl. Poult. Res.* 2010; 19 (2): 137-145.
5. Isikwenu JO. (2014) *Xylopia aethiopica* dried fruits (Grains of selim) as additive in starter broiler production. *Proc. 14th Euro. Poult. Conf.* 2014; 578. Stavanger, Norway.
6. FAO. Food and Agricultural Organization. Manuals of food quality control 4. FAO Food and Nutrient paper, United Nations Rome, Italy, *Microbiol. Analysis.* 1979; 14 (4): A1-F10.
7. Seal US, Erickson AW. Haematology, blood chemistry and protein polymorphisms in the white tail deer. *Comp. Biochem. Physio.* 1979; 39 (4): 695-713.
8. Coles EH. *Veterinary clinical pathology.* 4th Edition Published by WB Saunders Company, Philadelphia, USA. 1986.
9. AOAC. Official methods of analysis, 15th edition. Association of Official Analytical Chemists, Washington D. C.; 1990.
10. Sofowora A. *Medicinal plants and medicine in africa.* John Whilley Spectrum Books, Ibadan, Nigeria. 1993; 120-123.
11. Trease GE, Evans WC. *Textbook of pharmacognosy.* 1989. 12th ed. Ballere, Tinnall London.
12. SPSS (Statistical Package for the Social Sciences). 17.0. SPSS Inc. Chicago, IL, 60606.
13. Duncan DB. Multiple range and multiple F-tests. *Biometrics.* 1955; 11:1 – 42.
14. Okigbo RN, Anuagasi CL, Amadi JE, Ukpabi UJ. Potential inhibitory effects of some african tuberous plant extracts on *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans*. *Int. J. Integr. Biol.* 2009; 6 (2): 91-98.
15. Yadav RNS, Munin A. Phytochemical analysis of some medicinal plants. *J. Phyto.* 2011; 3 (12): 10-14.
16. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. *Afri. J. Biotech.* 2005; 4 (7): 685-688.
17. Kumar RS, Sivakuma T, Sunderem RS, Gupta M, Muruges K, Rajeshwa Y, Kumar MS, Kumar KA. Antioxidant and antimicrobial activities of *Bauhinia recemosa* L. stem bark. *Braz.J. Med. Biol. Res.* 2005;38: 1015-
18. Law M. Plant sterol and stanol margarines and health. *Br. Med J.* 2000; 320: 72-
19. Ostlund RE Jr. Phytosterols in human nutrition, *Ann. Rev. Nutr.* 2002; 22: 533-549.
20. Aji SB, Ignatius K, Ado AY, Nuhu JB, Abdulkarim A. Effects of feeding onion

- (*Allium cepa*) and garlic (*Allium sativum*) on some performance characteristics of broiler chickens. Res. J. Poult. Sci. 2011; 4:22-27.
21. Mansoub NH. Comparison of effects of using thyme and probiotic on performance and serum composition of broiler chickens. Advc. Environtl. Biol. 2011; 5: 2012 – 2015.
22. Pourali M, Mirghelenj SA, Kermanshahi D. (2010). Effect of garlic powder on productive performance and immune response of broiler chickens challenged with newcastle disease virus. Globa Veterinaria. 2010; 4: 616 – 621.
23. Tekeli A, Kutlu HR, Celik L. Effect of *Z. officinale* and propolis extracts on the performance, carcass and some blood parameters of broiler chicks. Current Res. Poult. Sci. 2011; 1:12 – 23.
24. Herawati . The effect of feeding red ginger as phytobiotic on body weight gain, feed conversion and internal organs condition of broiler. Int. J. Poult. Sci. 2010; 9(10): 963 – 967.
25. Onu PN. Evaluation of two herb spices as feed additives for finisher broilers. Biotech. Ani. Husbdry. 2010; 26: 383 – 392.
26. Javed M, Durrani F, Hafeez A, Khan RU, Ahmad I. Effect of aqueous extract of plant mixture on carcass quality of broiler chicks. ARPN J. Agri. Biol. Sci. 2009; 4:37 – 40.
27. Farinu GO, Ademola SG, Ajayi AO, Babatunde GM. Growth, haematological and biochemical studies on garlic and ginger-fed broiler chickens. Moor J. Agri. Res. 2004; 5:122-128.
28. Al-Mashhadani EA, Farah KA, Farhan YM, Al-Mashhadani HE. Effects of anise, thyme essential oils and their mixture on broiler performance and some on physiological traits. Egypt. Poult. Sci. 2011; 31: 481-489.
29. Toghyani M, Tohidi M, Gheisari AI, Tabeidian SA. Performance, immunity, serum biochemical and haematological parameters in broiler chick fed dietary thyme as alternative for an antibiotic growth promoter. Afri. J. Biotech. 2010; 9:6819 – 6825.
30. Al-Kassie GAM. Influence of two plants extracts derived from thyme and cinnamon on broiler performance. Pakistan Vet. J. 2009; 29: 169-173.
31. Bolukbasi SC, Erhan MK, Ozkan A. Effects of dietary thyme oil and vitamin E on growth, lipid oxidation, meat fatty acid composition and serum lipoproteins of broilers. South Afri. J. Ani. Sci. 2006; 36: 189 – 196.
32. Zhang K, Yan YF, Keen CA, Waldroup PW. Evaluation of microencapsulated essential oils and organic acids in diets for broiler chickens. Int. J. Poult. Sci. 2005; 4:612 – 619.
33. Sudrashan S, Fairoze N, Wildfred S, Shekar R. Effect of aqueous extract and essential oils of ginger and garlic as immunostimulant in chicken meat. Res. J. Poult. Sci. 2010; 3:58 – 61.
34. Josling P. Preventing the common cold with a garlic supplement: a double-blind placebo-controlled survey. Adv. Ther. 2001; 18: 189-193. PMID: 11697022.
35. Mansoub NH, Myandoab MP. The effect of different levels of

- 426 thyme on performance, carcass traits and blood parameters of broilers.
427 Annals Biol. Res. 2011; 2 (4): 379 – 385.
- 428 36. Jameel YJ. The effect of adding Thyme vulgaris and Cinnamomum
429 zeylanicum on production performance and some blood traits in broiler
430 chicken. M.Sc. Thesis in science of public health. College of Veterinary
431 Medicine. University of Baghdad. Iraq. 2008.
- 432 37. Mohamed AB, Al-Rubaei MAM, Jalil AQ. Effect of ginger
433 (Zingiber officinale) on performance and blood serum parameters of broiler.
434 Int. J. Poult. Sci. 2012;11: 143-146.
- 435 38. Rahimi S, Teymouri Zadeh Z, Karimi Torshizi MA, Omidbaigi R, Rokni
436 H. (2011). Effect of three herbal extracts on growth performance, immune
437 system, blood factors and intestinal selected bacterial population in broiler
438 chickens. J. Agri. Sci. Tech. 2011; 13: 527 – 539.
- 439 39. Horton GMJ, Fennell MJ, Prasad BM. Effect of dietary garlic (Allium sativum) on
440 performance, carcass composition and blood chemistry changes in broiler chickens. Can.
441 J. Anim. Sci. 1991; 71: 939-942.
- 442 40. El-Ghousein SS, Al-Beitawi NA. The effect of feeding of crushed
443 thyme (thymus vulgaris) on growth, blood constituents, gastrointestinal tract
444 and carcass characteristics of broiler chickens. J. Poult. Sci.
445 2009; 46: 100-104
- 446 41. Ademola SG, Farinu GO, Babatunde GM. Serum lipid, growth and haematological
447 parameters of broilers fed garlic, ginger and their mixtures.
448 World J. Agri. Sci. 2009; 5 (1): 99 – 104.
- 449 42. Al-Homidan AA. Efficacy of using different sources and levels of allium cepa, allium
450 sativum and zingiber officinale on broiler chick's performance. Saudi J. Biol. Sci. 2005;
451 12 (2): 96 - 102.