

Short Research Article

A Rapid Morphological Screening Procedure for Pea (*Pisum spp.*) Under Drought Stress in Greenhouse Settings

Aims: This work proposes a rapid morphological screening procedure for peas grown under drought stress aiming to provide phenotypic evaluations for further research.

Material and methods: The plant yield and morphological changes of two widely grown pea cultivars (*Aragorn* and *Banner*) to drought stress was evaluated in this work. A total of eighty seeds of each pea variety were grown in a randomized completed block with four replicated in a greenhouse located in Moscow, ID. Plants were subjected to optimal irrigation (1400 ml as a control) and two treatments that reduced optimal irrigation rates by 40% (840 ml, moderate drought) and 60% (560 ml, severe drought) to induce drought stress. The test was repeated.

Results: Varieties significantly ($P<0.05$) differed in their response to water deficiency. The variety 'Banner' appeared to be the most drought tolerant than *Aragorn* with high values at control, moderate and severe as (85.00±xx) cm, (87.21±xx) cm and (66.02±xx) cm respectively for total plant height, while the values for total dry weight were (3.65±xx) g, (2.18±xx) g and (1.26±xx) g respectively. Similarly, there were significant ($P<0.001$) differences among treatments on growth parameters, with highest values recorded at optimal irrigation (1400 ml) as (21.52±xx) cm, (62.18±xx) cm, (0.24±xx) g, (2.86±xx) g and (2.30±xx) seeds/pod for root length, shoot height, root weight, shoot weight and number of seeds respectively. The growth parameters decreased with a decrease of irrigation. but emergence and flowering dates were not affected by the treatments. Treatments were also significantly different for all measurements (root pod numbers, width, length and number of viable seeds, shoot height, root length, and shoot dry weight). Root weight was not significantly different between treatments. **Conclusions:** As drought and stress conditions are expected to increase as global climate change progresses, breeding for drought is a promising area. Therefore, *Banner* variety has been It was possible to observe the effect of drought in each treatment with significant differences between varieties and treatment ($p<0.001$). The variety 'Banner' appeared to be the most drought tolerant, which was reflected by both growth and yield parameters. 'Aragorn' had shown many drought stress symptoms even in the moderate drought treatment, such as very low yield. Our work also identified as a potential parental material to be used in breeding for drought tolerance.

Comment [E1]: Include the standard errors for all the parameter means

Keywords: drought, greenhouse plants, agronomic methods, stress tolerance, peas, growth parameters, irrigation, *Banner*, *Aragorn*, breeding.

1. INTRODUCTION

Over the past several decades, drought stress has become one of the biggest concerns in agriculture worldwide. Adaptation to abiotic stresses is an important requirement of agriculturally relevant crops and development of enhanced drought tolerance plays an essential role in plant breeding [1].

The development of drought tolerant plants is of increasing priority and relevance as the area affected by significant drought is expected to markedly increase in the future due to global climate change [2]. While progressive water withdrawal has been studied in the greenhouse for beans, it generally does not mimic actual field conditions [3].

While limited studies and techniques for simulating field drought stress in a greenhouse environment have been conducted [4], more studies are needed to adequately formulate a reliable protocol for legumes, especially peas. It is the aim of this research to provide a rapid morphological screening procedure for pea under drought stress in greenhouse settings.

The use of peas as a model crop for genetic studies has a very rich history, beginning with Mendel in the 1860s. *Pisum* spp. is adapted to cool, semi-arid to sub-humid growing conditions [5]. However, pea has been grown in diverse environments across the globe and the demand for production continues to increase [6]. Drought is of great concern in pea and it has been reported to result in a loss of between 38.5 and 43% production under different stress levels [7].

2. MATERIAL AND METHODS

Two dry pea cultivars, 'Aragorn' and 'Banner,' commonly grown in the U.S. and New Zealand and developed by ProGene Plant Research LLC in conjunction with NZ Plant Research LTD were utilized for this experiment. 80 seeds of each genotype were prepared by sterilization utilizing a 10% bleach aqueous solution, rinsed in sterile, distilled H₂O and air dried for a period of 24 hours.

Five seeds of each variety were planted in a 3.75 liter high density polyethylene/polypropylene polycan container (Anderson Pots, Oregon, USA) filled with 690 g of Sunshine Mix #1 growing media (Sun Gro Horticulture, Agawam, MA) and arranged in a complete randomized design with four replicates per cultivar. The experiment was repeated.

The peas were grown under controlled greenhouse conditions with temperatures ranging from 12 to 18°C at night and 21 to 27°C during daylight hours (MicroGrow Greenhouse Systems Growmate (Temecula, CA) utilizing 400 Philips HID Ceramalux High Pressure Sodium bulbs). The plants were subjected to a 12-hour photoperiod per 24 hours.

Three different irrigation regimes were tested: control, moderate drought and severe drought. Field capacity of the growing media served as the experimental control. Field capacity was determined by wetting 690 g of media until saturation was reached. The media was then allowed to drain for a period of 24 hours and oven dried to determine maximum moisture holding capacity (field capacity).

Previous field trials conducted by ProGene Plant Research [8] have shown that approximate reductions in irrigation of 40% and 60% caused visual drought stress symptoms on plots of field peas. Control plants were irrigated to the full field capacity of the growing media (1400 ml) and irrigated once every 7 days until flowering and then once every 3.5 days from flowering until maturity. Drought stress test pots received amounts of water decreased by 40% (840 mL) and 60% (560 mL) of field capacity, respectively, at the same watering interval.

Phenotyping methods consisted of the collection of the following agronomic notes: emergence and flowering dates, plant survival, shoot height and root length, shoot and root dry weight, number of pods per plant, pod length and width, and number of viable seeds produced per plant. Statistical analysis of the data, including the Fisher's LSD method, was

performed using Minitab 17 Statistical Software (Minitab, State College, PA) and Agrobase Generation II® (Agronomix Software Inc., Winnipeg, Manitoba).

3. RESULTS AND DISCUSSION

As shown in previous research, emergence and flowering dates were not affected by drought stress treatments [9]. All 'Banner' treatments (including control) flowered 52 days after planting, while all 'Aragorn' treatments (including control) flowered 54 days after planting. However, varieties differed in their response to water deficiency and plant survival was jeopardized. The 40% reduction from field capacity trial yielded a 20% rate of mortality for both varieties tested. The 60% reduction from field capacity trial resulted in 20% mortality in 'Aragorn' and 27% in 'Banner.' For all morphological measurements: shoot height, shoot dry weight and number of seeds per pod showed a consistent decrease due to drought for both varieties (Table 1).

Irrigation treatment	Variety name	Root length (cm)	Shoot height (cm)	Root weight (g)	Shoot weight (g)	Pod length (cm)	Pod width (cm)	Number of Seeds/pod
Control	Aragorn	22.54 ±2.1	61.21 ±2.6	0.32 ±0.04	2.44 ±0.2	2.99 ±0.67	0.29 ±0.07	0.60 ±0.4
	Banner	17.57 ±1.2	67.43 ±3.9	0.18 ±0.02	3.47 ±0.33	4.79 ±0.24	0.59 ±0.01	2.75 ±0.45
Moderate	Aragorn	22.79 ±2.58	55.27 ±1.6	0.27 ±0.03	1.55 ±0.14	1.55 ±0.03	0.067 ±0.01	0.00
	Banner	24.02 ±2.56	63.19 ±2.03	0.23 ±0.03	1.95 ±0.13	4.22 ±0.3.2	0.55 ±0.04	2.23 ±0.37
Severe	Aragorn	15.25 ±1.5	42.93 ±2.03	0.19 ±0.031	1.09 ±0.13	1.91 ±0.46	0.12 ±0.05	0.00
	Banner	19.48 ±1.48	46.54 ±2.52	0.19 ±0.02	1.07 ±0.1	3.92 ±0.39	0.59 ±0.02	2.00 ±0.33

Table 1 Morphological data means; showing: irrigation treatment, variety name, root length, shoot height, root weight, shoot weight, pod length, pod width, and number of seeds per pod.

Comment [E2]: The title should be written at the top of the table

A statistically significant ($P<0.001$) difference in shoot height was observed in both variety and treatment ($P<0.001$) comparisons. 'Banner' was taller than 'Aragorn' in all treatments and both varieties suffered from increasing stunt with relation to decreasing irrigation (Figure 1).





Figure 1 Plant total height (cm) of 'Banner' and 'Aragorn' across water treatments - control (1400 ml), moderate drought (840 ml) and severe drought (560 ml) stress. Total height included shoot and root measurements.

Comment [E3]: Bar chart is appropriate for this type of data. See below how it is looking. Line graphs are used to compare changes over same period of time, while Bar Charts are used to compare things between different groups.



'Banner' and 'Aragorn' exhibited a 6.3% and 9.8% reduction in plant shoot height, respectively, in the moderate drought treatment. Both had the least shoot height, at only

70% of their potential, when compared to the control plants, in the severe drought treatment. Shoot dry weight was also impacted by drought, being significantly different ($P < 0.001$) among treatment and between varieties. For the moderate and severe drought stress treatments 'Banner' shoot dry weight (Figure 2) was reduced by 44% and 70%, respectively, while 'Aragorn' was reduced by 55% and 69%, respectively.

Growth reduction was one of the major drought stress treatments results. It was also observed that leaf area was visually reduced. In a similar study involving wheat cultivars, leaf area was also reduced but root development was retarded in only one cultivar [10]. A reduction in plant height, lower leaf and stem dry weights were also shown, in addition to the production of a lower overall aerial biomass among drought-stressed plants grown in a greenhouse environment [4].





Figure 2 Total plant dry weight (g) of 'Banner' and 'Aragorn' across water treatments: control (1400 ml), moderate drought (840 ml) and severe drought (560 ml) stress. Total dry weight included shoot and root measurements.

Comment [E4]: Bar chart is appropriate for this type of data. See below how it is looking. Line graphs are used to compare changes over same period of time, while Bar Charts are used to compare things between different groups.



113

114 Root length did not significantly ($P>0.05$) differ among treatments and between varieties.
 115 'Aragorn' roots were not consistently affected on the moderate drought treatment but had an
 116 important reduction of 32% on root length on the severe drought; while 'Banner' roots were
 117 not consistently affected between the treatments. Root dry weight did not differ ($P>0.05$)
 118 significantly among treatments and between varieties. Overall, 'Aragorn' produced more
 119 roots than 'Banner' in all trials.

120 The number of pods per plant were highly significantly ($P<0.001$) different among treatments
 121 and also showed a significant ($P<0.05$) difference between varieties. In all treatments,
 122 'Banner' had more pods than 'Aragorn' but both varieties exhibited a reduction in the number
 123 of pods with increasing drought. 'Banner' displayed a 38.5% and 41.5% reduction in the
 124 overall number of pods as drought increased, respectively; while 'Aragorn' had a reduction of
 125 42.7% and 56%, respectively.

126 In addition, pods also became shorter and thinner with decreasing irrigation when compared
 127 to the control (Table 1). 'Aragorn' did not produce any viable seeds in either drought
 128 treatment while 'Banner' was able to produce viable seed throughout all treatments. 'Banner'
 129 suffered a reduction in the number of viable seeds by 19% in the moderate drought
 130 treatment and 27% in the severe drought treatment.

131 Yield measured by yield components (number of pods, pod length, width, and number of
 132 viable seeds per pod) was also reduced under simulated drought conditions, with simulated
 133 severe drought underperforming all other treatments. In a similar study, analysis of variance
 134 has indicated that there were significant differences between genotypes regarding
 135 agronomic and morphological traits of wheat under drought [11].

When both varieties are combined for the analysis in each treatment it is possible to observe the overall effect of drought on the investigated plant material. Significant differences can be then observed across treatments for all the evaluated parameters (Table 2).

Treatment/ Parameters	Control		Moderate		Severe	
Root length	21.52	A	19.57	A	15.57	B
Shoot height	62.18	A	54.37	B	40.22	C
Root weight	0.24	A	0.23	A B	0.17	B
Shoot weight	2.86	A	1.58	B	0.96	C
Number of pods	2.30	A	1.29	B	0.92	C

Table 2 Morphological data means across treatments (control, moderate and severe drought) combining both varieties; showing grouping information using Fisher LSD method and 95% confidence. Means that do not share a letter within a row are significantly ($P < 0.01$) different.

Our data suggests that plant characteristics can influence drought tolerance. Comparing the two varieties used in our study, 'Banner,' the variety that performed better under drought stress, has good early vigor emerging five to six days earlier than 'Aragorn' [8].

There are many metabolic changes that can enable a plant to withstand drought, including specific proteins such as chaperonins and dehydrins that are induced in certain plants under drought [12]. Water stress also induced the accumulation of soluble sugars in epicotyls between 2.8- and 5.1-fold in another study and the osmotic adjustment varied greatly depending on cultivar [13]. Moreover, temperature can contribute to drought stress and can induce significant changes in plant cell gene expression [14].

4. CONCLUSION

The present study has proposed a simple, rapid morphological screening procedure for pea under drought stress in greenhouse settings and provided evaluation of two commercial pea cultivars. 'Banner' appeared to be the most drought tolerant, which was reflected by both growth and yield parameters. 'Aragorn' had shown many drought stress symptoms even in the moderate drought treatment, such as very low yield. As drought and stress conditions are only expected to increase as global climate change progresses, breeding for drought is a promising area. Therefore, Banner variety has been identified as a potential parental material to be used in breeding for drought tolerance.

Our study has proposed a simple, rapid morphological screening procedure for pea under drought stress in greenhouse settings and provided evaluation of two commercial pea cultivars. We have shown that drought stress impacts growth regardless of the variety type. Future studies including more varieties and measurements of leaf water content and relative water content will be pursued. Of the two varieties tested, the early variety, 'Banner,' was clearly the most drought tolerant variety, producing viable seed in every moisture regime tested, a larger overall amount of pods per plant, viable seed, and overall plant biomass. Aragorn, which on average emerged 5-6 days later than Banner underperformed Banner in

Comment [E5]: 1. The title should be written at the top of the table
2. Include standard errors for the mean values on the table

Comment [E6]: This last part should be a note on the bottom of the table

Comment [E7]: The original conclusion was excellent you only needed to interchange and modify the last two sentences

~~nearly every respect, except for root production in which it exceeded Banner in all trials. 'Aragorn' displayed several drought stress symptoms even in the moderate drought treatment, and produced no viable seed in any treatment outside of the control. In the most extreme drought treatment, in which the plants received moisture equivalent to only 40% of field capacity of the growing medium, Banner suffered 27% mortality, exhibited shoot dry weight that was reduced by 70% overall, produced 41.5% fewer pods, which contained 27% fewer viable seeds. Comparatively, Aragorn suffered 20% mortality, exhibited shoot dry weight that was reduced by 69% overall, produced 56% fewer pods, which contained not a single viable seed under the same growing conditions. In the moderate drought treatment, in which the plants received moisture equivalent to only 60% of the field capacity of the growing medium, Banner suffered 20% mortality, a 6.3% reduction in average plant height, exhibited shoot dry weight that was reduced by 44% overall, produced 38.5% fewer pods, which contained 19% fewer viable seeds. Comparatively, Aragorn suffered 20% mortality, a 9.8% reduction in average plant height, exhibited shoot dry weight that was reduced by 55% overall, produced 42.7% fewer pods, which contained not a single viable seed under the same growing conditions.~~

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Witt S, Galicia L, Lisec J, Cairns J, Tiessen A, Araus JL, Fernie AR. Metabolic and phenotypic responses of greenhouse-grown maize hybrids to experimentally controlled drought stress. *Molecular plant*. 2011; ssr102.
2. de Paiva Rolla AA, Carvalho JDFC, Fuganti-Pagliarini R, Engels C, do Rio A, Marin SRR, Oliveira MCN, Beneventi MA, Corrêa F, Guimarães M, Farias JBC, Neumaier N, Nakashima K, Yamaguchi-Shinozaki K, Nepomuceno AL. Phenotyping soybean plants transformed with rd29A: AtDREB1A for drought tolerance in the greenhouse and field. *Transgenic research*. 2014; 23(1), 75-87.
3. Kavar T, Maras M, Kidric M, Sustar-Vozlic J, Meglic V. Identification of genes involved in the response of leaves of *Phaseolus vulgaris* to drought stress. *Mol Breeding*. 2008; 1(21), 159-172.
4. Pennypacker BW, Leath KT, Stout WL, Hill RR. Technique for simulating field drought stress in the greenhouse. *Agronomy journal*. 1990; 82(5), 951-957.
5. Coyne C J, McGee RJ, Redden RJ, Ambrose MJ, Furman BJ, Miles CA. Genetic Adjustment to Changing Climates: Pea (pp. 238-250). Wiley Blackwell: Chichester, UK. 2011.
6. EMBRAPA – Empresa Brasileira de Pesquisa Agropecuária. Centro Nacional de Pesquisa de Hortaliças. Ervilha para produção de grãos verdes. 2 Nov. 2008. <www.cnph.embrapa.br/cultivares/ervilha>.
7. Carvalho JA, Rezende, FC, Aquino RF, Freitas WA, Oliveira EC. Produção da ervilha cultivada em ambiente protegido sob diferentes tensões de água no solo. *Revista Brasileira de Engenharia Agrícola e Ambiental, Campina Grande*. 2012; 1(16), 44-50.

- 220 8. Braunwart K, Wood M. Drought Stress on Pea Crops. ProGene Plant Research; 2014.
- 221 9. Muchow RC. Phenology, Seed yield and WaterUse of Grain Legumes Grown under
222 Different Water Regimes in a Semi-arid Tropical Environment. FieldCrops Research. 1985;
223 11, 81-87
- 224 10. Klamkowski K, Treder ww. Response to Drought Stress of Three Strawberry Cultivars
225 Grown Under Greenhouse Conditions. J. Fruit Ornam. Plant Res. 2008; 180 (16):179-188.
- 226 11. Ahmadizadeh M, Nori A, Shahbazi H, Aharizad S. Correlated response of morpho-
227 physiological traits of grain yield in durum wheat under normal irrigation and drought stress
228 conditions in greenhouse. African Journal of Biotechnology. 2014; 10(85), 19771-19779.
- 229 12. Vaseca I, Akiscan Y, Demirevska K, Anders I, Feller U. Drought stress tolerance of red
230 and white clover: comparative analysis of some chaperonins and dehydrins. Scientia
231 Horticulturae, Canada. 2011; 4(130), 653-659.
- 232 13. Sánchez FJ, Andrés EF, Tenorio J, Ayerbe L. Growth of epicotyls, turgor maintenance
233 and osmotic adjustment in pea plants (*Pisum sativum* L.) subjected to water stress. Field
234 Crops Research. 2014; 2(86), 81-90.
- 235 14. Talalaiev O, Korduym E. Expression of small heat shock protein (sHSP) genes in the
236 garden pea (*Pisum sativum*) under slow horizontal clinorotation. Plant signaling & behavior.
237 2014; 9(4), e29035.