

1 Original Research Article
2 **Quality Assessment of Tubewell Water for**
3 **Irrigation and Impact on Soil and Crops in**
4 **Central Punjab, Pakistan**

5 .

6 **ABSTRACT**

7

Aims: The study was conducted to assess the quality of tubewell (TW) waters for irrigation and their effects on soils and crops.

Place and Duration of Study: Canal Command Area (CCA) of Chichawatni and Sahiwal tehsils of district Sahiwal, Punjab, Pakistan during 2012 and 2013.

Methodology: A total of 529 TW water samples were collected from nine distributaries of Lower Bari Doab Canal (LBDC); four on right side (RD); five on left side (LD). Twenty composite soil samples were collected at plough layer depth (0 to 15 cm) from the selected farmer fields irrigated with TW and canal water (CW). Soils were analysed for pHs, E_{Ce}, and water samples were analysed for EC, cations (Ca^{2+} + Mg^{2+} , Na^{+}) and anions (CO_3^{2-} , HCO_3^{-}) and then RSC and SAR were computed. Total dissolved salts (TDS) of TW waters were used to estimate the amount of salts added into soil with the irrigation. The salts addition into soil under different crops was also calculated using the consumptive use by crops. The yield data of crops were recorded from 20 farmer's fields irrigated with CW and TW waters to evaluate the effects of water quality on the yields.

Results: Data based on EC, RSC and SAR values showed that 71% TW waters were saline and >20% samples were sodic in nature. The quality of TW water in RD was

comparatively better than in LD due to its location between LBDC and Ravi River. The farmers used poor quality TW waters (TDS from 218 to 3309 mg L⁻¹) that added 0.66 to 10.05 t ha⁻¹ salts into soil with 0.30 m irrigation water. Similarly, 15.75, 27.80, 33.36, 44.48 and 55.99 t ha⁻¹ TDS were deposited in soil by wheat, cotton, potato, spring maize and rice, respectively, irrigated with TW water of TDS 3309 mg L⁻¹. The yields of crops reduced from 3 to 15% where usually TW water was used compared to CW irrigated fields.

Conclusion: Results showed that quality of 71% TW waters was not suitable for irrigation purpose due to higher concentration of soluble salts. The concentrations of chemical constituents in water from the LBDC aquifer system vary both depth and location wise. The application of TW irrigation having different salinities added substantial amount of soluble salts into soil that affected the soil quality. Resultantly, the yields of different crops reduced from 3 to 15%. Presently the farmers are using brackish TW water with CW but the continuous use of poor quality irrigation water would have serious consequences for sustainability of soil and crop productivity in the area.

Keywords: Tubewell water quality; Soil quality; crop yield; EC; SAR; RSC.

1. INTRODUCTION

Water quality is an important factor when considering an irrigation programme. The common water quality parameters are electrical conductivity (EC), sodium adsorption ratio (SAR) and residual sodium carbonate (RSC) [1]. The canal water supply is not sufficient to meet the water requirement of the crops raised under intensive cropping system of Punjab, Pakistan. Hence, farmers are forced to meet this deficit through tube wells water which, over 70% is of poor quality [2, 3].

Soluble salts in irrigation water, if present in sufficient quantity, affect soil quality and crops yields. Soil degradation has been defined as a reduction in soil quality in response to natural or anthropogenic causes, in a manner, that reduces current or potential productivity [4]. Soil degradation reduces the productivity of soil and water resources in many areas of the world, causing reductions in crop yields, higher production costs, and lower net income from agriculture. The economic impacts of soil degradation might be moderate when viewed from global perspective, but the impacts are severe in selected areas. [5]. Policy reforms and public investments are needed in those areas to reduce soil degradation, improve rural income, and enhance food security [6]. The most severe aspect of soil degradation on Asian lands is soil salinity that has occurred through the accumulation of salts, mainly deposited from saline irrigation water or through

mismanagement of available water resources [7]. The historical lands of Iran, Pakistan, Afghanistan, India and China are also subject to ancient and ongoing soil degradation processes which are subtle in some areas but evident and drastic in others.

The effect of different water qualities on soil health and crop yield is governed by many factors including water quality (amount and kind of salts), soil texture, crop species, climate (rainfall and temperature) and management practices. [8,9]. The results of a long term study of the effects of TW of EC 2.40 dSm⁻¹ and SAR 9.20 (m mole L⁻¹)^{0.5} on soil properties showed that soil EC increased from 2.98 to 4.55 dSm⁻¹ and SAR increased from 10.8 to 18.9 (m mole L⁻¹)^{0.5} after 7 years[9]. Similar results have been reported by other [10]. The negative effects of brackish water on the yields of wheat grain, paddy, berseem, maize fodder [9, 11] and sunflower [12] have been reported. The wheat grain yield from field irrigated with TW was found to be more affected by the sodicity compared with salinity of water [13].

Little attention is paid by farming community to keep the land productive in Pakistan. The pre-requisite for the safe use of poor quality water is to know the amount and kind of salts in irrigation water. Limited data on the quality status of TW waters in Lahore [14], Kasur [15], Gujarat [16], Rawalpindi [17], Jhang [13], Sahiwal [3] and other districts of Punjab is available. But systematic and comprehensive information on TW water quality and its implications on soil and crop productivity are not sufficient in the south Punjab of Pakistan. Therefore, a detailed survey of district Sahiwal was carried to collect TW water and soil samples along with yields data of crops grown in the area. The detailed information on ionic concentrations of water samples at village levels and remedial measures for the safe use of brackish water has been submitted elsewhere. In this report we will discuss (i) overall quality status of TW waters for irrigation and, (ii) quantification the effect of TW water irrigation on soil quality and some selected crops yields in the study area.

2. METHODOLOGY

2.1. Description of Study area

Sahiwal district (30°39'52"N 73°6'30"E) occupies central part of Punjab Province, Pakistan, and situated in a semiarid region that is flood irrigated with canal and tubewell waters. The parent material is alluvium formed by the Ravi and Sukh Bias rivers. It is highly productive and texture varies from silt loam to clay loam and clay type. Saline-sodic soils also exist in the district. The major crops grown are cotton, rice, and maize, sugar cane in Kharif and wheat and potato in Rabi season with cropping intensity > 200%. The annual rainfall ranges from 150 to 200 mm. The average maximum temperature during summer usually remains 40-45 °C. The winter is cold and dry and the average minimum temperature is 4-5 °C during month of January.

2.2. Sampling

Out of 531 villages of Sahiwal district, 261 villages were randomly sampled for 529 TW in Canal Command Area (CCA) during 2012 and 2013 (Fig. 1). The CCA comprised of nine distributaries of Lower Bari Doab Canal (LBDC); four on right side (RD) namely, 4R, 5R, 6R and 7R; five on left side (LD) namely, 5L, 9L, 11L, 12L and 14L. Out of total 261, 105 villages were sampled from RD and 156 from LD. The samples were collected in plastic bottle after 30 minutes of TW operation. The depths of TW ranged from 61 to 152 m. Twenty composite soil samples, 10 from each Chichawatni and Sahiwal tehsils, were collected at 0 to 15 cm depth from the selected farmer fields irrigated with TW and CW to monitor the effect of poor quality water on soil health. The basis of soil and crop yield samplings was to cover representative area under three prevailing crop rotations namely; wheat-cotton-wheat, wheat- rice -wheat and maize- potato-maize. Total cultivated area of Sahiwal district under above mentioned crops was 766188 acres.




Fig. 1 Map showing locations of tubewell water samples in LBDC command of district Sahiwal, Punjab, Pakistan.

2.3. Soil and Water Analyses

The analytical work was carried out at Soil and Water Testing Laboratory, Sahiwal. Soil samples were air-dried, ground and passed through 2 mm sieve and preserved in plastic bottles for analyses. Soils were analysed for pH_s, electrical conductivity of the saturated paste (EC_e), and water samples were analysed for EC, cations (Ca²⁺ + Mg²⁺, Na⁺) and anions (CO₃²⁻, HCO₃⁻) by methods described by [18]. Total Dissolved Salts (TDS), Residual Sodium Carbonate (RSC),

Sodium Adsorption Ratio (SAR) of water were computed [1] as given below and the criterion for the suitability of irrigation water is given in Table 1

$$\text{TDS (mg L}^{-1}\text{)} = \text{EC (dSm}^{-1}\text{)} \times 640.$$

$$\text{RSC (me L}^{-1}\text{)} = (\text{CO}_3^{2-} + \text{HCO}_3^-) - (\text{Ca}^{2+} + \text{Mg}^{2+}), \text{ where cations and anions in me L}^{-1}$$

$$\text{SAR (mmol L}^{-1}\text{)}^{0.5} = \text{Na}^+ / [(\text{Ca}^{2+} + \text{Mg}^{2+})/2]^{0.5}, \text{ where cations and anions in me L}^{-1}$$

Table 1. The criterion used for suitability of irrigation water [1].

Parameters	Fit	Marginally Fit	Unfit
EC (dSm-1)	<1.0	1.0 – 1.25	> 1.25
RSC(me L-1)	<1.25	1.25 – 2.25	> 2.25
SAR(mmol L-1) ^{0.5}	<6	6 – 10	>10

74

2.4. Estimation of Soluble Salts Addition to Soil with TW Water Irrigation

The quantities of soluble salts added into soil with the irrigation of TW of different qualities in the study area were estimated to assess their effects on soil quality. The basic equation [19] used to calculate the amount of salts addition with irrigation water is as under:

$$\text{TDS (tons per acre } \approx \text{ft)} = \text{TDS (mg L}^{-1}\text{)} \times 0.00136$$

$$\text{TDS (tons per hectare)} = \text{TDS (tons per acre)} \times 2.471$$

In the sampling area the TDS of TW waters ranged from 218 to 3309 mg L⁻¹ and this range was used to calculate that how much salts were added into soil with the application 0.30 m of water. Similarly, salts added to soil under different crops were calculated by considering consumptive use (CU) of wheat (425mm), cotton (750mm), rice (1500mm), spring maize (1200mm), autumn maize (750mm) and potato (900mm) [20] The highest level of water salinity (TDS 3309 mg L⁻¹) was used in these estimations. To make the understanding clear, the calculation for wheat is demonstrated.

$$\text{CU of wheat} = 425\text{mm} = 1.4167 \text{ ft.}$$

$$\text{TDS of TW water used} = 3309 \text{ mg L}^{-1}.$$

Putting values in the above equation we get

TDS (tons per acre ft of water.) = $3309 \times 0.00136 = 4.50$

As the CU of wheat is 1.4167 feet, So TDS will be $4.50 \times 1.4167 = 6.375$ tons per acre = $6.375 \times 2.471 = 15.753$

2.5. Crop Yield Data and Statistical analysis

The yield data of wheat, cotton, maize and potato crops were recorded, according to the standard procedures, from 10 farmer's fields from each tehsil irrigated with CW and TW to compare the effects of brackish water on the yields. Some information on inputs regarding source of irrigation and fertilizers type (organic and inorganic), and rate of fertilizers application were also collected. The scarcity of CW forced the farmers to use poor quality TW water and they mix CW and TW waters to raise the crops. It was noted that >90% farmers used Diammonium phosphate (DAP) as source of phosphorus in comparison of Single super phosphate (SSP). The sources of nitrogen were urea, Calcium ammonium nitrate (CAN). Now Ammonium sulphate (AS) is available in the market to replace urea and farmers are fully convinced as this fertilizer is acidic in reaction due to pH 4. The sources of potassium were potassium sulphate (SOP) and muriate of potash (MOP) and its use was limited to maize and potato crops. The maize and potato growers generally used 1.5 times nitrogen against the departmental recommendations. The comparisons among the treatment means were made by calculating standard error of the mean values [21].

3. RESULTS AND DISCUSSION

3.1. Quality Status of Tubewell Waters of District Sahiwal

Data (Fig.2) on characterization of TW water samples based on EC, RSC and SAR showed that 71% TW were saline and >20% samples were sodic in nature [13,14,15]. The quality of TW water in RD was comparatively better than in LD due to its location between LBDC and Ravi River. The TW situated along LBDC had sweet water due to the seepage of CW. The more TW water samples on LD were brackish than RD and about 6% of the total TW samples had extreme values of EC, RSC and SAR. The use of such waters caused soil salinity and affected crops yields. The yields reductions were not to that extent because of sandy aquifer without any marked clay layers [22], use of flood irrigation of TW with dilution of CW and sowing of semi tolerant crop like wheat. Based on soil and water analyses, the farmers were advised proper management strategies for sustainable crop production. It was realized that those farmers who followed the departmental recommendations had harvested 3.30 t ha^{-1} wheat in Chichawatni and 4.13 t ha^{-1} in Sahiwal where most of times brackish TW was used to irrigate the fields.

The concentrations of chemical constituents in water from the LBDC aquifer system vary both depth and location wise. The deep groundwater quality investigations carried out by [23] showed that the distribution of saline and fresh

groundwater zones in Bari Doab is a result of past and present hydrologic, climatic, and topographic factors. Among these, the present and former positions of stream channels, representing sources of recharge, the high bluffs of the bar uplands in the upper part of the Bari Doab, and differences in the permeability within the alluvial aquifer are the most important.




Fig.2. Characterization of tube well waters (n=529) on the basis of EC, RSC and SAR of district Sahiwal.

3.2. Impact of Tubewell Waters Irrigation on Soil Quality

The farmers of district Sahiwal used poor quality TW waters having EC 0.34 - 5.17 dSm⁻¹ with SD 0.88 (TDS ranged from 218 to 3309 mg L⁻¹) for crop production due to the scarcity of CW. The use of these water qualities contributed to the soil salinity [24] leading to soil degradation. It was estimated that a huge amount of salts were added into soil with the application of 0.30 m irrigation water of different qualities (Fig. 3). Similarly, it was calculated that 55.599 t ha⁻¹ soluble salts were deposited in soil under rice crop with the application of TW having TDS 3309 mg L⁻¹ (Table 2).

The effects of brackish irrigation water on soil salinity due to continuous use of poor quality TW was evaluated at farmer's field in two tehsils of district Sahiwal during 2012 and 2013 (Table 3). The EC of TW used for irrigation ranged from 1.00 to 3.62 dSm⁻¹. It was observed that soil EC increased from 2.06 to 3.62 dSm⁻¹ in Chichawatni and 2.40 to 3.97 dSm⁻¹ in Sahiwal tehsil with continuous use of TW. This data supported that substantial amounts of soluble salts were added to soils with the application poor quality TW irrigation as given in Fig. 3 and Table 2. However, the increase in soil salinity

was not to that extent as it was expected. This was due to leaching of salts with the application of heavy flood irrigations by the farmers, light to medium textured soils and rainfall (457 mm) during the rainy season [3]. It means the significance of rainfall and leaching fraction in maintaining the soil health while using brackish water should be duly accounted for [8]. Similarly, the results of a long term study conducted on the effects of brackish TW on soil quality showed that after growing 12 crops following berseem-rice-wheat- maize fodder-berseem crop rotation, the non-saline and non-sodic soil was converted to sodic soil (SAR 18.9) by the use of TW [9]. Similar results were also reported by others [10, 24].




Fig.3. Quantity of soluble salts added to soil with tubewell waters irrigation of 0.30 m depth of TDS 218 to 3309 mg L⁻¹

Table 2. Quantity of soluble salts added to soil with tubewell water irrigation having 3309 mg L⁻¹ TDS under different crops grown in district Sahiwal, Punjab, Pakistan

Crop	Consumptive use (mm)	Quantity of salts added (t ha ⁻¹)
Wheat	425	15.753
Cotton	750	27.801
Rice	1500	55.599
Potato	900	33.361
Spring Maize	1200	44.480

Table 3. Range, mean and standard deviation of ECe at farmer's fields irrigated with canal and tubewell water in two tehsils of Sahiwal district during 2012-13

Location	Source of irrigation	*Range	Mean	Standard Deviation
Chichawatni Tehsil	Canal water	1.75-2.41	2.06	0.24
	^a Tubewell water	2.99-4.01	3.62	0.41
Shiwal Tehsil	Canal water	1.95-2.80	2.40	0.36
	^b Tubewell water	3.65-4.36	3.97	0.27


*Mean value of 10 farmer's fields in each tehsil; ^aEC ranged from 1.00 to 2.82 with mean 1.9 ± 0.79 ; ^bEC ranged from 1.12 to 3.62 with mean $2.14 \pm 0.0.99$.

3.3. Impact of Tubewell Water Irrigation on Crop Yield

The effect of water quality on the yields of wheat, cotton, maize and potato crops was evaluated at farmer's fields of Chichawatni (Fig.4) and Sahiwal tehsils of the district (Fig. 5). The yields of crops reduced from 3 to 15% in the fields where usually TW was used compared to CW irrigated plots. The yields of crops were not severely affected by TW water having EC up to 5.17 dS m^{-1} in LD because the lithologic logs of bore holes have shown sandy aquifer without any marked clay layers[22] and also due to the management practices adopted by the farmers[25]. The yield decrease is very limited because the farmers did not use TW water alone. They occasionally mixed a limited quantity of available CW with TW water. Secondly, the soil texture of the study area is loam to loamy sand. Thirdly, the progressive growers of maize and potato used heavy inputs in terms of FYM, DAP, SSP AS, urea, gypsum and Sulphur to harvest 16 to 17 t ha⁻¹ potato and 8 to 10 t ha⁻¹ spring maize. It was noted during survey that the farmers harvested optimum yields of crops with brackish TW following viable management practices. For example, the addition of farm manures, use of balanced fertilizer, proper type of fertilizers (use of single supper phosphate on sodic soils) and light irrigation of poor quality water proved better. Furthermore, the response of crops to different water qualities differed greatly due to different salt tolerance limits of salts by crops [26]. The use of brackish TW water ($\text{EC } 2.40 \text{ dSm}^{-1}$; $\text{SAR } 9.20 (\text{mmol L}^{-1})^{0.5}$; $\text{RSC } 5.70 \text{ me L}^{-1}$) on


168 crops grown under berseem-rice-wheat-maize fodder-berseem rotation on permanent layout during 1987-88 to 1993-94
 169 reduced crops yields from 3 to 36% over CW [9]. Similar results were also reported by others [10, 25]. The soluble salts
 170 exerted both general and specific effects on plants which reduced crops yields.

171



172 Fig.4. Effect of water quality on the yields of crops at farmer's fields in Chichawatni tehsil of District Sahiwal. EC of TW
 173 ranged from 1.00 to 2.82 dS m⁻¹; SD for wheat, cotton, maize and potato yields in canal irrigated fields were ± 0.61 ,
 174 ± 0.77 , ± 0.19 and ± 1.85 ; and in tubewell irrigated fields were ± 0.57 , ± 0.72 , ± 0.13 and ± 1.20 , respectively.

175



176

177 Fig.5. Effect of water quality on the yields of crops farmer's fields in Sahiwal tehsil of District Sahiwal. EC of TW ranged
 178 from 1.12 to 3.62 dS m⁻¹; SD for wheat, cotton, maize, potato yields in canal irrigated fields were ± 0.91 , ± 0.73 ,
 179 ± 0.65 and ± 1.80 ; and in tubewell irrigated fields were ± 0.80 , ± 0.63 , ± 0.12 and ± 1.79 , respectively.

180

4. Conclusion

Data on quality status showed that more than 71% TW were saline and >20% were saline or sodic in nature. The concentrations of chemical constituents in water from the LBDC aquifer system vary both depth and location wise. It was estimated that application of TW irrigation having different salinities added substantial amount of soluble salts into soil that affected the soil quality. Resultantly, the yields of different crops reduced from 3 to 15% with the use of poor quality irrigation water. **Presently the farmers are using brackish TW water with CW but the continuous use of poor quality irrigation water would have serious consequences for sustainability of soil and crop productivity in the area.**

References

1. Richards LA. Diagnosis and Improvement of Saline and Alkali Soils. Handbook 60, USDA, Washington D.C; 1954.
2. Basharat M. Spatial and temporal appraisal of groundwater depth and quality in LBDC Command-Issues and Options. Pak. J. Engg. & Appl. Sci. 2012; 11:14-29.
3. Ishaq M. Annual Report of Soil and Water Testing Laboratory, Sahiwal. Directorate of Soil Fertility Research Institute, Lahore, Punjab, Pakistan; 2013-14.
4. Lal R. Cropping systems and soil quality. J. Crop Prod. 2003; 8: 33-52.
5. Faeth P. Building the case for sustainable agriculture. Environment. 1994; 36: 16-20.
6. Rosegrant MW, Paisner MS, Meijer S. 2003. The future of cereal yields and prices: Implications for research and policy. J. Crop Prod. 1994; 9: 661-690.
7. Klarin NG, Tateishi R, Harahshah H. Degradation of the dry lands of Asia. Center for Environmental Remote Sensing Chiba Univ., Japan; 1999.
8. Singh RB, Minhas PS, Gupta PK. Effect of high salinity and SAR waters on salinization, sodication and yield of pearl-millet and wheat. Agricultural Water Management. 1992; 21: 93-105.
9. Rashid M, Shakir MY, Jamil M, Ishaq M. Cyclic use of canal and tubewell waters for irrigation. J. Agric. Res. 1996; 34: 381-387.
10. Jakhar RA, Iqbal J, Hussain N, Dogar MS. A review of water quality research in Pakistan. Pakistan J. Soil Sci. 1994; 9: 10-17.

- 206 11. Rashid M, Shakir MY, Jamil M. Effect of brackish water on crop yields and properties of a soil treated with
207 amendments. Pakistan J. Soil Sci., 1994; 9: 87-90.
- 208 12. Hussain SA, Akhtar J, Haq MA, Ahmad R. Growth, yield and ionic concentration of two sunflower (*Helianthus*
209 *annuus* L) genotypes exposed to brackish water irrigation. Soil Environ. 2011; 30: 58-63.
- 210 13. Hannan A, Ahmad M, Niaz A, Ali I, Waheed T. Ground water quality characterization and its correlation with
211 wheat yield. Soil Environ. 2010; 29: 172-176.
- 212 14. Ali MS, Mahmood S, Chaudhary MN, Sadiq M. Irrigation quality of ground water of twenty villages in Lahore
213 district. Soil Environ. 2009; 28: 17-23.
- 214 15. Mehboob I, Shakir, MS, Mahboob A. Surveying tube well water suitability for irrigation in four tehsils of district
215 Kasur. Soil Environ. 2011; 30:155-159.
- 216 16. Zahid P, Kazmi SSH, Gill KH. Characterization of irrigation quality of ground water in Gujarat district. Pakistan J.
217 Soil Sci. 2003; 22: 48-54.
- 218 17. Rizwan K, Mahmood T, Abbas Z, Dilshad M, Lone MI. Ground water quality for irrigation in Rawalpindi district.
219 Pakistan J. Soil Sci. 2003; 22: 43-47.
- 220 18. Page AL, Miller RH, Keeney DR, editors. 1982. Soil Analysis. Part 2. Chemical and Microbiological Properties.
221 Agronomy Monograph No.9, 2nd ed. Madison, WI, USA, 1982
- 222 19. Rhoades JD, Kandiah A, Mashali AM. The use of saline waters for crop production. Irrigation and Drainage Paper
223 48, FAO, Rome; 1992.
- 224 20. Anonymous. Consumptive use of water for crops in Pakistan: Final Technical Report PL – 480; PARC, Islamabad,
225 Pakistan; 1982
- 226 21. Duncan BD. Multiple Range and Multiple F – Test. Biometrics. 1955; 11: 1-42.
- 227 22. Shakir AS, Rehman H, Khan NM, Qaz AU. Impact of canal water shortages on groundwater in the Lower Bari
228 Doab Canal system in Pakistan. Pak. J. Engg. & Appl. Sci. 2011; 9: 87-97.

229 23. Greenaman DW, Swarzenski WV, Bennett GD. Groundwater Hydrology of the Punjab, West Pakistan with
230 emphasis on problems caused by canal irrigation. West Pakistan Water and Power Development Authority,
231 WASID Bulletin 6; 1967

232 24. Qureshi RH, Barrett-Lennard EG. Saline Agriculture for Irrigated Land in Pakistan: A handbook, ACIA R,
233 Canberra, Australia; 1998.

234 25. Saleem Z, Rashid M, Ishaq M. Growing crops with brackish water without affecting the soil health. Pakistan J. Soil
235 Sci. 1993; 8: 41-46.

236 26. Ayers RS, Westcot DW. Water quality for agriculture. Irrigation and Drainage Paper 29, Rev.1. FAO, Rome;
237 1985.

238