

1

2 **A REVIEW OF HOST-PARASITE RELATIONSHIPS**

3

4 **ABSTRACT**

5 Host parasite relationships result from prolonged associations between
6 organisms living in a given environment. The nature and extent of the
7 association will determine the type of relationship existing between the co-
8 habiting organisms. Host-parasite associations usually give rise to four main
9 relationships namely parasitism, mutualism, commensalism and phoresis.

10 Key words: host, parasite, relationship, associations.

11

12

13 **SECTION ONE**

14 **1.1 INTRODUCTION**

15 Parasitology has been broadly defined as “a study of symbiosis, or literally
16 speaking “living together”[1] . Naturally speaking, parasitology is defined as
17 the scientific study of parasites. What then are parasites? A parasite is defined
18 by the advanced English dictionary as “an animal or plant living in or on
19 another and getting its food from it”[2].

20 Taking the broad definition of parasitology into consideration, the word
21 “symbiosis” as used, raises another question. The word itself is taken from
22 Ancient Greek language meaning “living together” [3] ie close and long term
23 interactions between two or more different biological species. In 1877, Albert
24 Bernard Frank used the word symbiosis to describe the mutualistic relationship
25 existing among lichens. This usage conforms to the definition of symbiosis
26 given by the Oxford advanced English dictionary depicting people living
27 together in a community. In 1879, Heinrich Anton de Bary, a German
28 Mycologist defined symbiosis as “living together of unlike organisms” [4, 5].

29 The meaning of the word symbiosis has become controversial among
30 Scientists. Whereas some Scientists believe symbiosis should refer to
31 relationships that are beneficial to both parties, (ie mutualistic relationships),
32 others believe it should apply to any type of persistent biological interactions [6,
33 7]. Consequently, four different types of symbiotic relationships have emerged
34 namely:

- 35 1. Parasitism
- 36 2. Mutualism
- 37 3. Commensalism
- 38 4. Phoresis

39 These relationships will be discussed at length as this essay continues.

40 1.2 TYPES OF SYMBIOSIS

41 **Obligate Symbiosis:** This is the type of relationship where both symbionts
42 entirely depend on each other for survival. For example, many lichens consist of
43 fungal and synthetic symbionts that cannot live on their own [6].

44 **Facultative Symbiosis:** This is the type of relationship where the organisms
45 may not necessarily live with each other in order to survive

46 **Ecosymbiosis:** Here, one organism lives on another eg mistletoe

47 **Endosymbiosis:** This is where one partner or symbiont lives inside the other eg
48 lactobaccili and other bacteria on humans or symbiodinium in corals [8].

49 **Conjunctive Symbiosis:** This is the type of relationship in which the two
50 organisms have bodily union ie attached to each other. If the opposite is the
51 case, it is called **disjunctive** symbiosis [9].

52 1.3 HOSTS

53 The word host as applied to parasitology has been defined as “an organism
54 infected with or is fed upon by a parasitic or pathogenic organism (eg
55 nematodes, fungi, virus etc). It is also described as an animal or plant that
56 nourishes or supports a parasite. The host does not benefit but instead is
57 harmed by the association.

58 1.4 TYPES OF HOSTS

- 59 1. **Definitive or primary host:** An organism in which a parasite reaches
60 sexual maturity eg the mosquito is the definitive host for the malaria
61 parasite, *plasmodium*.
- 62 2. **Intermediate (alternative or secondary) host:** An organism in which a
63 parasite develops but does not attend sexual maturity eg humans and
64 other vertebrate animals are intermediate host for *plasmodium*.
- 65 3. **Paratenic host:** A host which may be required for the completion of a
66 parasites life cycle but in which no development of the parasite takes
67 place eg the unhatched eggs of nematodes are sometimes carried in a
68 paratenic host such as a bird or a rodent. When a predator eats the
69 paratenic host, the eggs are ingested and it becomes infected.
- 70 4. **Accidental host:** One that accidentally harbours an organism that is not
71 ordinarily parasitic in the particular species.
- 72 5. **Dead-end host:** This is the host in which the disease cannot be
73 transmitted to another animal. Any host organism from which a parasite
74 cannot escape to continue its life cycle. Eg humans are dead end hosts for
75 trichinosis because the larvae encysted in the muscle and human flesh are
76 unlikely to be a source of food for other animals susceptible to the
77 parasite.
- 78 6. **Predilection host:** Is the host most preferred by the parasite.
- 79 7. **Reservoir host:** An animal or species that is infected by a parasite and
80 which serves as a source of infection for humans or other species.

81 **8. Transfer or transport host:** A host which is used until the appropriate
82 host is reached, but is not necessarily to complete the life cycle of the
83 parasite

84

85 **1.5 HOST-PARASITE SPECIFICITY**

86 A parasite can infect one or a limited number of hosts at a given time ie most
87 parasites occur on a restricted number of hosts. This gives rise to the concept of
88 specificity. Host specific parasites generally have a major host and then a few
89 less frequently used hosts in the absence of the major one [10, 11]. Even among
90 parasites that do not discriminate among hosts, there is preference for some
91 species hosts above others [12-14]. It is said that many parasite groups have a
92 drift toward greater host specificity. Host specificity is the characteristic of a
93 parasite that renders it capable of infecting only one or more specific hosts at a
94 time.

95 **1.6 HOST-PARASITE EVOLUTION/SPECIFICITY**

96 Natural selection tends to occur and favour the specialization of parasites to
97 their local environment or hosts [15-17]. The parasite ecosystem is a world of
98 competition between organisms where there is survival of the fittest. Thus the
99 most adapted and fitted host or parasite exists in greater abundance than the
100 least fitted. Host specialization is said to be promoted by host-dependent fitness

101 tradeoffs which is dependent on the relative availability and predictability of
102 hosts [19, 16, 20, 21, 12, 22, 23].

103 A parasite should specialize if the advantages of using one single host
104 species in a profitable manner outweigh the benefits of interacting less
105 profitably with several less frequent host species [19, 16, 20, 21, 22, 23]. In
106 other words, lack of adequate hosts will promote parasite generalization [11a],
107 while abundance of hosts will make parasites to specialize to the specific
108 environmental conditions [13, 11b].

109 It is believed that host parasite interactions, and thus host specificity take
110 place simultaneously at several “host” levels. This is probably while such
111 interactions are especially difficult to explain. Studies carried out by Georgi et
112 al [24], using ectoparasitic mites, (*Spinturnicidae*) which infest colonial bats,
113 revealed that parasite specificity may be mediated by three main mechanisms:

114 **1. Dispersal capacity of the parasite** which depends on the number of
115 hosts it can physically encounter during its life.

116 **2. Host preference**

117 **3. Ability to successfully transmit** and establish a population on a new
118 host.

Considering the third mechanism, it is said that highly specific parasites are expected to exhibit a higher reproductive success or survival on traditional or native host species than on less closely related ones [25-27].

SECTION TWO

2.1 PARASITISM

Parasitism is defined as a relationship in which one of the participants (the parasite), either harms its host or in some sense lives at the expense of the host [1]. Wikipedia describes parasitism as a non mutual symbiotic relationship between species, where one species, (the parasite), benefits at the expense of the other (the host). Traditionally, a parasite referred primarily to organisms which were visible to the naked eye, otherwise known as macro-parasites (eg helminthes), but nowadays, parasites include microscopic organisms such as viruses, and bacteria [28] which are referred to as micro-parasites. The word “parasite” was derived from a Latin word “parasitus” which means “one who eats at the table of another” [29].

Although parasites may inflict harm on their hosts, it is not in the best interest of the parasite to kill its host. A parasite which kills its host has

139 invariably committed “suicide”. Some of the ways parasites inflict harm on their
140 hosts include:

- 141 • Boring a hole into the host eg Schistosomes
- 142 • Digging into hosts skin or other tissues eg hookworm larvae
- 143 • Stimulation of damaging inflammatory or immune response eg
144 microfilariae
- 145 • Robbing of the host of nutrients eg tapeworm, hookworm
- 146 • A combination of two or more of the above conditions

147 Unlike predators, parasites are usually smaller than their hosts and will often
148 live in or on their hosts for an extended period of time. Both parasitism and
149 predators are special cases of consumer resource interactions [30].

150 Parasites display a high degree of specialization, and reproduce at a
151 faster rate than their hosts. Examples of parasitism include interactions
152 between vertebrate hosts and diverse animals such as tapeworm, flukes, the
153 plasmodium species and fleas.

154 **2.2 TYPES OF PARASITES**

- 155 • **Ectoparasites:** Are parasites living on the surface of their hosts eg bed
156 bugs, mites, ticks etc
- 157 • **Endoparasites:** Are parasites living within the body of their hosts eg
158 schistosomes, tape worm, ascaris etc

159 • **Obligate parasites:** Cannot complete their life cycle without spending at
160 least part of the time in parasitic relationship eg *plasmodium*. However
161 many obligate parasites have free living forms which can exist outside the
162 host for some period of time in the external environment in a protective
163 egg shell or cyst eg hookworm larvae, ascaris, *Entamoeba histolytica*.

164 • **Facultative parasites:** These are not normally parasitic but become so
165 when they are accidentally eaten or enter a wound or other body orifices
166 eg certain free living amoeba such as *N.fowleri* and free living nematodes
167 of the genus *micronema* [31]. Infection of humans by any of these
168 facultative parasites is always very fatal.

169 • **Accidental/incidental parasites:** This occurs when a parasite enters or
170 attaches to the body of species of host different from its usual preferred
171 host eg nematodes parasitic in insects can live for a short time in the
172 intestines of a bird or rodent. Fleas can live for a while in dogs or
173 humans. Accidental parasites usually do not survive in the wrong host but
174 in some cases they can be extremely pathogenic eg *Toxicara*, *baylis*
175 *ascaris*.

176 Parasitism usually results from a long history of evolutionary symbiosis
177 between the parasites and the hosts in which both parties are fully
178 adapted. It is no wonder then why accidental parasitism is fatal for both
179 host and the parasites because neither of the two parties is adapted for the
180 co-existence.

- 181 • **Permanent parasites:** These are parasites which live their entire adult
182 lives within or on their hosts
- 183 • **Temporary or intermittent parasites:** These feed on their hosts and
184 then leave eg mosquitoes, bed bug. They are also called micropredators
185 because they also prey on several different hosts or the same host at
186 several discrete times.
- 187 • **Mesoparasites:** Are those parasites living in an intermediate position ie
188 half ectoparasites and half endoparasites
- 189 • **Epiparasites:** Are parasites which feed on other parasites. This is
190 sometimes referred to as hyperparasitism eg a protozoan living in the
191 digestive tract of a flea living on a dog.
- 192 • **Social parasites:** Are parasites which take advantage of interactions
193 between members of a social group of organisms such as ants or termites
194 eg *Phengaris arion*, a butterfly whose larvae employ mimicry to
195 parasitize certain species of ants [32].

196 **2.3 TYPES OF PARASITISM**

197 **KLEPTOPARASITISM:** In this type of relationship, parasites
198 appropriate the food gathered by the host eg brood parasitism practiced
199 by many species of cuckoo and cowbird which do not build nests of their
200 own but rather deposit their eggs in nests of other species and abandon
201 them there. The host behaves as a “baby sitter” as they raise the young

202 ones as their own. If the host bird ventures to remove the Cuckoos eggs,
203 some cuckoos will return to attack the nest to compel the host bird to
204 comply with their wish [33]. In the case of the cowbird, the host's brood
205 is not necessarily harmed but this is not so with the cuckoo which may
206 remove one or more of the host's eggs to avoid detection or the young
207 cuckoo may heave the hosts eggs and nestlings out of the nest entirely.
208 What a wicked act!

209 **INTRASPECIFIC SOCIAL PARASITISM:** This may occur in the
210 form of parasitic nursing where some members of the relationship take
211 milk from unrelated females eg in wedge capped capuchins, higher
212 ranking females sometimes take milk from low ranking females without
213 any reciprocation. That is to say high ranking females benefit at the
214 expense of the low ranking ones [34].

215 **CHEATING OR EXPLOITATION:** Parasitism can also occur as
216 isolated cheating or exploitation among more generalized mutualistic
217 interactions eg broad classes of plants and fungi exchange carbon and
218 nutrients in common mutualistic mycorrhizal relationships. However,
219 some plant species known as mycohetrotrophs "cheat" by taking carbon
220 from a fungus without donating it.

221 **PARASITOIDS:** These are organisms whose larval development takes
222 place inside or on the surface of another organism (the host) leading to the

death of the later [35]. This differentiates parasitoids from true parasites which normally do not kill their hosts. Thus parasitoid relationship is similar to predation where the host is always killed. Parasitism differs from parasitoid relationship in the sense that parasitoids generally kill their hosts [36]. Parasitoidism occurs in a similar variety of organisms to that in which parasitism occurs. A parasite can reduce the host's biological fitness in a variety of ways:

- parasitic castration of the host ie impairment of the hosts secondary sex characteristics
- modification of the hosts behavior

Parasites can also increase their own fitness by exploiting the host for resources necessary for their own survival such as food, water, heat, habitat and transmission.

ADELPHO-PARASITISM: An adelpho parasite is one in which the host species is closely related to the parasite, often being a member of the same family or genus eg the citrus blackfly parasitoid, *Encarsia perplexa* whose unmated females may lay haploid eggs in the fully developed larvae of their own species. These result in the production of male offsprings. Secondly, the marine worm *Bonellia viridis* has a similar reproductive strategy, although the larvae are planktonic [37].

AUTOINFECTION: Is the infection of a primary host with a parasite, particularly a helminth, in such a way that the complete life cycle of the parasite occurs in a single organism without passing through other hosts i.e the primary host is at the same time the secondary host. Examples include *Strongyloides stercoralis*, *Enterobius vermicularis*, *Taenia solium* and *Hemenolepsis nana*. *Strongyloides* for example can cause premature transformation of a non infective larva to infective lava, which can then penetrate the intestinal mucosa (internal autoinfection) or the skin of the perineal area (external autoinfection). Thus infection can be maintained by repeated migratory cycle for the rest of the person's life.

2.4 HOST DEFENSES AGAINST PARASITES

The host responds to parasitism in a variety of ways ranging from morphological to the behavioural. Some of these ways include:

- 1. Toxins:** Some plants produce toxins which are antiparasitic to inhibit the growth of parasitic fungi and bacteria [38].
- 2. Immune systems:** Vertebrate animals develop complex immune systems which fight parasitic organisms to get rid of them. In humans parasitic immunity involves IgE.
- 3. Behavioural defenses:** For example sheep avoid open pastures during spring when roundworm eggs are known to accumulate enmasse over

the previous years. Secondly some infected fruit flies ingest alcohol as a form of self medication against blood borne parasites [39].

2.5 EVOLUTION OF PARASITES

Biotrophic parasitism is said to be a common mode of life that has arisen independently many times in the course of evolution. It is also believed that as many as half of all animals have at least one parasitic phase in their life cycles [40] and it is also frequent in plants and fungi. Secondly, almost all free living animals are hosts to one or more parasitic organisms at one time or another [40]. A study [41] has shown that holes in the skull of several specimens might have been caused by Trichomonas-like parasites.

Furthermore, parasites have been known to evolve in response to the defense mechanisms of their hosts. As a consequence of their host defenses, some parasites evolve adaptations that are specific to a particular host taxon, specializing to the point where they infect only a single species. Such parasites may pay dearly over time if the host species become extinct. Consequently, many parasites evolve to infect a variety of more or less closely related host species with different success rates.

282 Host defenses also evolve in response to parasitic attacks. In theory,
283 parasites may have advantage in this evolutionary arms race because parasite
284 generation time is commonly shorter ie hosts reproduce less quickly than
285 parasites and therefore have fewer chances to adapt than their parasites do over
286 a given range of time.

287 In some cases a parasite may co-evolve with its host taxa. It is said that long
288 term co-evolution may lead to a relatively stable relationship tending towards
289 commensalism or mutualism since it is in the best interest of the parasite that
290 the host remains alive. A parasite may evolve to become less harmful for its host
291 or a host may evolve to cope with the unavoidable presence of a parasite- to the
292 extent that the parasites absence causes the host harm. For example it is known
293 that animals infected with parasitic worms are often clearly harmed, such
294 infections may also reduce the prevalence and effects of auto immune disorders
295 in animal hosts, humans inclusive [42].

296 Competitions between parasites often occur and this tends to favour faster
297 reproducing, and hence more virulent parasites. Parasites which kill the host in
298 the course of their life cycle, in order to enter a new host, evolve to be more
299 virulent or even change the behavior or other properties of the host to make it
300 more vulnerable to predators. Parasites that reproduce largely to the offspring of
301 the previous host, tend to become less virulent or mutualist, so that its hosts
302 reproduce more effectively [43].

The presumption of shared evolutionary history between parasites and hosts can sometimes explain how host taxa are related. For instance, the relationship between flamingos and storks or their relatives and ducks, geese and their relatives has been controversial. It has been said that the fact that flamingos share parasites in common with ducks and geese is evidence or proof that these groups may be more closely related to each other than either is to the storks.

Parasitism has been used to explain the evolution of secondary sex characteristics seen in breeding males throughout the animal kingdom eg the plumage of male peacocks and manes of male lions. According to this theory, female hosts select males for breeding based on such characteristics because they indicate resistance to parasites and other diseases.

2.6 PARASITES ADAPTATIONS

Parasites are adapted to infect hosts that exist within their same geographical area (sympatric host) more effectively than hosts found outside their own geographical area (allopatric hosts). This phenomenon is said to support the so called “Red Queen hypothesis” which states that interactions between species (such as hosts and parasites) lead to constant natural selection for adaptation and counter adaptation [44]. Experiments conducted by the later authors, using two snail populations from different sources substantiated the fact that parasites were more infective to sympatric hosts than they were to allopatric hosts ie

324 although the allopatric snails were equally infected, by the digenetic
325 trematodes(parasites), the infectivity was much less when compared to the
326 sympatric snails. Hence the parasites were found to have adapted to infecting
327 local populations of snails [44].

328 **PARASITIC TRANSMISSION**

329 Since parasites inhabit living organisms (hosts), they are faced with numerous
330 problems emanating from the host which will mount many forces aimed at
331 repelling or destroying these invaders. Consequently, parasites develop several
332 strategies to evade these host defense mechanisms to ensure their movement
333 from one host to the other. This is referred to as parasitic transmission or
334 colonization. Some endoparasites infect their host by penetrating its external
335 surface (eg hookworm larvae), while others must be ingested in food by the host
336 (eg *Entamoeba histolytica*). Once they are inside the host, adult endoparasites
337 (eg tapeworm, ascaris) must shed their offspring to the external environment so
338 as to infest other hosts. Many adult endoparasites live in the host's
339 gastrointestinal tract, where the eggs can be shed along with the hosts excreta or
340 faeces. Examples here include tapeworms, thorny headed worms, and most
341 flukes. Some other parasites like malaria parasites (plasmodium) or
342 trypanosomes use insect vectors to transmit their infective stages. Furthermore
343 some larval stages of endoparasites infect sites other than the blood or
344 gastrointestinal tract eg muscle tissue. In such cases, larval endoparasites

require their hosts to be consumed by the next host (predators) in the parasites life cycle in order to survive and to reproduce. On the alternative, some larval endoparasites may shed free living transmission stages that migrate through the host's tissue into the external environment where they actively search for or await ingestion by other hosts. The above mentioned strategies are used variously by larval stages of tapeworms, thorny headed worms, flukes and parasitic round worms.

Furthermore, some ectoparasites eg monogenian worms, depend on direct contact between hosts eg lice. Some ectoparasites may shed eggs which may survive off the host (eg fleas), or wait in the external environment for an encounter with a host (eg ticks). Some aquatic leeches locate hosts by sensing movements and only attach when certain temperatures and chemical cues are present.

Host behavior: Some parasites modify hosts behavior to make transmission to other hosts more likely. For instance, in California salt marshes, the fluke *Euhaplorchis Californienses* reduces the ability of its killifish host to avoid predators [45]. This parasite matures in egrets which are more likely to feed on infected killifish than on uninfected fish. Another example is the protozoan *Toxoplasma gondii*, a parasite which matures in cats, though it can be carried by other animals. Uninfected rats avoid cat odours, where as infected rats are

attracted to cat odours which causes their being easily devoured and hence transmission [46].

2.7 ROLES OF PARASITES IN THE ECOSYSTEM

Although parasites are often omitted in the depiction of food webs, they usually occupy the top position of every food web. Thus they function like keystone species, thereby reducing the dominance of superior competitors and allowing competing species to co-exist. Many parasites require multiple hosts of different species to complete their life cycles and rely on predator-prey or other ecological interactions to get from one host to another. Thus the parasite in an ecosystem reflects the health of that system.

2.8 IMPORTANCE OF PARASITES

- They account for as much as more than half of life's diversity.
- They perform an important ecological role (by weakening prey) that ecosystems would take some time to adapt to.
- Without parasites, organisms may eventually tend to asexual reproduction thereby diminishing the diversity of sexually dimorphic traits [47].
- They provide an opportunity for the transmission of genetic material between species. On rare occasions, this may facilitate evolutionary

385 changes that would not otherwise occur, or taken longer time to occur
386 [28].

387

388

389 SECTION THREE

390 3.1 MUTUALISM

391 Mutualism is the type of relationship where two organisms of different species
392 exist together with each one benefitting. A similar interaction between
393 organisms of the same species is known as cooperation. Mutualism differs from
394 interspecific competition in which each species experiences reduced fitness and
395 exploitation, or parasitism where one species benefits at the expense of the
396 other. Mutualism is one aspect of symbiotic relationships.

397 Examples of mutualism include:

- 398 • Relationship between ungulates (eg bovines) and bacteria within their
399 intestines. The ungulates benefit from the cellulose produced by the
400 bacteria, which facilitates digestion, while the bacteria benefit from the
401 abundant nutrient present in the host environment.
- 402 • Humming bird Hawkmoth and Dianthus. Here, the hawkmoth drinks
403 from the dianthus and in the process helps to bring about pollination.

- The Oxypecker (a kind of bird) and the rhinoceros or zebra. Oxypeckers land on rhinos or zebras and eat ticks or other parasites that live on their skin. The birds get food while the beasts get pest control. Also when there is danger, the oxypecker fly upward and scream a warning which helps the animal to run away.
- The bee and the flower. Bees fly from flower to flower sucking nectar which serves as food. In the process bees bring about cross pollination which benefits the plant.
- The spider crab and the algae. Spider crabs live in shallow areas of the ocean floor, and green brown algae live on the crabs back, thus making the crabs blend in with their environment thereby becoming unnoticeable to predators. The algae gets good place to live while the crab gets camouflage
- Humans and bacteria. A certain kind of bacteria lives in the intestines of man and other animals. The bacteria eat the food humans cannot digest and partially digest it, allowing the human to complete the job. The bacteria benefit by getting food while the human benefits by being able to achieve full digestion.

3.2 IMPORTANCE OF MUTUALISTIC RELATIONSHIPS

1. Mutualistic relationships are important for terrestrial ecosystem function since more than 48% of land plants rely on mycorrhizal relationships with fungi to provide them with inorganic compounds and trace elements.

2. Mutualism is thought to have driven the evolution of much of the biological diversity we see, such as flower forms (which is important for pollination mutualism) and co-evolution between groups of species [48].

Despite its importance in ecology, mutualism has received less attention from Scientist than other relationships such as predation and parasitism [49-50].

3.3 TYPES OF MUTUALISTIC RELATIONSHIPS

Mutualistic relationship has been described as a form of “biological barter” [51] in which species trade resources (eg carbohydrates and inorganic compounds or services, such as gamete, offspring dispersal or protection from predators.

Resource-resource mutualism: This is probably the most common form of mutualism where one type of resource is traded for a different resource. Examples include:

a) Mycorrhizal association between plant roots and fungi in which the plant provides carbohydrates to the fungus while the later provides inorganic phosphates and nitrogenous compounds.

b) Rhizobia bacteria that fix nitrogen for leguminous plants (family fabaceae) in return for energy containing carbohydrates [52].

Service-Resource relationship: These are also common. Examples include:

a) The Oxypecker eats ticks on the zebra's skin. Whereas the bird gets food, the zebra gets service of pest control.

b) Pollination in which nectar or pollen (food resource are traded for pollen dispersal (service)

c) Ant protection of aphids where the aphid trade sugar-rich honey dew, a by-product of their mode of feeding on plant sap) in return for defense against predators such as ladybugs.

d) Phagophiles feed (resource) on ectoparasites thereby providing anti pest service as in cleansing symbiosis.

e) Elacatinus and Globiosoma, genus of globies also feed on ectoparasites of their client while cleaning them [53].

f) Zoochory- an example where animals disperse the seeds of plants. This is similar to pollination in that the plant produces food resources (eg fleshy fruits, over abundance of seeds) for animals that disperse the seeds (service).

Service-service relationship: Strict service-service relationships are very rare for reasons which are not clear [51]. Examples of service to service relationships include:

- 464 a) Relationship between sea anemones and anemone fish in the family
465 Pomacentridae. The anemone provides the fish with protection from
466 predators, while the fish defends the animal against butterfly fish which
467 eats anemones. However, it is believed that there is more to this
468 relationship than service-service mutualism. For instance waste ammonia
469 from the fish feed the symbiotic algae that are found in the anemones
470 tentacles [54-55]. Thus what appears as service- service relationship has a
471 service-resource component.
- 472 b) Relationship between some ants in the genus *Pseudomyrmex* and trees in
473 the genus *Acacia* such as the Whistling thorn and Bullhorn *Acacia*. The
474 ants nest inside the plants thorns thereby obtaining shelter whereas the
475 plant gets protection from attacks by herbivores, which they frequently
476 eat, thereby introducing service-service relationship) and competition
477 from other plants by trimming back vegetation that would shade the
478 *Acacia*. In addition, another service-resource component is obvious since
479 the ants regularly feed on lipid-rich food bodies called Beltian bodies that
480 are found on the *Acacia* plant
- 481 c) In the neotropics, the ant, *Myrmelachista Schumani* builds its nest in
482 special cavities in *Duroia hirsute*. Plants in the vicinity that belong to
483 other species are killed with formic acid. This selective gardening can be
484 so aggressive that small areas of the rain forest are dominated by *Duroia*

485 hirsute. These peculiar perches are known by the local people as “devils
486 gardens” [56].

487 d) *Cordia* species trees in the Amazonian rain forest have a kind of
488 partnership with allomerus species ants, which make their nests in
489 modified leaves. The ants often destroy the trees flowerbud to make more
490 living space available. As the flowers die, more leaves develop and take
491 their place, thus creating more room for the ants.

492 e) Another type of allomerus species ants lives with the *Hirtella* sp tree in
493 the same forest; but unlike in the former relationship, when the tree wants
494 to make flowers, the leaves harbouring the ants dwellings begin to wither
495 and shrink, thus forcing the ants to flea thereby leaving the trees flowers
496 to flourish free from ants attack [56].

497 **3.4 HUMANS AND MUTUALISM**

498 Mutualistic relationships between humans and other species abound in life:

499 **a) Humans and gut flora:** The gut flora helps man to digest food efficiently
500 [57].

501 **b) Head lice and Man:** It is apparent that head lice confer some immunity
502 to man thereby helping to reduce the threat from body louse-borne lethal
503 diseases.

504 **c) Humans and domesticated animals:** Dogs and sheep were among the
505 first animals to be domesticated by man and they are beneficial to him.

d) Man and some agricultural varieties of maize: The later are unable to reproduce without human intervention. First the leafy sheath does not fall open, and secondly, the seed head (the “corn on the cob”) does not shatter to disperse the seeds naturally unless man intervenes

e) In traditional agriculture, some plants have mutualists as companion plants, providing each other with shelter, soil fertility and or natural pest control. For example, beans may grow up corn stalks as trellis, while fixing nitrogen in the soil for the corn. This phenomenon is applied in the Three Sisters farming [59].

f) The Boran people of Ethiopia and Kenya traditionally use a whistle to call the honey guide bird. If the later is hungry, it usually guides them to a bee’s nest where they (Boran) harvest the honey leaving some for the birds to eat [60].

g) In Laguna Brazil, a population of bottle nose dolphins communicates through body language with local net using fishermen in order for both to catch schools of mullet [61].

526

527

528

529 SECTION FOUR

530 **4.1 COMMENSALISM AND PHORESY**

531 Commensalism simply means “eating at the same table”. It is a type of
532 symbiotic relationship where one partner benefits whereas the second partner
533 (the host) is neither helped nor harmed. Commensal relationships mainly
534 involve feeding on food “wasted” or otherwise not consumed by the host.

535 Examples of commensalism include:

536 a) **Remora sharks and Whales:** The remora sharks have adhesive disk on
537 the dorsal surface of their head which they use to attach to larger animals
538 such as whales which tend to be sloppy eaters. When food floats away
539 from the whale’s mouth, the remora shark can unhitch itself and collect
540 the scraps of food from the host.

541 b) **Barnacles and Whales:** Barnacles are crustaceans whose adults are
542 sedentary. The motile larvae find a suitable surface and then undergo
543 metamorphosis to the sedentary form. The barnacles adhere to the skin
544 of a whale or shell of a mollusk and are transported to areas with new
545 sources of food.

c) **The titan triggerfish (*Balistoides viridescens*) and smaller fish:** The former fish creates feeding opportunities for smaller by moving large rocks which are too big for the smaller fish to shift.

d) **Humans and protistans:** Humans harbour several species of commensal protistans such as *Entamoeba gingivalis* which lives in the mouth where it feeds on bacteria, food particles and dead epithelial cells but never harm healthy tissues. Adult tape worms though generally regarded as parasites may not have known ill effects on their hosts [62].

4.2 TYPES OF COMMENSALISM

Facultative commensalism: This is a situation where the commensal may not necessarily participate in the relationship to live eg stalked ciliates of the genus *verticella* are frequently found on small crustaceans but they can survive equally on sticks on the same pond.

Obligate commensalism: This is a situation where the commensals necessarily need each other to survive eg some related ciliates such as *Epistylis* spp cannot survive without the presence of other organisms especially crustaceans.

4.3 PHORESIS

This is the relationship in which two organisms are simply “travelling together” and there is no physiological or biochemical dependence on the

part of each participant. The two organisms are known as phoronts. Usually the smaller organism is usually carried by the larger organism (the host).

Examples of phoresic relationship include:

- a) Bacteria on the hairs of a fly
- b) Fungous spores on the feet of beetle
- c) Mites on insects such as beetles, flies or bees.
- d) Pseudo scorpions on mammals [63]
- e) Millipedes on birds [64].
- f) The *Dermatollia hominis* larvae usually live beneath the skin of warm blooded animals including man. The eggs are usually carried by other insects such as mosquitoes and are deposited on the host's skin as the mosquito perches to feed. The eggs quickly hatch and the larvae burrow their way into the skin.

Like commensalism, phoresis can be facultative or obligate depending on the existing environmental conditions.

4.4 OTHER RELATIONSHIPS

INQUILINISM

This is a type of relationship where one organism uses the other as a permanent housing or place of abode. Examples include:

- a) Epiphytic plants (eg Orchids) that grow on trees [65].

586 b) Birds that live in holes in trees.

587

588

589 **METABIOSIS**

590 This is a relationship in which one organism creates or prepares a suitable
591 environment for the other. Examples include:

592 a) Maggots which feast and develop in corpses

593 b) Hermit crabs which use gastropod shells to protect their bodies

594 **AMENSALISM**

595 This is the type of relationship that exists where one species is inhibited or
596 completely obliterated and the other is not affected. This type of relationship
597 is common in the natural world. An example is a sapling growing under the
598 shadow of a mature tree. The mature tree usually robs the sapling of
599 necessary sunlight and other nutrient (eg rain water). The mature tree
600 remains unaffected while the sapling dwindles and dies. The mature tree will
601 even make use of nutrients arising from the decaying sapling.

602 **SYNNECROSIS**

603 This is a rare type of symbiosis in which the interaction between species is
604 detrimental to both organisms involved. [9]. It is a temporal condition since

the interaction will eventually lead to death of the two partners.

Consequently, evolution selects against synnecrosis hence it is uncommon in

life and the term is rarely used. [66].

CONCLUSION

Host parasite relationships occur as a result of prolonged evolutionary

associations between organisms ie organisms developing or living with each

other in the same environment for a long time. The extent of association

determines the type of relationship which may result.

623

624

625

626 REFERENCES

627 1. Larry S,R and Gerald D.S . Foundations of Parasitology (6th Edition).

628 2000.pp 4-7.

629 2. The New Oxford Advanced English Dictionary.

630 3. Liddel and Scott. Greek English Lexicon.

631 4. Wilkinson, (2001). Symbiosis.Earth and Environmental Science. Earth

632 Science series

633 5. Douglas,A. Symbiotic Interactions, Oxford (Oxfordshire): Oxford

634 University press 1994, p 1

635 6. Douglas A.E. The Symbiotic Habitat, New Jersey; Princeton University

636 Press 2010 pp 5-12

637 7. Martin, B.D; Schwab,E. “Symbiosis: Living together in chaos”. Studies

638 in the History of Biology 2012, 4(4):7-25

639 8. Ahmadjian and Paracer (2000). The Role of Symbiosis in the Evolution

640 of biosphere. p12

- 641 9. Dorland (2007) “Symbiosis” Dorlands illustrated Medical Dictionary.
642 Philadelphia Elsevier Health Sciences, . Credo reference web. 17
643 Sept 2012
- 644 10. Poulin R (1992) Evolutionary ecology of parasites. From individual
645 to communities. Chapman & Hall, London
- 646 11. Tripet F, Jacot A, Richner H (2002a) Larval competition affects the
647 life histories and dispersal behavior of an avian ectoparasite.
648 Ecology 83:935–945
- 649 12. Tripet F, Richner H (1997) The coevolutionary potential of a
650 ‘generalist’ parasite, the hen flea *Ceratophyllus gallinae*.
651 Parasitology 115:419–427
652
653
- 654 13. Soler JJ, Møller AP, Soler M (1999) A comparative study of host
655 selection in the European cuckoo *Cuculus canorus*. Oecologia
656 118:265–276
- 657 14. Johnson KP, Williams BL, Drown DM, Adams RJ, Clayton DH
658 (2002) The population genetics of host specificity: genetic
659 differentiation in dove lice (Insecta: Phthiraptera). Mol Ecol
660 11:25–38
- 661 15. Combes C (1991) Evolution of parasite life cycles. In: Toft CA,
662 Aeschlimann A, Bolis L (eds) Parasite-host associations.
663 Coexistence or conflict? Oxford University Press, London, pp
664 62–82
- 665 16. Thompson JN (1994) The coevolutionary process. University of
666 Chicago Press, Chicago
- 667 17. Kawecki TJ (1998) Red queen meets Santa Rosalia—arms races and
668 the evolution of host specialization in organisms with parasitic
669 lifestyles. Am Nat 152:635–651
- 670 18. De Meus et al, 1998 (Delete)
- 671 19. Jaenike J (1990) Host specialization in phytophagous insects. Annu
672 Rev Ecol Syst 21:243–273
- 673 20. Combes C (1995) Interactions durables: ecologie et évolution du
674 parasitisme. Masson, Paris
- 675 21. Combes C (1997) Fitness of parasites—pathology and selection
676 [Review]. Int J Parasitol 27:1–10

- 677 22. Norton DA, De Lange PJ (1999) Host specificity in parasitic
678 mistletoes (Loranthaceae) in New Zealand. *Funct Ecol* 13:552–
679 559
- 680 23. McCoy KD, Boulinier T, Tirard C, Michalakakis Y (2001) Host
681 specificity of a generalist parasite: genetic evidence of
682 sympatric host races in the seabird tick *Ixodes uriae*. *J Evol*
683 *Biol* 14:395–405
- 684 24. Giorgi MS, Arlettaz R, Christe P, Vogel P (2001) The energetic
685 grooming costs imposed by a parasitic mite (*Spinturnix myoti*)
686 upon its bat host (*Myotis myotis*). *Proc R Soc Lond B Biol Sci*
687 268:2071–2075
- 688 25. Norton DA, Carpenter MA (1998) Mistletoes as parasites: host
689 specificity and speciation. *Trends Ecol Evol* 13:101–105
- 690 26. Timms R, Read AF (1999) What makes a specialist special? *Trends*
691 *Ecol Evol* 14:333–334
- 692 27. Tompkins DM, Clayton DH (1999) Host resources govern the
693 specificity of swiftlet lice: size matters. *J Anim Ecol* 68:489–
694 500
- 695 28. Combes, (2005). *The Art of being a parasite*. University of Chicago
696 press.www.press.uchicago.edu
- 697 29. Henry and Scott. *Greek English Lexicon*
- 698 30. Getz, W (2011). Biomass transformation webs provide a unified approach
699 to consumer-resource modeling. *Ecology Letters*, doi:10.1111/j.1461-
700 0248.2010.01566.x
- 701 31. Gardiner C.H, Koh D.S, Cardella T.A (1980). *Micronema* in man: Third
702 fatal infection. *Am J Trop Med Hyg* 30:586-589
- 703 32. Thomas J, Karsten S, Simona B, Francesca B, and Emilio B (2010).
704 “Corruption of ant acoustical signals by mimetic social parasites”
705 *Communicative and integrative Biology* 3(2):169-171

- 706 33. Parasitism: Bullies of Wild Life, the Bird World Wild life Magazine,
707 1997.
- 708 34. O Brien T.G, (1998).”Parasitic nursing behavior in the wedge-capped
709 capuchin monkey (*Cebus olivaceus*). *Am J Primatology* 16(4):341-344
- 710 35. Charles H, Godfray J (2004). “Parasitoids”. *Current Biology Magazine*
711 14(12): R468
- 712 36. Bug Life, 2013“The differences between Parasites and Parasitoids”. Bug
713 Life. Retrieved 2010- 07-19
- 714 37. Larry G, and Mark Wheelis (1991). *The Cartoon guide to Genetics*.
715 Harper Collins, 1991
- 716 38. Overview of Plants Diseases, 2013
- 717 39. Milan N,F, Cacao B.Z, Schlenke T.A (2012).” Alcohol Consumption as
718 self medication against Blood- Bourne parasites in the Fruit fly”. *Current*
719 *Biology* 22(6):488-93
- 720 40. Price W.A (1980), *Evolutionary Biology of Parasites*. Princeton
721 University Press, Princeton.
- 722 41. Wolff E.D.S, Steven W.S, John R.H, David J.V (2009). “Common Avian
723 Infection Plagued the Tyrant Dinosaurs”. In Hansen, Denis Marinus.
724 *PLoS ONE* 4 (9):e7288.

42. Rook G.A.W.(2007). “The Hygiene hypothesis and the increasing prevalence of Chronic Inflammatory disorders”. *Trans Roy Soc Trop Med Hyg* 101(11): 1072-4
43. Claude Combes (2005). *The Art of being a parasite*, Univ of Chicago Press, 2005
44. Lively M.M, and Dybdahyl M.F (2000). ‘ Parasite Adaptation to locally common Host Genotypes”. *Nature*, Vol 405. 8 June 2000.
45. Lafferty K.D, and Morris A.K (1996). Altered behavior of parasitized killifish increases susceptibility to predation by bird final hosts” *Ecology* 77.
46. Berdoy M, Webster J.P, Macdonald D.W (2000). Fatal attraction in rats infected with *Toxoplasma gondii*”. *Proc Biol. Sci* 267 (1452): 1591-4
47. Holts R.D.(2010). Parasitism. *Wikipedia, the free encyclopedia*
48. Thomson J.N, (2005). *The Geographic mosaic of coevolution*. Chicago, IL:University of Chicago Press.
49. Brostein J.L, (1994). Our Current Understanding of mutualism. *Quarterly Review of Biology* 69(1):31-51
50. Begon M, Harper J.I, and Tronsend C.R (1996). *Ecology, Individuals, populations and communities*, (3rd ed). Blackwell Science Ltd, Cambridge Massachussetts, USA.
51. Ollerton J (2006). “Biological barter”. *Patterns of Specialization compared across different mutualisms*. Pp 411-435 in: Waser, N.M and

- 747 Ollerton, J (Eds)-Plant Pollinator Interactions: From Specialization to
748 generalization. University of Chicago Press.
- 749 52. Denis R.F and Kiers E.T (2004). Why are most rhizobia beneficial to
750 their plant host, rather than parasitic? *Microbes and infection* 6(13):
751 1235-1239.
- 752 53. Soares M.C, Cote I.M, Cardoso S.C and Bshary R (2008) “The cleaning
753 goby mutualism: a system without punishment, partner Switching or
754 tactile stimulation”. *J Zool* 276 (3):306-312
- 755 54. Porat D and Chadwick-Furman N.E (2004). Effects of anemone fish on
756 giant sea anemones: expansion behavior, growth, and survival.
757 *Hydrobiologia* 530:513-520.
- 758
- 759 55.Porat D and Chadwick-Furman N.E (2005). Effects of anemone fish on
760 giant sea anemones: ammonium uptake, zooxanthella content and tissue
761 regeneration. *Mar Freshw Behav- Phy*, 38,, 43-51.
- 762 56.Piper R (2007). Extra ordinary Animalas: An Encyclopedia of Curious
763 and unusual Animals, Greenwood Press.
- 764 57. Sears C.L (2005). “A dynamic partnership: celebrating our gut flora”.
765 *Anaerobe* 11(5): 247-51
- 766 58. Lozsa and Apari, 2012(Delete)
- 767 59.Mt. Pleasant J (2006). “ The Science behind the three sisters mound
768 system: An agronomic assessment of an indigenous agricultural system in

- 769 the North- east. In: John E Staller, Robert S Tykot and Bruce F BENZ.
 770 Histories of maize: multidisciplinary approach to the prehistory,
 771 linguistics, biogeography, domestication and evolution of maize,
 772 Amsterdam, pp529- 537
- 773 60. Gibbon J Whitefield, forwarded by Odum Eugene P (2010). Keeping all
 774 the pieces: Perspectives on Natural History and the environment. Athens,
 775 Georgia. University of Georgia Press, pp41-42.
- 776 61. [Http://newsdiscovery.com/animals/whales-dolphin](http://newsdiscovery.com/animals/whales-dolphin)
- 777 62. Insler G.D, and Robberts L.S (1976). Hymenolepsis diminuta: Lack of
 778 pathogenecity in the healthy rat host. Exp Parasitology 39:351-357
- 779 63.Durden L.A (2001). “ Pseudoscorpions associated with Mammals in
 780 Papua New Guinea” Biotropica 23(2):204-206
- 781 64.Tajovy Karel et al (2001). “Millipeds (Diplopoda) in Dogs Nest”. Eur J
 782 Soil Biol Vol 37:321-323
- 783 65. Hogan C.M (2011). Commensalism. Topic Ed, M, Mcginley, Ed-in-chief
 784 C.J Cleveland. Encyclopedia of Earth. National Council for Science and
 785 the Environment. Washington D.C.
- 786 66. Lidicher W.Z (1979). A Clarification of Interaction in Ecological
 787 Systems”. BioScience 29:475-477.
- 788
- 789