

Cypermethrin-induces oxidative stress to the freshwater ciliate model: *Paramecium tetraurelia*.

ABSTRACT

Synthetic Pyrethroids are considered to be safe over other insecticides; however, data indicate that their use may pose risk to environmental biota, especially, aquatic organisms. Therefore, the aim of the present study was to investigate the effect of cypermethrin, a widely used insecticide and one of the most common contaminants in freshwater aquatic system on the oxidative stress biomarkers of the freshwater ciliate *Paramecium tetraurelia*.

After the treatment of paramecium cells with increasing concentrations of cypermethrin (0.05, 0.5, 1, 2 µg/l), we followed up the growth kinetics, generation time and the response percentage. Also, we studied the variation in biomarkers of stress such as: proteins, GSH content, GST and CAT activities.

Our results showed a significant decrease in the proliferation of cell, we denote a difference of nearly 1,900 cells between the control cells and those treated with (2µg/l) at the fourth day of treatment. This finding is correlated by the decrease in generation number and velocity and the increase in generation time. Also, we noted an inhibition in the response percentage: it varies from (20%) to (54%) for 0.5µg/l and 2µg/l respectively.

The rate of total proteins increased in dose dependent manner and very highly significant for the two highest concentrations (1 and 2 µg/l).

The monitoring of biomarkers revealed a depletion in GSH content in a proportional and dose dependent manner (it is 7.34188758µmol/mg Pro for the control whereas it is 2.41682134 µmol/mg Pro for 2µg/l) accompanied by an increase in the GST activity (we note an increase of the order of 1.62932472 µM/min/mg Pro for the highest concentration compared to the control which is of the order of 0.59883133 µM/min/mg Pro) .In parallel, a strong induction of the CAT activity was noted specially for the highest dose.

Keywords: Cypermethrin, *Paramecium tetraurelia*, growth, oxidative stress, GST, GSH, CAT.

1. INTRODUCTION

The long-term ecological hazard associated with the use of organochlorine, organophosphate and carbamate compounds propelled the introduction of new generation of pesticides with a lesser degree of persistence. As a consequence, the use of pyrethroids as insecticidal and anti-parasitic formulations has markedly increased as a viable substitute and currently accounts for over 30 % of insecticides used globally [1]

Indeed, synthetic pyrethroids are synthetic chemical analogs and derivatives of pyrethrins, they represent the third largest class of chemical insecticides after organophosphates and chloronicotinyl insecticides [2]. The pyrethroids have been divided into two types (type I and type II) on the basis of their chemical structure and toxic manifestation. In fact, type I pyrethroids are those which lack α-cyano moiety and give rise to the tremor syndrome (T syndrome) while type II pyrethroids are those which contain α cyano moiety and cause choreoathetosis/ salivation (CS)syndrome [3]. Their general site of action is biological membranes by alteration of sodium transport but they also affect chloride and calcium channels [4].

Several studies have indicated that this class of insecticides is highly toxic to a number of non-target organisms such as: bees, fish and aquatic invertebrates [5][6] [7][8] [9] [10] [11] [12] [13] [14]. For instance, these pesticides have been found to induce alterations in the hematological profiles of *Channa punctatus* and *Prochilodus lineatus* [15] [16]), reproduction and physiology of *Cyprinus carpio* [17] and *Atlantic salmon* [18]. Furthermore, [19] demonstrated that type II pyrethroids could increase SOD activity in zebrafish larvae after 8h exposure, which suggest that oxidative stress could be induced and played an important role in developmental toxicity in fish.

Cypermethrin (CYP), the alpha-cyano-3-phenoxybenzyl ester of 2,2-dimethyl-3-(2,2-dichlorovinyl)-cyclopropane-carboxylic acid is the most widely used type II pyrethroid insecticide, it is commonly used in urban and agricultural environments[20].

Cypermethrin is very highly toxic to fish and aquatic invertebrates. Many studies reported that this compound is metabolized and eliminated significantly more slowly by aquatic organisms than by mammals or birds [21].

The environmental contaminants affect aquatic ecosystems by inducing oxidative damage as a sensitive and specific biomarker and causing cell death, via the enhancement of intracellular reactive oxygen species (ROS) and perturbation of antioxidant efficiency [22].

Cypermethrin produces drastic effects on both invertebrates[23] and vertebrates [24]. [25] reported its potential to induce hepatic oxidative stress, DNA damage and apoptosis in adult zebrafish *Danio rerio*. As well, [20], indicated the behavioral morphological deformities and the induction of biomarkers of oxidative damage due to sublethal concentration of cypermethrin on tadpoles of *D.melanostictus*. Also, experiments conducted with *Ceriodaphnia dubia* showed the increase of the toxic effect with increasing concentrations and exposure time [26]. Moreover, *Brachionus calyciflorus* and *Thamnocephalus platyurus* have, also, demonstrated high sensitivity to permethrin, resmethrin and cypermethrin[27].

The use of *Paramecium* species as a model of survey has been reported by several authors in some disciplines; in genetic, because its sequencing genome is well known, researchers used *Paramecium tetraurelia* for genetic analyses, gene expression and mutation[28] [29]. In physiology, paramecia is used in general for studying the role, the function and the cell organization [30]. In ecotoxicology: *Paramecium* species were used to study environmental qualities and toxic effects of industrial, domestic and agricultural chemicals[31] [32] [33] [34] [35] [36] [37] [38]. Further, the unicellular ciliate facilitates the study of physiological process and cytotoxicity of pollutants, that's why, they are well suited to being included in the increasing panel of organismic systems that could meet the 3Rs (aimed at Reducing, Refining and Replacing tests on vertebrate organisms in toxicological studies) and sensitive to such environmental compounds. This sensitivity is due to their simple eukaryotic single-cell and organism organization which exposes their receptors to external environment, making them respond to environmental stimuli [39]. Moreover, their easy culture and maintenance and their short cell-cycle provide results in a short time. For all these reasons, ciliates, especially *Paramecium* species, have been exploited as excellent tools for environmental biomonitoring, either as bioindicators of pollution or bioassays to evaluate the effect of toxic compounds [40] [41] [42] [43].

Thus, the present work was carried to investigate the cytotoxicity of cypermethrin at different sub lethal concentrations on population growth and some biomarkers of oxidative stress of the ciliated protozoan *Paramecium tetraurelia*.

2. MATERIAL AND METHODS

2.1. Test organisms:

The biological model used in our study is a unicellular microorganism *Paramecium tetraurelia*.

2.2. Test chemical:

The insecticide used for our experiments is cypermethrin (Figure 1) that belongs to the chemical family of pyrethroids type II.




Figure 1. Chemical structure of Cypermethrin

2.3. Treatment:

The habitual culture of *Paramecium tetraurelia* was done in the culture medium described by [35] at Ph 6.5 and 28±2°C. Cells were transplanted each three days for keeping the youthful state of culture [35] [36] [38].

Paramecium tetraurelia were incubated with the tested insecticide concentrations in aliquots of 10 ml, the retained concentrations were 0.05, 0.5, 1 and 2 µg / l.

Two modes of treatment have been adopted: For growth kinetics, the insecticide treatment was performed before the transplantation of *Paramecium* cells (at t=0). For the enzymatic assays, the treatment was carried at the end of the exponential growth phase (t=96 H) [44].

2.4. Parameter measurement:

2.4.1. Growth kinetic:

For growth experiments, the culture was done at 28°C in test tube using 10 ml of culture medium. For each tube we added 13 cells of *paramecia*. The growth kinetic study was realized by the daily cell counting after fixation with lugol under optic microscope type LEICA DM 1000.

Based on the data, the number, the time and the velocity of generation were calculated by the following formula:

$$n = (\log N_t - \log N_0) / \log 2$$

$$k = n/t$$

$$g = 1/k$$

Where n is the number of generation, N_t is the population in time t, N₀ is the initial number of cells, k is velocity of generation and g is the generation time.

2.4.2. Response percentage:

The response percentage was calculated to evaluate the toxicity of xenobiotics via the inhibition of cell growth after 96 H of exposure.

Positive values indicate an inhibition of growth while negative values indicate a stimulation of growth[45].

The assessment of the response percentage is calculated according to the following formula:

$$\text{Response (\%)} = \frac{(N_c - N_e)}{N_c} \times 100$$

Where N_c is the number control cells, N_e is the number of treated cells.

2.4.3. Total protein estimation:

Total protein is determined by the method of [46]. It is a colorimetric method using BSA as standard. The absorbance is measured at a wavelength of 595 nm using spectrophotometer type Jenway 3600.

2.4.4. Estimation of glutathione (GSH) content:

GSH content was quantified using the method of [47]. Cells are mixed in 1 ml EDTA (0,02M). 0,2 ml of ASS was added to 0,8 ml of homogenate. After agitation, the homogenate was centrifuged. The assay mixture contains 1 ml tris/EDTA buffer (0.02 M, pH 9,6), 0.025 ml of 5,5'-dithiobis-2-nitrobenzoic acid (DTNB) and the *paramecium* sample. The reaction was monitored at 412 nm and the amount of GSH was expressed as µmol/mg of proteins.

2.4.5. Determination of glutathione S-transferase (GST) activity:

The GST activity was measured according to the method of [48]. the homogenization of samples was done in 1 ml of phosphate buffer (0,1 M, pH 6) and centrifuged (14000 rpm/30 min). The final reaction contain 1,2 ml CDNB (1 mM) / GSH

(5mM) and the sample. The absorbance was measured spectrophotometrically at 340 nm. The result was expressed as $\mu\text{mol/ min/ mg}$ of proteins.

2.4.6. Determination of catalase (CAT) activity:

The CAT activity was determined spectrophotometrically at 240 nm by calculating the rate of degradation of H_2O_2 [49]. Samples are mixed in 1 ml of phosphate buffer then centrifuged at 15000 g. At 0,025 ml of supernatant we added 0,75 ml of phosphate buffer and H_2O_2 . The result was expressed as $\mu\text{mol/ min/ mg}$ of proteins.

2.5. Statistical analysis:

The obtained results are represented by the average \pm Standard Error. Statistical analysis of data is performed using Minitab student t-test.

3. RESULTS

3.1. Effect of Cypermethrin on growth kinetic:

The growth kinetic provides information about the toxic effect of a specific substance. Figure (2) represents the effect of cypermethrin on the variation of paramecium cells number (control and treated) versus time.




Figure 2. Effect of Cypermethrin on *Paramecium tetraurelia* growth (T 28°C pH: 6.5)

Different chosen concentrations inhibited the population growth in a dose-dependent manner especially for the highest concentration that inhibits strongly after 4 days of treatment. Indeed, we denote a difference of nearly 1,900 cells between the control cells and those treated with this concentration (2 $\mu\text{g/l}$) at the fourth day of treatment.

The proliferation of *Paramecium tetraurelia* was significantly affected by the action of cypermethrin as the generation number and concentration of cypermethrin are inversely proportional (table1). The generation time gradually increased with the increase of Cypermethrin concentrations.

Table 1. Effect of cypermethrin on *Paramecium tetraurelia* generation number (n), and generation time (g) at 96 H:

Cypermethrin concentrations (µg/l)	Generation number (n)±SE	Generation time (g)±SE
Control	8.07±0.10	11.90±0.15
0.05	7.75±0.04	12.40±0.06
0.5	7.70±0.04	12.47±0.07
1	7.54±0.16	12.74±0.28
2	6.90±0.20	13.85±0.41

Each value is mean of four assays ± Stander Error

Generation velocity (n/H)




Figure 3. Effect of cypermethrin on the velocity of generation of *Paramecium tetraurelia* (t=96 H)

The decrease in the velocity of generation compared to the control shows the negative response of *Paramecium tetraurelia* to the increasing concentrations of cypermethrin. Result revealed that insecticide has slowed the generation velocity of exposed paramecia in a dose dependent manner.

3.2. Response percentage:

The results obtained concerning the response percentage confirm those of kinetics growth.




Figure4. Effect of Cypermethrin on the response percentage at 96 H (n= 4)

Response percentage

We denote that the inhibitory effect was dose-dependent and proportional to the increasing concentrations.

Thus, the response percentage was positive and show a strong inhibition of microorganisms growth .In fact, it varies from (20%) to (54%) for 0.5µg/l and 2µg/l in which more than half of population is inhibited.

3.3. Protein estimation:

We note a dose dependent increase of total protein rate in the presence of xenobiotic. The statistical analysis indicates very highly significant differences ($P<0.001$) for the two highest concentrations (1 and 2 µg/l).

Table 2. Effect of cypermethrin on the rate of total proteins in *Paramecium tetraurelia* versus time

Cypermethrin concentrations µg/l	Rate of total proteins µM/mg of tissues ±SE
Control	0.283±0.011
0.05	0.294±0.009
0.5	0.339±0.009
1	0.620±0.022 ***p
2	0.772±0.019 ***p

*** $P=0.001$

3.4. Estimation of glutathione (GSH) content:

Figure (5) illustrates the variations of total GSH content in *Paramecium tetraurelia* exposed to increasing concentrations of cypermethrin.




Figure 5. Variations of GSH content in *Paramecium tetraurelia* exposed to increasing concentrations of cypermethrin



The result shows that this nonenzymatic antioxidant tends to decrease in dose dependent manner. Thus, the GSH rate is (7.34188758µmol/mgPro) for the control whereas it is (2.41682134 µmol/mgPro) for paramecia treated with the highest concentration (2µg/l) ie three times less. The statistical analysis reveals a significant difference ($P<0.050$) for the highest concentration compared to the control.

3.5. Determination of glutathione S-transferase (GST) activity:




Figure 6. Variations of GST activity in *Parametium tetraurelia* exposed to increasing concentrations of cypermethrin.



The results concerning the variation rate of GST show a strong induction of the activity of this enzyme particularly for the highest concentration. Indeed, this induction is in dose-dependent manner: we note an increase of the order of (1.62932472 μM/min/mg Pro) for the highest concentration compared to the control which is of the order of (0.59883133 μM/min/mg Pro), that is to say, three times higher.

The statistical study show a significant difference ($P < 0.050$) for the highest dose compared to the control.

3.6. Determination of catalase (CAT) activity:





Figure 7. Variations of CAT activity in *Parametium tetraurelia* exposed to increasing concentrations of cypermethrin.



The effect of cypermethrin on CAT activity is illustrated in figure (7). The results show a significant increase for the second concentration (0.5µg/l) and a very highly significant increase ($P < 0.001$) of the activity of CAT in cells treated with the highest concentration (2µg/l) compared with the control.

4. Discussion

The problem of environmental contamination by the excessive use of pesticides cannot be neglected [50]. Extensive application is usually accompanied with serious problems and health hazard. It is established that many chemicals, in common use, can produce some toxic effects on biological systems when tested on various type of experimental models through their mode of action or by production of free radicals that damage all cell compounds [51]. In fact, these chemicals act as pro-oxidants [52] [53] [54] [55] [56] [57] [58].

Oxidative stress is believed to occur when there is an imbalance in the biological oxidant-to-antioxidant ratio; this can result in oxidative damage to lipid, proteins, carbohydrates and nucleic acids. In most cases, the abnormal generation of ROS, which can result in significant damage to cell structure, is considered an important signal of oxidative damage [59].

Pyrethroids group of pesticides is the most commonly used in agriculture today [2]. However, it has been reported by several investigations conducted in various animal species that this pesticides cause oxidative damage [60] [61] [62] through the generation of ROS and can alter the antioxidants or free oxygen radical scavenging enzymes systems in animals, especially, in aquatic organisms [63].

Cypermethrin, is probably the most used pyrethroid. Studies showed that the excessive use can adversely affect most physiological processes [64].

Protists are eukaryotic unicellular organisms and their position in the food web makes them excellent models for predicting the effects of chemicals on aquatic communities. Ciliated protozoa represent a basic component of aquatic environment, where they play critical roles both quantitatively and qualitatively [65].

In this context, the ciliate assay has become a valuable tool for detection of environmental disturbance and for assessment of the trophic state [65] [66].

Paramecium is one of the most commonly used ciliated for laboratory research to investigate the direct toxicity of compounds [32] [35] [67] [68] [69].

That is why we chose *Paramecium tetraurelia* as a biological model for elucidating cypermethrin toxicity.

In this study, we were interested in the first time at the effect of cypermethrin on population growth. Our result showed an inhibition in the growth of microorganisms especially for the highest concentrations. Similar results were reported in studies [35] [36] [70] that investigate the effect of different chemicals on the physiology and morphology of *Paramecium* sp.. [31] Reports that toxics may affect the survival of protozoa in a variety of ways, as the concentration of toxicants in the cell membrane increase and destroy their integrity causing cell death. Toxic affects freshwater ciliates; these effects are perceptible at the population level by reducing the number of cells and on the cellular level by a structural behavioral and physiological damage.

The findings concerning growth kinetics were confirmed, on the one hand, by the decrease in generation number, the decrease in velocity as well the increase in generation time which mean that the proliferation and cellular metabolism were affected [33] [71] [72] [73] [74]. On the other hand, the positive value of response percentage demonstrated the inhibitory effect of cypermethrin. Indeed, cypermethrin as a lipophilic compound can penetrate into cell, disturbing phospholipid orientation and causing changes in fluidity of membrane [75].

Proteins are one of the major energy reserves present in all organisms, these reserves will be affected by toxicant exposure [20]. In this work, we noted an increase of total protein rate in a dose dependent manner and very highly significant for 1µg/l and 2µg/l. This finding is in agreement with those of [76] [77] who showed an increase in the rate of total proteins of paramecia treated with Bifenazole and Proclaim.

Our hypothesis is that this increase could be related to the induction of the detoxification process elaborated by this control system which is composed of enzymes, proteins and antioxidant molecules [78].

The antioxidant defense systems are present in all aerobic cells and neutralize the intermediate chemical reactions produced endogenously and/or metabolism of xenobiotics. The antioxidant system activity may undergo an increase or depletion under the effect of a chemical stress [79]. the cells are equipped with both the enzymatic and nonenzymatic

antioxidants for combating oxidative stress, which may be either due to increased production of free radical or impaired antioxidant defense or both [80].

Alteration of antioxidant enzymes by cypermethrin has also been reported to be one of the mechanisms of toxicity.

The Glutathione is the major non enzymatic radical scavenger in the animal cells; it is the most abundant thiol, which scavenges residual free radicals resulting from oxidative metabolism and escaping decomposition by the antioxidant enzymes [81]. During the metabolic action of GSH, its sulfhydryl group becomes oxidized resulting with the formation of the corresponding disulfide compound, GSSG (oxidized form) [2] [82]. In this work, we noted a significant depletion in a concentration and dose dependent manner. The decrease in total GSH level may be due to the presence of free radicals produced by the insecticide [2] [44] [77]. In addition, GSH also participates in the detoxification of xenobiotics as a substrate for the enzyme GST, so, it plays a crucial key role in cellular defense against pesticides toxicity [83].

The GST is enzyme of biotransformation that catalyzes the conjugation of electrophilic substrates to the thiol group of GSH, producing less toxic forms and also lipid peroxides [84] [85]. In our study, the increase in GST activity was in dose dependent manner and significant for the highest concentration (2µg/l). The induction of GST activity may be beneficial to handle a stress condition and indicates protection against cypermethrin. [20] Reported an increase in the GST activity by cypermethrin in tadpoles of *D. melanostictus*. [86] Suggested that increase in GST activity is involved in metabolic detoxification of butachlor in *Rhamdia quelen*. Also, [87] reported that this enzyme seems to be implicated in the detoxification of cypermethrin in amphibian larvae.

Catalase is the most important mechanisms against toxic effects of oxygen metabolism. It catalyzed the conversion of hydrogen peroxide into water. This antioxidant enzyme can, therefore, alleviate the toxic effect of ROS [85]. Present study clearly showed a dose-frequency-dependent increase in catalase activity in individuals treated by different concentrations probably due to the intensification of antioxidant activity in *Paramecium*. CAT is one of the most active enzymes and its level change first following induction of oxidative stress [80]. The present result is consistent with those of [53][54][72] who reported intensification in the Catalase activity in many animal models when treated by pesticides. These results indicate the activation of protective mechanisms necessary for the scavenging of the produced reactive oxygen radicals.

Cytochrome P450 monooxygenases heme-thiolate enzymes catalyzing various reactions, but are best known for their monooxygenase activity, inducing reactive or polar groups into xenobiotics or endogenous compounds [88]. Insect genomes revealed a large expansion of the P450 gene family. Elevated level of P450 activity has frequently been observed in pyrethroid resistant insects populations [89]. Several CYP genes were also linked to pyrethroid resistance [90] [91]. These findings validates many P450 as pyrethroid metabolizers [92] [93] [94] [95]. The living cell, *Paramecium tetraurelia*, contains a large number of gene families that are involved in processes associated with sensing and responding to environmental cues, such as: P450. Our hypothesis is that CYP450 enzymes are also implicated in the detoxification process of pyrethroids in *Paramecium* but this requires extensive studies to accurately determine this implication and its mechanisms.

Type II pyrethroids seems be toxic to *Paramecium tetraurelia*, an organism that does not poses a voltage sensitive sodium channel. Likewise, it is established that type II pyrethroids stimulated *Paramecium tetraurelia* back-swimming behavioral, an avoidance behavioral response that is controlled exclusively by Ca⁺⁺ uptake via voltage sensitive calcium channels associated with the cilia: [96] have characterized the action of pyrethroids on ciliary calcium channel in *Paramecium tetraurelia*. The study was conducted with deltamethrin, the results revealed that the toxic effect of deltamethrin is structurally related, dose dependent and enhanced by depolarization and provide substantial evidence that type II pyrethroids act as potent calcium channel agonists on the ciliary voltage sensitive channel of *Paramecium tetraurelia*. Furthermore, The effect could be due to the Ca⁺⁺ accumulation in the cell which leads to free radical mediated cell damage [97].

5. CONCLUSION

In summary, under the current experimental conditions, cypermethrin is toxic to the freshwater ciliate *Paramecium tetraurelia*. Exposure to low concentrations of cypermethrin showed significant adverse on growth accompanied with the induction of oxidative damage supported by the decrease in GSH content and the intensification of the antioxidant enzymes such as GST and CAT. It showed be mentioned that other biomarkers of oxidative stress and lipid peroxidation have to be measured, it is the same for the detection and estimation of ROS. A genotoxic study may provide more answers concerning the effects of cypermethrin on *Paramecium tetraurelia*.

REFERENCES

- [1] Prasanthi K, Muralidhara, Rajini P S. Fenvalerate-induced oxidative damage in rat tissues and its attenuation by dietary sesame oil. *Food Chemical Toxicology* . 2005; (43): 299-306.
- [2] El-Demerdash F M. Lipid peroxidation, oxidative stress and acetylcholinesterase in rat brain exposed to organophosphate and pyrethroid insecticides. *Food Chemical Toxicology*. 2011; (49): 1346-1352. doi: 10.1016/J.FCT.2011.03.018.
- [3] Ramesh C. Chlorinated hydrocarbons and pyrethrins/ Pyrethroids. *Reproductive and Developmental Toxicology*. 2011; (38):487-501.ISBN: 978-0-12-382032-7.
- [4] Steven B et al,. Binary mixture of pyrethroids produce differential effects on Ca^{++} influx and glutamate release at isolated presynaptic nerve terminals from rat brain. *Pesticide Biochemistry and physiology*. 2011; (99): 131-139.
- [5] Saha S, Kaviraj A. Acute toxicity of synthetic pyrethroids cypermethrin to some freshwater organisms. *Bull. Environmental Contaminant Toxicology*; 2008; (80): 49-52.
- [6] Ronco E A, Carriquiriborde P, Natale G S, Martin M L, Mugni H, Bonetto C. Integrated approach for the assessment of biotech soybean pesticides impact on low order stream ecosystems of the Pampasic Region, In: J.Chen. C. Guo(Eds). *Ecosystem Ecology Research Trends*. Nova Science. Hauppauge ; 2008:209-239.
- [7] Carriquiriborde P, Diaz J, Mugni H, Bonetto C, Ronco E A. Impact of cypermethrin on stream fish populations under field use in biotech-soybean production. *Chemosphere*. 2007; (68): 613-621.
- [8] Crossland N O. Aquatic toxicology of cypermethrin.II. Fate and biological effects in pond experiments. *Aquatic Toxicology* 1982; (2) 205-222.
- [9] Xiangguo S et al,. Developmental toxicity of cypermethrin in embryo-larva stages of zebrafish. *Chemosphere*. 2011;(85) :1010-1016. Doi: 10.1016/j.chemosphere.2011.07.024
- [10] Slaninova A, Smutna M, Modra H, Svobodova Z. A review; oxidative stress in fish induced by pesticides. *Neuro Endocrinol. Lett.* 2009; (30):2-12.
- [11] Zhang et al,. Time-dependent oxidative stress responses of crucian carp (*Carassius auratus*) to intermediated injection of extracted microcystins. *Bulletin of Environmental contamination and toxicology*. 2009; (82): 574-578.
- [12] Jin et al,. Developmental toxicity of cartap on zebrafish embryos. *Aquatic toxicology*. 2009. (95): 339-346.
- [13] Rizwan et al,. In vivo cytogenetic and oxidative stress-inducing effects of cypermethrin in freshwater fish, *Channa punctata Bloch*. *Ecotoxicology and Environmental Safety*. 2011; (74): 150-156. doi: 10.1016/j.ecoenv.2010.08.036
- [14] Saxana K, Seth N. Toxic effects of cypermethrin on certain hematological aspects of freshwater fish, *Channa punctatus*, *Bulletin of Environmental Contamination and Toxicology*. 2002; (69): 364-369
- [15] Parma M.J, Loteste M, Campana M, Bacchetta C. Changes in hematological parameteres in *Prochilodus lineatus* (Pisces, Prochilodontidae) exposed to sublethal concentrations of cypermethrin. *Environmental Biology*. 2007 ; (28) : 147-149.
- [16] Aydin R, Koprucu K, Dorucu M, Koprucu S, Pala M. Acute toxicity of synthetic pyrethroid cypermethrin on the common carp (*Cyprinus carpio L*) embryos and larvae. *Aquacult. Int.* 2005; (13): 451-458.
- [17] Koprucu K, Aydin R. The toxic effects of pyrethroid deltamethrin on the common carp (*Cyprius carpioL*) embryos and larvae. *Pesticide Biochemistry and physiology*. 2004; (80): 47-53.
- [19] Gu A et al,. Exposure to fenvalerate causes brain impairment during zebrafish development. *Toxicological Letter*. 2010; (197): 188-192.
- [20] Muniswamy D, Shambanagouda Marigooudar R. PatilVineetkumar, Ramzsh H. Behavioral, morphological deformities and biomarkers of oxidative damage as indicators of sublethal cypermethrin intoxication on tadpoles of *D.*

melanostictus (Schneider, 1799). Pesticide Biochemistry and Physiology . 2012; (103)127-134.<http://dx.doi.org/10.1016/j.pestbp.2012.04.009>.

[21] Bradbury S M, Coats J R, Toxicokinetics and toxicodynamics of pyrethroid insecticides in fish. Environmental Toxicology Chemistry. 1989; (8) 373-380.

[22] Regoli F, Frenzilli G, Bacchetti R, Annarumma F, Scarcelli V, Fattarini, D. Time-course variation in oxyradical metabolism, DNA integrity and lysosomal stability in mussels, *Mytilus galloprovincialis* during a field translocation experiment. Aquatic Toxicology. 2004; (68): 167-178.

[23] Gowlan B T, Moffat C F, Stagg R M, Houlihan D F, Davies I M. Cypermethrin induces glutathione S-transferase activity in the shore, *Carcinus maenas*. Marine Environment . 2002; (54): 169-177.

[24] Das B K, Mukherjee S C. Toxicity of cypermethrin in *Labeo rohita* fingerlings: biochemical, enzymatic and hematological consequences. Comparative Biochemistry and Physiology. 2003; (C:134): 109-121.

[25] Yuanxiang J, Shansahn Z, Yue P, Linjun S, Weiping L, Zhengwei F. Cypermethrin has the potential to induce hepatic oxidative stress, DNA damage and apoptosis in adult zebrafish *Danio rerio*. Chemosphere. 2011; (82): 398-404. doi: 10.1016/j.chemosphere.2010.09.072.

[26] Shen M, Kumar A, Ding S, Groke S. Comparative study on the toxicity of pyrethroids, alpha-cypermethrin and deltamethrin to *Ceriodaphnia dubia*. Ecotoxicology and Environment Safety. 2011; 1-5. doi:10.1016/j.ecoenv.2011.07.018.

[27] Sanchez-Fortun S, Barahona MV. Comparative study on the environmental risk induced by several pyrethroids in estuarine and freshwater invertebrate organisms. Chemosphere. 2005;(59): 533-559.doi: 10.1016/j.chemosphere.2004.12.023.

[28] Haynes W J, Ling K Y, Preston R R, Saimi Y, King C. The cloning and molecular analysis of pawn-B in *Paramecium tetraurelia*. Genetics. 2000; (155): 1105- 1117.

[29] Vayssie L, Skouri F, Sperling L, Cohen J. Molecular genetics of regulated secretion in *Paramecium*. Biochimie. 2000 ;(82:4): 269-88.

[30] Hemmersbach R et al,. *Paramecium* a model system for studying cellular graviperception. Advances Space. 2001; 27(5): 893-898.

[31] Madoni P. The acute toxicity of Nockel to freshwater ciliates. Environmental Pollution . 2000; (109): 53-59.

[32] Miyoshi Net al,. Use of *Paramecium* species in bioassays for environmental risk management : Determination of IC50 values for water Pollutants. Journal Health Science. 2003; 49: (6): 429-435.

[33] Rouabhi R, Berrebbah H, Djebbar M R. Toxicity evaluation of flycycloxuron and diflubenzuron on the cellular model, *Paramecium* sp.. African Journal Biotechnology. 2006; 5(1): 45-048.

[34] Mortuza M G, Takahashi T, Kosaka T, Hosoya H. Effect of industrial sludge toxicity on the cell growth of green *Paramecium*, *Paramecium bursaria*. J cell. Anim. Biol. 2010;3 (4): 62-66.

[35] Azzouz Z, Berrebbah H, Djebbar M R. Optimazation of *Paramecium tetraurelia* growth kinetics and its sensitivity to combined effects of azoxystrobin and cyproconazole. African Journal of Microbiology Research. 2011; 5(20) : 3243-3250.

[36] Benbouzid H, Berrebbah H, Berredjem M, Djebbar MR. Toxic effects of phosphoramidate on *Paramecium* sp. With special emphasis on respiratory metabolism, growth and generation time. Toxicological and environmental chemistry. 2012; 94 (3):557-565.

[37] Bouaricha H, Brrebbah H, Grara N, Djebbar M R. Response of *Paramecium* sp. with respect to an insecticide (Proclaim) : growth, content of MDA, AChE activity and respiratory metabolism. Journal of Applied Sciences Research . 2012; 8(8):4172- 4180.

[38] Boulassel A, Djebbar M R, Rouabhi R, Berrebbah H. Physiological and biochemical changes observed in alternative cellular model :*Paramecium tetraurelia* treated with paracetamol. International Journal of Biosciences. 2013 ; 3(9) : 132-141.

- [39] Amaroli et al., Nitric oxide production inhibited by xenobiotic compounds in the protozoan *Paramecium primaurelia*. Ecological Indicators. 2010; (10):212-216 . doi: 10.1016/j.ecolind.2009.04.012
- [40] Matsubara E, Harad K, Inone K, Koizumi A. Effects of perfluorinated amphiphiles on backward swimming in *Paramecium caudatum*. Biochemical and Biophysical Research communication. 2006; (339): 554-561. Doi: 10.1016/j.bbrc.2005.11.048
- [41] Kozai N, Ohnuk T, Koka M, Satoli T, Kamiya T. Behavioral of *Paramecium sp.* In solutions containing Sr and Bp: Do *Paramecium sp.* Alter chemical forms of those metals?. Nuclear Instruments and Methods in Physics Research. 2011; (B269): 2391-2398). Doi:10.1016/j.nimb.2011.02.052
- [42] Hussain MM, Amanchi NR, Solanki VR, Bhagavathi M. Low cost bioassay test for assessing cytopathological and physiological responses of ciliate model *Paramecium caudatum* to carbofuran pesticide. Pesticide biochemistry and physiology. 2008; (90): 66-70. doi: 10.1016/j.pestbp.2007.07.006
- [43] Venkateswara Rao, Arepalli S K, Gunda V G, Bharat Kumar J. Assessment of cytoskeletal damage in *Paramecium caudatum*: An early warning system for apoptotic studies. Pesticide biochemistry and physiology. 2008; (91): 75-80
- [44] Azzouz Z. Etude des effets toxiques d'un fongicide (AmistarXtra) et d'un herbicide (Glyphosate) sur la biologie et le comportement de *Paramecium tetraurelia*. 2012. Ph.D. Thesis ,Badji Mokhtar University, Annaba. Algérie. French.
- [45] Wong C K, Cheung M H Y. Toxicological assessment of coastal sediments in Hong Kong using a flagellate *Dunaliella tertiolecta*. Environmental pollution. 1999;(105): 175-83.
- [46] Bradford M M A. Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. Analytical Biochemistry. 1976; (72): 248-254.
- [47] Wechbeker G, Cory. Ribonucleotide reductase activity and growth of glutathione depleted mouse leukemia 1210 cell in vitro. Cancer letters. 1988; (40): 257-264.
- [48] Habig W H, Pabst M J, Jakoby W B. Glutathione S-transferases : the first enzymatic step in mercapturic acid formation. Journal of Biological chemistry. 1974;(249): 7130-7139.
- [49] Regoli F, Principato G. Glutathione, glutathione dependant and antioxidant enzymes in mussel *Mytilus galloprovincialis* exposed to metal under field and laboratory conditions: implication for the biomarker. Aquatic Toxicology. 1995;(31): 143-164.
- [50] Palanivelu V, Vijayavel K, Ezhilarasibalasubramanian S, Balasubramanian M P. Impact of fertilizer (urea) on oxygen consumption and feeding energetics in the freshwater fish *Oreochromis mossambicus* .Environmental Toxicology and Pharmacology.2005; (19): 351-355.
- [51] Khan M S. Protective effect of black tea extract on the levels of lipid peroxidation and antioxidant enzymes in liver of mice with pesticide induced liver injury. Cell Biological Function. 2006; (24): 332-372.
- [52] Limon-Pacheco J, Gonshebat M E. The role of antioxidants and antioxidant related enzymes in protective responses to environmentally induced oxidative stress. Mutation. 2009; (674):137-147.
- [53] Zeriri I, Tadjine A, Grara N, Belaouchet N, Berrebbah H, Djebbar MR. Potential toxicity of an insecticide of the family of carbaamates on a bioindicator model of the pollution the earthworm *Octodrilus complanatus* (Oligochaeta, humbricidae). Annals of Biological Research. 2012; 3(11): 5367-5373.
- [54] Belhaouchet N, Djebbar M R, Meksem L, Grara N, Zeriri I, Berrebbah H. Evaluation of the biomarkers of the oxidative stress induced by a biopesticide: The Spinosad on an alternative model, *Helix aspersa*. Journal of Applied Sciences Research. 2012 ; 8(8) : 4199-4206.
- [55] Grara, Noudjoud. Evaluation de la toxicité de certains polluants industriels sur un animal bioaccumulateur (gastéropode *Helix aspersa*) : Cas des métaux. 2011. Ph.D .Thesis, Badji Mokhtar University, Annaba. Algérie. French.

- [56] Tlidjen S, Meksem L, Bouchelaghem S, Sbartaï H, Djébar, M R. Effect of the herbicide Calliofop 36EC on the growth, antioxidant enzyme response and respiratory metabolism in two aquatic plant *Elodea Canadensis* and *Lemna minor*. *Advances in Environmental Biology*. 2012; 6 (9): 2514-1530.
- [57] Khaldi F, Berrebbah H, Djébar M R. Study of atmospheric pollution emitted rated a plant of a fertilizers (Algeria) by the use of bioindicator plants: lichens. *Advances in Environmental Biology*. 2012; 6 (5): 1823- 1833.
- [58] Issaad G, Djébar M R, Grara N, Berrebbah H, Chagra A. Oxidative stress, chlorophyll content and ROS production and localization in *Triticum durum* Seed. *Annals of Biological Research*. 2013; 4(5):11-15.
- [59] Brazilia A, Yamamoto K I. DNA damage responses to oxidative stress. *DNA Repair*. 2004; (3):1109-1115.
- [60] El-Demerdash F M. Lambda-cyhalothrin- induced changes in oxidative stress biomarkers in rabbit erythrocytes and alleviation effect of some antioxidant. *Toxicology in Vitro*. 2007;(21): 392-397.
- [61] Giray B, Gurbay A, Hincal F. Cypermethrin- induced oxidative stress in rat brain and liver in prevented by vitamin E or allopurinol. *Toxicology letters*. 2001;(118):139-146.
- [62] Liu H, Zhao M, Zhang C, Ma Y, Liu W. Enantio selective cytotoxicity of the insecticide bifenthrin on a human aamion epithelial (FL). cell. Line. *Toxicology*. 2008; (253:1-3): 89-96.
- [63] Livinstone D R. Contaminant- stimulated reactive oxygen species production and oxidative damage in aquatic organisms. *Marine Pollution Bulletin*. 2001;(42): 656-666.
- [64] Azizullah A, Nasri A, Richter P, Lebert M, Hader D P. Evaluation of adverse effects of two commonly used fertilizers, DAP and urea, on mobility and orientation of the green flagellate *Euglina gracilis*, *Environmental and Experimental Botany*. 2011; (74):140-150.
- [65] Madoni P, Guiseppa Roeo M. Acute toxicity of heavy metals towards freshwater ciliated protests. *Environmental pollution*. 2006; (141): 1-7.
- [66] Cairns J, Pratt J R. The scientific basis of bioassays. *Hydrobiologia*. 1989;188(189):5-20.
- [67] Cernichiari E, Muscatine L, Smith D C. Maltose excretion by the symbiotic algae of *Hydra viridus*. *Proceeding of the Royal Society B: Biological Science*. 1969;(173): 557-576.
- [68] Benboouzid H, Berrebbah H, Berredjem M, Djébar M R. Effect of novel phosphoramidate on growth and respiratory metabolism of *Paramecium Aurelia*. *Journal of Natural Science Biology and Medicine*. 2012; 3(1): 48-51.
- [69] Rouabhi R, Berrebbah H, Djébar M R . Toxicity evaluation of flucycloxuron and diflubenzuron on the cellular model, *Paramecium sp*. *African Journal of Biotechnology*. 2005; 5 (1): 045-048.
- [70] Rouabhi R, Berrebbah H, Djébar M R. Toxic effect of a pesticide, Diflubenzuron on freshwater micro invertebrate (*Tetrahymena pyriformis*). *Chemical Journal Applied Environmental Biology*. 2006; 12 (4): 514- 517.
- [71] Venkateswara J, Srikanth K, Arepalli S K, Gunda V G. Toxic effects of acephate on *Paramecium caudatum* with special emphasis on morphology, behavior and generation time. *Pesticide Biochemistry and physiology*. 2006;(86): 131-137.doi:10.1016/j.pestbp.2006.02005.

- [72] Klauke N, Blanchard M P, Platter H. Polyamine triggering of exocytosis in paramecium involves an extracellular Ca^{2+} (polyvalentcation)- Sensig receptor, SublasmalemmalCa-Store Mobilisation and Store- Operated Ca^{2+} . Influx via unspecific cation channels. The Journal of Membrane Biology. 2000; (15): 141-156.
- [73] Yoshiaki I Wadate. Photolysis of caged calcium in cilia induces ciliary reversal in *Paramecium caudatum*. Journal of Experimental Biology. 2003; (206): 1163-1170.
- [74] Sbartaï I, Berrebbah H, Rouabhi R, Sbartaï H, Guy S, Djebbar M R. Behavior of *Paramecium sp.*, treated with Bifenazole with special emphasis on respiratory metabolism, protein and generation time. American-Eurasian Journal of Toxicological Science. 2009; 1(1): 13-18.
- [75] Vaalavirta L, Tahti A. Astrocyte membrane Na^{+} , K^{+} -ATPase and Mg^{2+} -ATPase as targets of organic solvent impact, Life science. 1995; (57):2223-2230
- [76] Sbartaï I. Toxicité d'un hydrazine (Bifenazte) et d'un oxadiazine (Indoxacarbe) observée chez un modèle cellulaire d'eau douce: *Paramecium sp.* 2013. Ph.D. Thesis, Badji Mokhtar University. Annaba. Algérie. French.
- [77] Bouaricha H. Evaluation du stress oxydatif induit par le Proclaim: Essai comparatif sur deux modèles biologiques (*Helix aspersa* et *Paramecium sp.* 2013. Ph. D. Thesis, Badji Mokhtar University, Annaba. Algérie. French.
- [78] Nzengue Y. Comparaison des mécanismes de toxicité redox du Cadmium, du cuivre et du zinc : place des métallo thionines et de P53. 2008. Ph.D. Thesis, JOSEPH FOURIER – GRENOBLE 1 University, France. French.
- [79] Winston G W, Di Giulio R T. Prooxidant and antioxidant mechanisms in aquatic organisms. Aquatic Toxicology. 1991; (19): 137-161.
- [80] Ojha A, Yaduvanshi S K, Sivastava. Effect of combined exposure of commonly used organophosphate pesticide on lipid peroxidation and antioxidant enzymes in rat tissues; Pesticide Biochemistry and Physiology. 2011; (99). 148-156.
- [81] Leve De L, Kaplowitz N. Glutathione metabolism and its role in hepatotoxicity. Pharmacology Therapeutics. 1991; (52): 287-305.
- [82] Meister A, Anderson M E. Glutathione. Annual Review of Biochemistry. 1983; (52): 711-760.
- [83] Halliwell B, Gutteridge J M C. Glutathione in Metabolism In: Halliwell B, Gutteridge J M C (Eds). In free radicals in biology and medicine. Clarendon Press. Oxford. UK. 1999: 146-150.
- [84] Mosialou E, Ekstrom G, Adang A E, Morgenstern R. Evidence that rat liver microsomal glutathione transferase is responsible for glutathione- dependent protection against lipid peroxidation. Biochemical Pharmacology. 1993; (45): 1645-1651.
- [85] Mansour A S, Mossa A H. Lipid peroxidation and oxidative stress in rat erythrocytes induced by chlorpyrifos and the protective effect of zinc. Pesticide Biochemistry Physiology. 2009;(93): 34-39.
- [86] Charlene C M et al,. Roundup effects on oxidative stress parameters and recovery pattern of *Rhamdia quelen*. archives environmental contamination and toxicology. 2010. Available: <http://dx.doi.org/10.1007/s00244-010-9574-6>.
- [87] Greulich K, Hoque E, Pflugmacher S. Uptake, metabolism, and effects on detoxification enzymes of isoproturon in spawn and tadpoles of amphibians. Environmental Toxicology Safety. 2002; (52): 256-266.
- [88] Feyereisen R. Insect CYP genes and P450 enzymes. In:Gibbert,LI(Ed). Insect Biochemistry and Molecular Biology. Elsevier Academic Press. 2012; 236-316.
- [89] Hemingway J, Hawkes NJ, Maclarioll L, Ranson H. The molecular basis of insect resistance in mosquitoes. Insect biochemistry and Molecular Biology. 2004; (34): 653-665

- [90] Irving H, Riverson JM, Ibrahim SS, Labio NF, Wondji CS. Positional cloning of rp2QTL associates the P450 genes CYP6Z1, CYP6Z3 and CYP6M7 with pyrethroid resistance in the Malaria vector *Anopheles funestus*. Heredity. <http://dx.doi.org/10.1038/hdy.2012.53>
- [91] Wondji et al., Mapping a quantitative trait locus (QTL) conferring pyrethroid resistance in the African Malaria Vector *Anopheles funestus*. BMC Genomics. 2007: 7-34.
- [92] Stevenson et al., Cytochrome P450 6M2 from the Malaria vector *Anopheles gambiae* metabolizes pyrethroid : sequential metabolism of deltamethrin revealed. Insect Biochemistry and Molecular Biology. 2011; (38): 492-502
- [93] Duangkaew et al., Characterization of mosquito CYP6P7 and CYP6AA3 : differences in substrate preference and kinetic properties. Archive Insect Biochemistry and Physiology. 2011;(76). 236-248
- [94] Boonsuepsakul S, Luepronnchai E, Rongnoparut P. Characterization of *Anopheles minimus* CYP6AA3 expressed in a recombinant baculovirus system. Archive Insect Biochemistry and physiology. 2008; (69): 13-21
- [95] Estomih Nkyat T, Akhouayri I, Kisinza, David JP. Impact of environment on mosquito response to pyrethroid insecticides : Facts, evidences and prospects. Insect Biochemistry and Molecular Biology. 2013; (43): 407-416. <http://dx.doi.org/10.1016/J.bmp.2012.10.006>
- [96] Symington SB, Zhang A, Karstens W, Van Houten J, Marshall Clark J. Characterization of Pyrethroid Action on Ciliary Calcium Channel in *Paramecium tetraurelia*. Pesticide Biochemistry and Physiology. 1999;(65): 181-193
- [97] Kadouk A, Matsumura F, Enan E. High affinity binding of 3-verapamil to rat brain synaptic membrane in antagonized by pyrethroid insecticides. Journal of Environmental science and Health. 1994; (29): 855-871