

# Effects of injected N-Methyl-N-Nitrosourea (MNU) in albino mice on the histology and haematology of selected organs of the circulatory, lymphoid and digestive systems

A. A. Osowole<sup>1</sup>, A. A. Oni<sup>2\*</sup>, F. O. Onadje<sup>2</sup>, K.C. Onyegbula<sup>3</sup>, A. T. Hassan<sup>2</sup>

<sup>1</sup>Department of Chemistry, University of Ibadan, Ibadan. Nigeria.

<sup>2</sup>Department of Zoology, University of Ibadan, Ibadan. Nigeria.

<sup>3</sup>Department of Chemical Pathology, College of Medicine, University of Ibadan, Ibadan. Nigeria.

## ABSTRACT

**Aims:** This study was performed to assess the effects of N-methyl-N-nitrosourea (MNU) on the hematology and histology of selected organs in albino mice.

**Study design:** Eighty male and female mice, eight-week old, were subdivided into four groups of twenty each.

**Place and Duration of Study:** Department of Zoology, University of Ibadan between January and July 2011.

**Methodology:** Forty male and female mice received a single dose of 50mg/kg MNU in 1mL of 0.9% NaCl via intra-peritoneal injection. One mouse from each group was sacrificed each month and processed using standard histological techniques. At the end of the study, blood samples collected from two mice per group were processed using standard hematological procedures.

**Results:** MNU induction impacted different morphological abnormalities in induced mice: vacuolization (small intestine), multi-nucleation (spleen, liver), cytoplasmic pallor (heart, liver), increased lymphocytic activity (spleen, thymus), necrosis (heart, liver), and destruction of intestinal wall (small intestine). Increased nuclear-cytoplasmic ratio, nuclear pleomorphism, enlarged nuclei sizes and cytoplasmic degeneration were also common features observed. One of the induced mice developed a neck tumor eight weeks after induction. The histological section of the tumor showed complete necrosis. Except for increased nuclei sizes in the liver, these abnormalities were not observed in control mice. Hematological analysis did not show any significant differences ( $p=0.05$ ) in packed cell volume, hemoglobin, red blood cell count and monocytes between the induced and control mice.

**Conclusion:** Exposure to MNU caused varying degrees of abnormalities over a 28-

\* Tel.: +234-805-530 8669; fax: Nil  
E-mail address: adeyoni2006@yahoo.co.uk

**week period in induced mice.** The study duration however appeared insufficient for the development of any pronounced hematological effects in induced mice.

*Keywords: [Carcinogenesis, Histopathology, Hematology, N-methyl-N-nitrosourea, Swiss Albino mice, Tumors].*

## 1. INTRODUCTION

Chemically induced carcinogenesis models in animals especially rats and mice are widely used for studying the biology of cancer and for developing and evaluating cancer treatment strategies. N-methyl-N-nitrosourea (MNU) is one of the widely used chemicals in several studies of chemical carcinogen models. It has been predominately used in tumor induction and in the investigation of a variety of novel chemo-preventive and treatment agents (Macejova and Brtko, 2001). The works of several authors have established MNU as a potent carcinogen in the induction of diverse tumors on varying strains of rats and mice. Grubbs et al., (1981) examined its effects on hamsters. The induction of mammary tumours in Sprague-Dawley rats has also been well reported (Moon et al., 1992; Rivera et al., 1994; Takahashi et al., 1995 and Thompson et al., 1995). da Silva Franchi et al., (2003) examined its effects on Wistar rats, while its effects on the hematology of Wistar rats were reported by Akanni et al., (2010).

However, there is paucity of information as regards its effects in the Swiss albino mice strain, and thus its susceptibility and suitability in carcinogenesis models involving MNU. Differences in susceptibility may occur across different strains, which is indicative of genetic variation in response (Mori et al, 1985; Festing, 2010). In particular, differences may occur across strains in the target organs that may be affected by MNU. For instance, while MNU induced mammary gland carcinomas in Sprague Dawley rats, (Moon et al, 1992; Rivera et al, 1994; Thompson et al, 1995); thymic lymphomas were observed in Wistar rats and P53+/- mice. Splenic hemangiomas and small intestine adenomas were also reported respectively (da Silva Franchi et al, 2003; Morton et al, 2008). MNU was also found to induce cataracts in 0-20 day old Sprague Dawley rats (Yoshizawa et al, 2000). The time required for the effects to be observed is another important factor. The objective of this two part study (the first of which is reported here), is therefore to provide information on the effects of MNU induction in the Swiss albino mice strain. Furthermore, reports on the **hematology of tumor bearing rats and mice, particularly in Swiss albino mice appear** to be scarce. Hence this study also seeks to provide information in this regard.

Although there have been variations in the dosage of MNU administered, the use of a single dose of 50mg/kg has been used to induce carcinomas by most authors (Rivera et al, 1994; Thompson et al, 1995; Pazos et al, 1998). This dosage is ideal for applications in which rapid tumor induction with high yield is desired (Thompson and Adlakha, 1991). This thus informed the use of this dosage in our study. To achieve the objective stated above, a histological evaluation of selected organs of the circulatory (heart), lymphoid (thymus, spleen) and the digestive systems (liver, and small intestine) in mice given a single intra-peritoneal dose of MNU were carried out. Additional information on the effects of the MNU on some haematological parameters [Packed Cell Volume (PCV), Haemoglobin (Hb), Red Blood Cell (RBC), White Blood Cell (WBC) and its differentials (Lymphocytes, Neutrophils, Monocytes and Eosins), Platelets and Plasma] in the mice was also evaluated.

## 64 2. MATERIAL AND METHODS

65

### 66 2.1 Experimental animals.

67 Eighty (80) eight week old albino mice made up of forty males and forty females were  
68 used for this study. The mice were obtained from the Department of Anatomy, College of  
69 Medicine, University of Ibadan, Ibadan, Nigeria. They were later housed ten each in iron  
70 cages with wooden chips for bedding in a well-ventilated room and maintained on 12 h light /  
71 12 h dark cycle in the animal house of the Department of Zoology. These mice were fed with  
72 standard laboratory diet and had access to water *ad libitum*. They were acclimatized for a  
73 period of two weeks prior to the commencement of the experiment.

74

### 75 2.2 Chemical carcinogen.

76 N-methyl-N-Nitrosourea (MNU Sigma Chemical Co., St. Louise Mo, Catalogue No. 684-  
77 93-5) was purchased from Sigma Aldrich and was used to initiate the carcinogenesis  
78 process.

79

#### 80 2.2.1 Preparation mode for the carcinogen

81 A single dose of 50mg/kg of MNU was administered to the mice via intra-peritoneal  
82 injection. The carcinogen (10mg) was dissolved in 1ml of 0.9 % sodium chloride. Based on  
83 the average weight of 24.47g and 20.14g for males and females respectively, the mice were  
84 thus given a calculated dosage of approximately 0.1ml MNU for both the males and females.

85

### 86 2.3 Experimental Design

87

#### 88 2.3.1 Induction with the carcinogen MNU

89 Mice of both sexes were randomly divided into four groups of twenty each (MI, FI, MC and  
90 FC). Groups MC (male control) and FC (female control) served as the control (members of  
91 this group were not induced nor given placebo) while groups MI, (male induced) and FI  
92 (female induced) received 0.1ml of MNU via intra-peritoneal injection. After the carcinogen  
93 induction, the mice were kept on basal diet and water *ad libitum* and were subsequently  
94 weighed weekly to monitor changes in their weight. From the second week after induction,  
95 animals were palpated twice weekly to monitor possible tumor appearance. Complete gross  
96 examination was carried out for detection of tumor. The animals were monitored for 28  
97 weeks. One mouse from each of the four groups was sacrificed for histological examination  
98 at the end of every four weeks. This was to allow for sufficient sample size for the second  
99 phase of this study. Although less common, studies involving the use of one mouse per time  
100 point/group for histological evaluation in cancer studies do occur in literature (Lu and  
101 Meistrich, 1979; Barres et al, 2010). In all, a total of twenty eight mice comprising seven  
102 mice from each of the four groups were sacrificed over the 28-week study period and  
103 processed for histological examination. In addition, a tumor bearing mouse which died

before the end of the study period was also processed for histological examination, bringing the total number of mice examined to twenty nine. The mice were anaesthetized with chloroform (except for the dead mouse) and subsequently dissected to obtain the selected organs: liver, spleen, thymus, small intestine and heart, which were then stored in 40% formal saline prior to histological analysis. Histological analysis was carried out using the following equipment: a Leica rotary microtome model 2125 RT, a Leica tissue embedding machine model EG 1160, a Raymond Lamb tissue floatation bath model E652 and hot plate model E18, as well as a Tissue-Tek II tissue processor model 4634. The prepared slides were viewed using an Olympus CX31 microscope with computer attachment. Results of representative slides are however presented in this paper. Except for the sacrificed animals, all other mice were monitored weekly for changes in their body weight, hence total number of mice in each of the four groups (n) varied from 20 at the beginning of the study period, to 13 mice at week 28 respectively. The male induced group (MI) however contained 12 mice instead of 13, due to mortality in the tumor bearing mouse. The remaining mice were used for the second part of this study (not reported here).

### **2.3.2 Collection of blood samples and measurement of hematological parameters**

At the end of the 28-week study period, two mice from each of the four groups (eight in all) were sacrificed out of the fifty one mice remaining. This was to provide some preliminary information on the probable effects of the carcinogen on hematological parameters. However, only two mice per group were sacrificed for this purpose. This was to allow for an adequate sample size for the second stage of this study which is currently in progress. Blood samples were collected from the retro-orbital sinus of the mice with heparinised 70ml micro-haematocrit capillary tubes into a vial containing 0.5ml Ethylene Diamine Tetra Acid (EDTA). White blood cell (WBC) counts, RBC counts, PCV, platelets and hemoglobin (Hb) content of the blood obtained from the mice were analyzed at the hematology unit of the Department Of Veterinary Medicine, University Of Ibadan, Nigeria using the method of Moore, (1958) and Stevens, (1997).

### **2.3.3 Statistical Analysis.**


T-Test was used to determine if there were any statistically significant differences (at  $p = 0.05$ ) between the means of the control and test group for each of the hematological indices determined, as well as for the body weight. Sexes were analysed separately.

## **3. RESULTS**

### **3.1 Body Weights and Clinical Signs**

Members of both groups gained weight progressively during the study period (Figure 1), with the induced and control males reaching a maximum weight of 40.03g and 40.37g respectively, at the end of the study period. The female control group presented a mean body weight ( $\pm$  SD) of  $28.48g \pm 0.95$ , which was higher than the mean body weight of the female induced group ( $27.24g \pm 0.73$ ). However this difference was not significant at  $p = 0.05$ . Furthermore, the average weight of the induced male mice ( $34.38g \pm 0.94$ ) was higher

149 than the male control mice ( $34.23\text{g} \pm 1.08$ ) by a difference of  $0.15\text{g}$ , which was also not  
 150 statistically significant at  $p = 0.05$ . The induction with MNU appears not to have caused any  
 151 changes in the behavior of both the control and the induced mice. Majority of the induced  
 152 male and female mice appeared to be healthy and showed no observable behavioral  
 153 differences from their respective control groups. The induced mice also did not show any  
 154 form of labored or rapid breathing patterns. One of the male induced mice was suspected to  
 155 have developed a tumor in the cervical region barely eight weeks after the carcinogen  
 156 induction. Except for the mouse with tumor, no mortality was observed in the induced or  
 157 control mice.



158

159

160 Fig. 1. The average weight of both the induced and control albino mice over a 28-week study  
 161 period.

162 *Mean  $\pm$  S.D = Mean values  $\pm$  Standard deviation of twenty mice per group.*

163 *However there was variation in N over the study period due to the exclusion of animals sacrificed for histology at the*  
 164 *end of every four weeks (seven in each group over the study period). One mouse also died in the group MI before*  
 165 *the end of the study period.*

166

### 167 3.2 Histopathology.


168 There were varying degrees of lesions and abnormalities in the induced animals when  
 169 compared to the control. Abnormalities were observed in all the seven induced male and  
 170 female mice respectively, which were sacrificed over the study period. Except for increased  
 171 nuclei sizes in the liver, all the control mice sacrificed (seven in each group) did not record  
 172 any abnormalities or lesions. Representative figures are however used here for the  
 173 comparison. The figures presented represent some of the abnormalities observed in the  
 174 organs of different mice examined during the study period. Figure 2 shows the histological  
 175 section of the thymus of a female control mouse with normal architecture and well delineated  
 176 cells. Vesicular nuclei, nuclear pleomorphism, increased nucleo-cytoplasmic ratio and  
 177 increased lymphocytic activity (Figure 3) were some abnormalities observed in the thymus of  
 178 the induced mice. While normal architecture with well delineated striated muscles was

179 observed in the heart of the control mice (Figure 4), the morphological changes observed in  
180 the heart of the induced mice included necrosis and scanty lymphocyte infiltration (Figure 5).

181 The spleen of the control showed normal architecture (Figure 6), while some of the  
182 abnormalities observed in the induced mice were cytoplasmic degeneration, increased  
183 lymphocytic activity and nuclear pleomorphism (Figure 7). The liver of the MNU treated mice  
184 showed abnormalities such as necrosis and cytoplasmic degeneration associated with  
185 cytoplasmic pallor (Figure 9). There were also dilation of the sinusoids, and disintegration of  
186 hepatic cells including necrosis of hepatocytes, and enlarged nuclei in the liver of the treated  
187 mice.


188 Other observable features were; serious disarrangement and degeneration / distortion of  
189 the regular hexagonal shape of the hepatocytes. Furthermore, most of the cytoplasms have  
190 lost their eosinophilic colours, the radial architecture of the cells has also been altered and  
191 the cells thus lack distinct outline and appear reduced in size indicating distortion of the  
192 hepatic architecture. Some of these features were however, observed in the histological  
193 sections of the control mice (Figure 8). The histological sections of the small intestine of the  
194 control mice showed well defined architecture with unaltered intestinal wall (Figure 10).  
195 However, degeneration of the intestinal wall of the small intestine was observed as  
196 illustrated in Figure 11. The histological section of the cervical tumor observed in one of the  
197 exposed mice barely eight weeks after exposure showed complete necrosis (Figure 13).

198




199


200 Fig.2. Control Thymus Female showing normal architecture (Mag. x 400).



201  
 202 Fig.3. Induced Thymus Female showing Vesicular Nuclei (VN), irregularly shaped nuclei or  
 203 Nuclear Pleomorphism (NP), increased Nucleo-Cytoplasmic Ratio (NCR) and increased  
 204 lymphocytic activity (L) (dark colored cells) (Mag. x 400).




205  
 206 Fig.4. Heart Control Male showing normal architecture with well delineated striated muscle  
 207 and evenly distributed cardiomyocytes (Mag. x 400).




208

209 Fig.5. Heart Induced Male Showing Necrosis and scanty lymphocytes (Mag. x 400).




210

211 Fig.6. Spleen Control Male showing normal architecture (Mag. x 400)




212

213 Fig.7. Induced Spleen Male showing Cytoplasmic Degeneration (CD), increased  
214 Lymphocytic activity (L) and Nuclear-Pleomorphism (NP) (Mag. x 400).




215

216 Fig.8. Histological section showing normal appearance of female liver (Mag. x 400)




217

218 Fig.9. Histological section of the liver of an induced female showing necrosis and  
 219 cytoplasmic degeneration associated with cytoplasmic pallor (Mag. x 400).




220

221 Fig.10. Histological section of the small intestine of male control mice showing normal  
 222 architecture and unaltered intestinal wall (Mag. x 100).




223

224 Fig.11. Histological section of the small intestine of an induced male showing corrosion of  
225 the intestinal wall (C) (Mag. x 100).



226

227 Fig.12. Suspected tumor in the cervical region in Male mouse induced with MNU.



228

229 Fig.13. Histological section of the above tumor showing necrosis (Mag. X 400).

230

### 231 3.3 Hematological Analysis

232

233 The hematological parameters that were considered in this study are shown in Table 1. The  
234 analysis shows that there were no significant differences ( $P = 0.05$ ) amongst all the  
235 parameters considered in this study between the control and induced mice. Generally,  
236 eosinophils showed higher values in females ( $0.75 \pm 0.35$ ;  $0.75 \pm 0.35$ ) than males ( $0.5 \pm$   
237  $0.71$ ;  $0.5 \pm 0.0$ ), although levels within the induced and control in both sexes respectively  
238 were the same (Table 1). White blood cell count, lymphocytes and plasma levels were not

239 significantly different at  $p = 0.05$  in both induced and control mice (both sexes). Similarly,  
240 packed cell volume, hemoglobin content, red blood cells, neutrophils and monocytes showed  
241 no significant differences at  $p = 0.05$  (Table 1).  
242  
243  
244  
245  
246  
247  
248  
249  
250

251 **Table 1: Hematological Parameters of MNU induced and control Albino Mice**

252

| Test samples   | PCV (%)   | Hb (g/dl)  | RBC (L)   | WBC X10 <sup>3</sup> (μL) | Platelets X10 <sup>4</sup> (μL) | Lymphocytes | Neutrophils | Monocytes | Eosinophils | Plasma    |
|----------------|-----------|------------|-----------|---------------------------|---------------------------------|-------------|-------------|-----------|-------------|-----------|
| Fc             | 32        | 10.35      | 5.32      | 14.93                     | 25.35                           | 67          | 30.5        | 2         | 0.5         | 7.9       |
|                | 39.5      | 12.95      | 6.4       | 5.95                      | 11.6                            | 67          | 29          | 3         | 1           | 7.5       |
| Means±SD       | 35.75±5.3 | 11.65±1.84 | 5.86±0.76 | 10.44±6.35                | 18.48±9.72                      | 67±0        | 29.75±1.06  | 2.5±0.71  | 0.75±0.35   | 7.7±0.28  |
| F <sub>i</sub> | 37        | 12.15      | 6.15      | 13.25                     | 23.05                           | 60          | 39          | 1         | 0.5         | 6.75      |
|                | 39        | 12.80      | 6.35      | 4.43                      | 8.7                             | 68          | 29          | 2         | 1           | 7.55      |
| Means±SD       | 38±1.41   | 12.48±0.46 | 6.25±0.14 | 8.84±6.24                 | 15.88±10.3                      | 63.75±6     | 34±7.07     | 1.5±0.71  | 0.75±0.35   | 7.15±0.57 |
| Mc             | 31        | 10.05      | 4.89      | 7.75                      | 13.8                            | 67          | 31.5        | 0.5       | 1           | 7.7       |
|                | 38        | 12.60      | 6.4       | 7.4                       | 9.2                             | 70          | 29          | 1         | 0           | 7.5       |
| Means±SD       | 34.5±4.95 | 11.33±1.8  | 5.65±1.07 | 7.58±0.25                 | 11.5±325.3                      | 68.5±2.12   | 30.25±1.77  | 0.75±0.35 | 0.5±0.71    | 7.6±0.14  |
| M <sub>i</sub> | 31        | 10.15      | 5.21      | 11.38                     | 15.9                            | 67          | 32          | 0.5       | 0.5         | 7.6       |
|                | 39        | 13         | 6.35      | 2.98                      | 6.3                             | 69          | 28          | 2.5       | 0.5         | 7.5       |
| Means±SD       | 35±5.66   | 11.56±2.00 | 5.78±0.81 | 7.18±187.8                | 11.1±2146.6                     | 68±1.41     | 30±2.83     | 1.5±1.41  | 0.5±0.0     | 7.55±0.07 |

253

254

255

256 *F<sub>c</sub>*-Female control; *F<sub>i</sub>*-Female induced; *Mc*-Male control and *M<sub>i</sub>*- Male induced; *PCV*- Packed Cell Volume; *Hb*- Hemoglobin; *RBC* - Red Blood Cell and *WBC*- White Blood  
 257 Cell. (n=2)

258

259

260

#### 4. DISCUSSION

MNU is a very potent and reliable carcinogen with a severe toxicity in terms of the duration taken to induce cancer and bring about effects in organs of organisms, when compared to other known carcinogens (Franchi *et al.*, 2003). It was not clear why the male induced mice presented a slightly higher mean body weight when compared to their male control counterparts, although these differences were not significant at  $p = 0.05$ . However, though rarely observed, similar findings occur in literature. Swiss Albino mice inoculated with Ehrlich Ascites Carcinoma had a maximum body weight gain of 20% compared to Cisplatin treated controls. No reasons were however given for this observation by Sathistha *et al.*, (2008). However it may probably indicate some degree of resistance by this strain of mice to the effects of the carcinogen induction. When organisms are exposed to toxicants, their organs tend to accumulate such toxicants (Pelgrom *et al.*, 1995). The organism may metabolize the toxicant and excrete it. However, it is the accumulation of the toxicant in the organs of the induced mice, such as the liver, spleen and heart amongst others that is responsible for the abnormalities observed in these visceral organs as shown in this study.

The thymus is a specialized organ of the immune system. The only known function is the production and "education of T-lymphocytes (T cells) which are critical cells of the adaptive immune system. It is derived from the pharyngeal pouch and in relationship with other lymphatic organs takes a central super ordinate place (da Silva Franchi *et al.*, 2003). Induction with MNU in P53+/- mice resulted in the effacement of the thymic cortico-medullary architecture by diffuse sheets of lymphoblasts with large euchromatic nuclei, moderate to high number of mitotic figures, infiltration of lymphoblasts through the thymic capsule and presence of cluster or sheets of lymphoblasts in the spleen, liver and other visceral organs (Morton *et al.*, 2008). Increased lymphocytic activity was observed in the thymus of the female induced and the spleen of male induced mice respectively. Morton *et al.*, (2008) also observed enlarged nuclei sizes in the thymus of the induced mice. Both studies also reported the presence of thymic lymphomas in the strains used. Enlarged nuclei sizes and other abnormalities such as pleomorphic nuclei were also observed in the thymus in this study. These cellular abnormalities may therefore suggest a gradual progression towards the distortion reported by the above authors.

The heart and spleen of the induced mice also showed a number of abnormal features; an indication of distortion and degeneration as a result of exposure to the carcinogen. Although, reports of the effects of injected MNU on the histology of the heart seem very limited, some of the distortions have however been recorded by Palazzi and Kergozien-Framery (2009). These authors also reported other degenerations in the heart which included myofibrosis, distortion of muscle fibres and mineralization. Minardi and Maltoni (1998) reported splenic hemosiderosis, epitheliomorphous peritoneal mesothelioma and biphasic (solid epithelial and fusocellular pattern) as features of abnormalities in the spleen of an MNU induced rat. However, there were no histopathological changes recorded in the heart and spleen of rats exposed to MNU intravenously in a six month study conducted by Murata *et al.*, (2006).

The liver is an important organ of metabolism, detoxification, storage and excretion of xenobiotics and their metabolites and is especially vulnerable to damage. Its activeness makes it more susceptible to carcinogen intoxication (Fazeli *et al.*, 2006). It is the primary target for carcinogen effect of more than two hundred chemicals (including pesticides, food additives, pharmaceuticals and industrial intermediates) tested in long-term toxicity safety assessment assays (Ward *et al.*, 1996). Most tumor initiating agents either generate or are metabolically converted to electrophilic reactants which bind covalently to cellular DNA (Slaga *et al.*, 1986; DiGiovannio, 1992). Free radicals and these modified DNA bases have been strongly implicated in carcinogenesis in general (Floyd, 1990; Malins, 1993). A high correlation has been found to occur between the dose of administered carcinogens such as MNU and the levels of total DNA adducts in the liver. The liver is hypothesized to have a

greater capacity for metabolizing toxicants (Izzotti *et al.*, 1999). MNU is a well known liver carcinogen which induces multiple cellular, molecular, and biochemical changes (Samaras *et al.*, 2010).

This study showed that MNU induction resulted in extensive degenerative changes in the liver varying from necrosis, multinucleation and increased nucleo-cytoplasmic ratio, cytoplasmic degeneration and pallor (Figures 11 & 12). Hyperplasia of the hepatic cells, congestion of central vein, and dilation of sinusoids were also observed in most of the histological sections of the induced mice. Similar observations have been reported by Shashi and Thapar (2000) in albino rabbits. The results of the histopathology of the liver were very consistent with early reports by Yamamoto *et al.*, (2002) in F344/DuCrj rats and Crj: BDF1 mice; and Brzoska *et al.*, (2003) in male Wistar rats. In rabbits, Wang *et al.*, (1993), also reported necrotic areas in the liver cells associated with vacuolar changes and proliferation of connective tissue fibres. Some of these observations have also been reported in the liver of rats exposed to formaldehyde (Bandman, 1994; and Teng *et al.*, 2001). The increased sizes of the nuclei noticeable in both groups can be attributed to the increased activity going on in the cells since the liver is a very active metabolizing organ.

MNU exposure to the small intestine of albino mice caused an ulcerating effect as the walls of the small intestine (epithelial) were indented and eroded. Other morphological features noticed were nuclear pleomorphism, cytoplasmic pallor and vacuolization. Although there is paucity of reports showing the effects of injected MNU on the small intestine of albino mice, the morphological changes observed did not differ significantly from the abnormalities found in the other organs examined. Yoshizawa *et al.*, (2000) observed vacuolation in the posterior cortex of the lens of neo-natal Sprague Dawley rats, as cataract development progressed. One male mouse was suspected to have developed a tumor in the cervical region at 60 days post induction. The tumor appeared to be encapsulated within a thin sheet of fibrous connective tissue. Further studies are however required to confirm the tumor and its origin. The histo-pathological examination of the tumor showed complete necrosis. This result is very similar to the findings of several authors on tumors observed in different strains of rats and rabbits. (Shashi and Thapar, 2000 on albino rabbit; Brzoska *et al.*, 2003 and da Silva Franchi *et al.*, 2003 on Wistar rats and Morton *et al.*, 2008 on P53+/- mice)..

In comparison with other studies, Rivera *et al.*, (1994) administered MNU to female Sprague Dawley rats at a dosage of 50 mg/kg in animals of age 50, 80 and 110 days old. About 90% developed mammary tumors. Moon *et al.*, (1992) administered a single intravenous dose of 50 mg/kg in 120 day old Sprague Dawley rats. Their results showed a cancer incidence of 75-95% which were mainly mammary carcinomas. da Silva Franchi *et al.*, (2003) varied the dosage of MNU (intra-peritoneal) between 80-240 mg/kg in eight week old Wistar rats and found thymic lymphomas and splenic hemangiomas in the intermediate to high dose category (160-240 mg/kg). Morton *et al.*, (2008) also obtained thymic lymphomas in P53+/- mice. MNU resulted in cataract formation in 0-20 day old Sprague Dawley rats 7-30 days after exposure to a single intraperitoneal injection of 100 mg/kg (Yoshizawa *et al.*, 2000). In contrast, our study on MNU induction in albino mice using a single intra-peritoneal dose of 50 mg/kg showed that percentage tumor induction was minimal at the specified dose. This suggests that higher doses and/or a longer time frame may be required for carcinomas to develop in this strain of mice. It may also indicate that this strain is less susceptible than other strains such as Sprague Dawley and Wistar rats and may also likely explain the frequent use of the above strains in carcinogenesis models.

There were no significant differences in hematological parameters between the control and induced mice ( $p = 0.05$ ). These results were then compared with the normal range of values expected in mice as stated by the Research Animal Resources of the University of Minnesota and the Iowa State University (Retrieved 17<sup>th</sup> February, 2012 from [www.ahc.umn.edu/rar/refvalues.html](http://www.ahc.umn.edu/rar/refvalues.html) and [www.lar.iastate.edu/index.php?option=com\\_content&](http://www.lar.iastate.edu/index.php?option=com_content&)). The normal range for hematological

parameters in mice as given by the above were 10.2 – 16.6 g/dl for hemoglobin, white blood cell count (6-15 x 1000), lymphocytes, (55-95%), monocytes (1-4), platelets (160-410 x 1000) and eosinophils (0-4). Normal ranges for PCV, neutrophils and plasma were however not stated. Except for platelets in both male induced and control, and eosinophils in male controls, the range of hematological values observed in the control and induced animals of both sexes were mainly within the range of normal values stated above. However it is also pertinent to note that the ranges given above are not firm boundaries as age, sex, strain, sampling techniques and methodology may account for some degree of variation [www.ahc.umn.edu/rar/refvalues.html](http://www.ahc.umn.edu/rar/refvalues.html).

The results of the hematological analysis were then compared with studies by other authors (Chakraborty *et al*, 2010; Raman *et al*, 2010) who observed significant differences in some hematological parameters between control and tumor bearing mice. Hemoglobin for instance, in the studies by Raman *et al*, (2010) had values of  $12.92 \pm 0.15$  in control mice compared to  $5.78 \pm 0.4$  g/dl in tumor bearing mice. Lymphocytes were  $66.4 \pm 1.36\%$  in control compared to tumor bearing mice with  $26.2 \pm 1.16\%$ , while PCV values in both control and tumor bearing mice were much lower than obtained in this study at 16.86 and 25.98%. Values for PCV in studies by Akanni *et al*, (2010) at 24% and 31% for tumor bearing and control mice respectively were also lower than obtained in our study. Studies by Chakraborty *et al*, (2010) showed a similar pattern as that of Raman *et al*, (2010). Lymphocytes in normal unexposed mice were  $67.13 \pm 1.41$  compared to tumor bearing mice at  $18.54 \pm 0.28\%$ , while red blood cell count in control and tumor bearing mice in studies of Sathistha *et al*, (2008) were  $5.03 \pm 0.28$  and  $2.14 \pm 0.34$  respectively.

The disparity between the hematological results in this study and that presented by the above authors may therefore be due to the fact that these authors examined tumor-bearing rats; while in this study, only one of the induced mice had advanced to the tumor-bearing stage (figures 12 and 13) during the study period. It is hypothesized that the duration of this study was probably not sufficient to have resulted in significant differences among the hematological parameters between the induced and the control animals when compared with some previous studies. The fact that the hematological values were mainly within normal range in both induced and controls' further supports this assertion. It is also illustrative of the challenges in human cancer diagnosis and treatment, in which symptoms often appear when the disease is at an advanced stage, thus making treatment of the disease challenging.

#### 4. CONCLUSION

Reports abound on MNU induction in Sprague Dawley rats, less commonly in Hamsters, and Wistar rats. Our studies provide information on MNU induction in albino mice, which is illustrative of differences in the susceptibility of various strains to the carcinogen induction. Percentage tumor induction was minimal at a dose of 50 mg/kg, with only one mouse suspected to have developed a tumor in the cervical region at about 60 days post induction. No acute toxicity was also observed except for the mouse with tumor.

410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454

**ACKNOWLEDGEMENTS**

One of us, Aderoju Osowole thanks TWAS-COMSTECH for the research grant 09-137 RG/PHA/AF/AC\_C-UNESCO FR: 3240231226 used to purchase the chemicals for the study; and TWAS-DFG for a fellowship to Germany to undergo training in anti-cancer studies.

**COMPETING INTERESTS**

Authors have declared that no competing interests exist.

**AUTHORS' CONTRIBUTIONS**

Author A initiated the study and provided funds via the research grant. Author B managed the experimental, draft writing, and corrections of the manuscript, Author C carried out the experimental work and prepared first draft, Author D carried out the histology, while Author E conceptualized and supervised the study. All authors read and approved the final manuscript."

**ETHICAL APPROVAL**

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable.

**REFERENCES**

Akanni, E.O, Oloke, J.K., Fakunle, E.E. (2010). Haematological characterization in N-nitroso-N-ethylurea induced tumour bearing rats on oral administration of pleurotus pulmonarius and pleurotus ostreatus metabolites. IJRRAS, 3, 343-348.

Bandman, A. (1994). Hazardous substances, galogen and oxygen containing substances. Chimia, 48, 124-128.

Barres V., Ouellet, V., Lafontaine J., Tonin, P.N., Provencher D.M., Mes-Messon, A.M. An essential role for Ran GTPase in epithelial ovarian cancer cell survival. Molecular Cancer 9, 272.

Brzoska, M.M., Moniuszko-Jakoniuk, J., Pilat-Marcinkiewicz, B., Sawicki, B. (2003). Liver and kidney function and histology in rats exposed to cadmium and ethanol. Alcohol Alcohol, 38, 2-10.

Chakraborty, A., Kumar, P., Ghosh, K., Roy, P. (2010). Evaluation of a Schiff base copper complex compound as potent anticancer molecule with multiple targets of action. Eur. J. Pharmacol, 647, 1-12.

455 da Silva Franchi, C.A., Bacchi, M.M., Padovani, C.R., de Camargo, J.L. (2003). Thymic  
 456 lymphomas in Wistar rats exposed to N-methyl-N-nitrosourea (MNU). *Cancer Sci*,  
 457 94, 240-243.

458 DiGiovanni, J. (1992). Multistage carcinogenesis in mouse skin. *Pharmacol Ther*, 54, 63-  
 459 128.

460 Fazeli, S.A, Davarian, A., Azarhoush, R., Golalipour, M.J. (2006). Histopathologic changes  
 461 of rat liver following formaldehyde exposure. *Pak. J. Biol. Sci*, 9, 2137-2140.

462 Festing, M.F.W. (2010). In bred strains should replace out bred stocks in toxicology, safety  
 463 testing and drug development. *Toxicol. Pathol*, 38, 681-690.

464 Floyd, R.A. (1990). Role of oxygen free radicals in carcinogenesis and brain ischemia. *Fed*  
 465 *Am. Soc. Exp. Biol. J*, 4, 2587-2597.

466 Grubbs, C.J., Becci, P.J., Thompson, H.J., Moon, R.C. (1981). Carcinogenicity of N-methyl-  
 467 N-nitrosourea and N-ethyl-N-nitrosourea when applied to a localized area of the  
 468 hamster trachea. *J. Natl. Cancer Inst*, 66, 961-965.

469 Iowa State University. Vice President for Research and Development. Normal values in the  
 470 mouse. Hematology. Retrieved 17<sup>th</sup> February, 2012 from  
 471 [www.lar.iastate.edu/index.php?option=com\\_content&](http://www.lar.iastate.edu/index.php?option=com_content&)

472 Izzotti, A., Camoirano, A., Cartiglia, C., Grubbs, C.J., Lubet, R.A., Kelloff, G.J., De Flora, S.  
 473 (1999). Patterns of DNA adduct formation in liver and mammary epithelial cells in  
 474 rats treated with 7,12-Dimethyl-benz(a)anthracene, and selective effects of  
 475 chemopreventive agents. *Cancer Res*, 59, 4285-4290.

476 Lu, C.C, Meistrich, M.L. (1979). Cytotoxic effects of chemotherapeutic drugs on mouse  
 477 testis cells. *Cancer Res*, 39, 3575-3582.

478 Macejova, D., Brtko, J. (2001). Chemically induced carcinogenesis: a comparison of 1-  
 479 Methyl-1-Nitosourea, 7, 12- DiethylBenzanthracene, Diethylnitroso-amine and  
 480 Azoxymethan Models (Minireview). *Endocrine Regulations*, 35, 53-59.

481 Minardi, F., Maltoni, C. (1998). Results of long-term carcinogenicity bioassays of ceramic  
 482 fibres ("Fiberfrax") on Sprague-Dawley rats. *Eur. J. Oncol*, 3, 241-249.

483 Malins, D.C. (1993). Identification of hydroxyl radical lesions in DNA base structure:  
 484 biomarkers with a putative link to cancer developments. *J. Toxicol Environ Health*,  
 485 40, 247-261.

486 Moon, R.C., Kelloff, G.F., Detrisac, C.J., Steele, V.E., Thomas, C.F., Sigman, C.C. (1992).  
 487 Chemoprevention of MNU-induced mammary tumours in the mature rat by 4-HPR  
 488 and tamoxifen. *Anticancer Res*, 12, 1147-1153.

489 Moore, R. (1958). The white blood cell count in the indigenous people of East Africa.  
 490 *J.Trop.Med.Hyg*, 61, 70-72.

491 Mori, S., Takeuchi, Y., Toyama, M., Makino, S., Ohhara, T., Tochino, Y., Hayashi, Y. (1985).  
 492 Amitrole: Strain differences in morphological response of the liver following  
 493 subchronic administration to mice. *Toxicol. Lett*, 29 (2-3), 145-152.  
 494

495 Morton, D. Bailey, K.L., Stout, C.L, Weaver, R.J, White, K.A., Lorenzen, M.J., Ball, D.J.  
 496 (2008). N-Methyl-N-Nitrosourea (MNU): A positive control chemical for p53+/-  
 497 mouse carcinogenicity studies. *Toxicol. Pathol*, 36, 926-931.

498 Murata, S., Kominsky, S.L., Vali, M., Zhang, Z., Garrett-Mayer, E., Korz, D., Huso, D., Baker,  
 499 S.D., Barber, J., Jaffee, E., Reilly, R.T., Sukumar, S. (2006). Ductal Access for  
 500 Prevention and Therapy of Mammary Tumors. *Cancer Res*, 66, 638-645.

501 Palazzi, X., Kergozien-Framery, S. (2009). Use of rasH2 transgenic mice for carcinogenesis  
 502 testing of medical implants. *Exp. Toxicol. Pathol*, 61, 433-441.

503 Pazos, P., Lanari, C., Elizalde, P., Montecchia, F., Charreau, E.H., Molinolo, A.A. (1998).  
 504 Promoter effect of Medroxyprogesterone Acetate (MPA) in N-methyl-N-nitrosourea  
 505 (MNU) induced mammary tumors in BALB/c mice. *Carcinogenesis* 19, 529-531.

506 Pelgrom, S., Lock, R., Balm, P., Wendelaar Bonga, S. (1995). Integrated physiological  
 507 response of tilapia, *Oreochromis mossambicus* to sublethal copper exposure.  
 508 *Aquat. Toxicol*, 32, 303-320.

509 Raman, N., Jeyamurugan, R., Senthilkumar, R., Raj Kapoor, B., Franzblau, S.G. (2010). In  
 510 vivo and in vitro evaluation of highly specific thiolate carrier group (II) and Zinc (II)  
 511 complexes on Ehrlich ascites carcinoma tumor model. *Eur. J. Med. Chem*, 45,  
 512 5438-5451.

513 Rivera, E.S., Andrade, N., Martin, G., Melito, G., Cricco, G., Mohamad, N., Davio, C., Caro,  
 514 R., Bergoc, R.M. (1994). Induction of mammary tumors in rat by intraperitoneal  
 515 injection of MNU: histopathology and estral cycle influence. *Cancer Lett*, 86, 223-  
 516 228.

517 Samaras, V., Rafailidis, P.I., Mourtzoukou, E.G., Peppas, G., Falagas, M.E. (2010). Chronic  
 518 bacterial and parasitic infections and cancer: a review. *J Infect Dev Ctries*, 4, 267-  
 519 281.

520 Sathistha, M.P., Shetti, U.N., Revankar, V.K., Pai, K.S.R. (2008). Synthesis and antitumor  
 521 studies on novel Co(II), Ni(II) and Cu(II) metal complexes of bis(3-acetylcoumarin)  
 522 thiocarbohydrazone. *Eur. J. Med. Chem*, 43, 2338-2346.

523 Shashi, A., Thapar, S.P. (2000). Histopathology of fluoride-induced hepatotoxicity in rabbits.  
 524 *Fluoride*, 34, 34-42.

Slaga, T.J., O'Connell, J., Rostein, J., Patskan, G., Morris, R., Aldaz, C.M., and Conti, C.J. (1986). Critical genetic determinants and molecular events in multistage skin carcinogenesis. *Symp. Fundam. Cancer Res*, 39, 31-44.

Stevens, M.L. (1997). *Fundamentals of Clinical Hematology*. W.B Saunders. Philadelphia, PA. in: Banaee, M. Mirvagefi, A.R., Rafei, G.R. and Majazi Amiri, B. (2008). Effect of sublethal diazinon concentration on blood plasma biochemistry. *Int. J. Environ. Res*, 2, 189-198.

Takahashi, H., Uemura, Y., Nakao, I., Tsubura, A. (1995). Induction of mammary carcinomas by the direct application of crystalline N-methyl-N-nitrosourea onto rat mammary gland. *Cancer Lett*, 92, 105-111.

Teng, S., Beard, K., Pourahmad, J., Morideni, M., Easson, E., Poon, R., O'Brien, P.J. (2001). The formaldehyde metabolic detoxification enzyme systems and molecular cytotoxic mechanism in isolated rat hepatocytes. *Chem-Biol. Interact*, 130-132, 285-296.

Thompson, H.J. and Adlakha, H. (1991). Dose-responsive induction of mammary gland carcinomas by the intra-peritoneal injection of 1-Methy-1-nitrosourea. *Cancer Res*, 51, 3411-3415.

Thompson, H.J, McGinley, J.N., Rothhammer, K., Singh, M. (1995). Rapid induction of mammary intraductal proliferations, ductal carcinoma in situ and carcinomas by the injection of sexually immature female rats with 1-methyl-1-nitrosourea. *Carcinogenesis*, 16, 2407-2411.

University of Minnesota, Research Animal Resources. Reference values for laboratory animals. Normal Hematology Values. Retrieved 17<sup>th</sup> February, 2012 from [www.ahc.umn.edu/rar/refvalues.html](http://www.ahc.umn.edu/rar/refvalues.html)

Wang J, Zheng, Z.A., Zhang, L.S., Cao, D.M., Chen, K.Z, Lu, D. (1993). An experimental study for early diagnostic features in fluorosis. *Fluoride*, 26, 61-65.

Wilbur, W.M. (2009). *The World of the Cell*, seventh ed. Beaker, San Francisco.

Ward, J.M., Shibata, M.A., Devor, D.E. (1996). Emerging issues in mouse liver carcinogenesis. *Toxicol. Pathol*, 24, 129-137.

Yamamoto, S., Kasai, T., Matsumoto, M., Nishizawa, T., Arito, H., Nagano, K., Matsushima, T. (2002). Carcinogenicity and chronic toxicity in rats and mice exposed to chloroform by inhalation. *J. Occup. Health*, 44, 288-293.

Yoshizawa, K., Oishi, Y., Nambu, H., Yamamoto, D., Yang, J., Senzaki, H., Miki, H. and Tsubura, A. (2000). Cataractogenesis in Neonatal Sprague Dawley rats by N-Methyl-N-nitrosourea. *Toxicol. Pathol*, 28, 555-564.

561  
562  
563  
564  
565  
566  
567